

UNCLASSIFIED

US ARMY
PUBLIC HEALTH COMMAND
(Provisional)



WIPE SAMPLE INTERPRETATION

E2S2 Conference
Denver, 14-17 June 2010

George Murnyak, CIH
USAPHC

Environmental Health Risk Assessment Program



UNCLASSIFIED

Report Documentation Page

Form Approved
OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE JUN 2010	2. REPORT TYPE	3. DATES COVERED 00-00-2010 to 00-00-2010			
4. TITLE AND SUBTITLE Wipe Sample Interpretation		5a. CONTRACT NUMBER			
		5b. GRANT NUMBER			
		5c. PROGRAM ELEMENT NUMBER			
6. AUTHOR(S)		5d. PROJECT NUMBER			
		5e. TASK NUMBER			
		5f. WORK UNIT NUMBER			
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. Army Public Health Command ,5158 Blackhawk Road ,Aberdeen Proving Ground,MD,21010-5403		8. PERFORMING ORGANIZATION REPORT NUMBER			
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)		10. SPONSOR/MONITOR'S ACRONYM(S)			
		11. SPONSOR/MONITOR'S REPORT NUMBER(S)			
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES Presented at the NDIA Environment, Energy Security & Sustainability (E2S2) Symposium & Exhibition held 14-17 June 2010 in Denver, CO.					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 26	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Purpose

UNCLASSIFIED



- Describe the rationale and logic used to assess health risks associated with chemical contamination on indoor surfaces
- USAPHC developed TG 312 for *office environments*
- Will not address issues related to collecting a “good” wipe sample



UNCLASSIFIED

Background and Scope

UNCLASSIFIED



- Few published health-based wipe sample standard/guidelines
- The development of TG 312 was an evolutionary process over time
- Guide is written in two parts;
 - First part basic concepts/explanation for general preventive medicine community
 - Second part detailed discussion of methodology for health risk assessors



UNCLASSIFIED

UNCLASSIFIED Evolution of Technical Guide 312



- Pesticide residues at military housing
- Johnston Atoll Chemical Agent Disposal System (JACADS)
- Developed screening levels for construction/demolition workers
- Research laboratory converted to office
- Explosive residues in storage buildings
- Past herbicide research in laboratory





- Contrast the health risk interpretation:
 - Drinking water
 - Food consumption
 - Surface wipe samples
- Basic EPA Risk Methodology equates health risk to magnitude of chemical intake.
- How to estimate an Average Daily Intake (ADI) from available environmental data?



EPA Health Risk Fundamentals



Health Effects	Human Health	Formula
Cancer risk	<p style="text-align: center;">ILCR (Incremental Lifetime Cancer Risk)</p>	<p style="text-align: center;">ILCR = Chemical Intake X Cancer Slope Factor</p>
Noncancer	<p style="text-align: center;">HQ (Hazard Quotient)</p>	<p style="text-align: center;">HQ = $\frac{\text{Chemical Intake}}{\text{Reference Dose}}$</p>



Cancer Example

UNCLASSIFIED



ILCR = chemical intake X cancer slope factor

example:

A person incidentally ingesting sediment containing arsenic with a calculated intake of 7.23E-08 mg/kg-day



$$7.23\text{E-}08 \text{ mg/kg-d} \times 1.5 \text{ (mg/kg-d)}^{-1} = \mathbf{1.08\text{E-}07}$$



UNCLASSIFIED

Non Cancer Example- HQ

UNCLASSIFIED



example:

A worker incidentally ingests surface water with a calculated intake of $3.66\text{E-}07$ (mg/kg/day) of thallium



$$\text{HQ} = \frac{3.66\text{E-}07 \text{ (mg/kg/d)}}{7.00\text{E-}05 \text{ (mg/kg/d)}}$$

$$\text{HQ} = 5.2 \text{ E-}03$$



UNCLASSIFIED

Drinking Water Example

UNCLASSIFIED



- Measure chemical concentration
- Estimate daily water intake
- Concentration x consumption = mg



UNCLASSIFIED


Example



- Measure concentration in fish tissue
- Estimate fish consumption
- Concentration x consumption = mg



Wipe Sample Example



- Assume perfect sampling results of $50\mu\text{g}/100\text{ cm}^2$
- How do we use this surface sampling information to estimate intake?



Sampling Scenario

Child Day Care Center




UNCLASSIFIED



Sampling Scenario

Locked Mechanical Room



Potential Exposure Pathways




- ✓ Direct dermal contact
- ✓ Indirect ingestion from “mouthing behavior”
- ✓ Inhalation of settled particles resuspended from surface
- X Inhalation of semi-volatiles absorbed to surfaces (e.g., laminated, plastic)





Wipe Sample Interpretation

Exposure Assessment



$$ADI_{ing} = \frac{(SA \times Fd \times FT_{ss} \times Cs \times Ff \times FT_{sm} \times EV_{ing}) \times EF \times ED \times 10^{-3}}{BW \times AT}$$



Incidental Ingestion (fingers)

$$PD_{\text{ingestion (fingertips)}} = SA \cdot Fd \cdot FT_{SS} \cdot C_S \cdot F_f \cdot FT_{SM}$$


PD_{ingest}	Potential ingestion dose (mg/event)
SA	Exposed skin surface area per event (cm ² /event)
Fd	Fraction exposed skin surface area that actually contacts the surface (unitless)
FT_{SS}	Fraction transferred from surface to the skin (unitless)
C_S	Contaminant surface loading (mg/cm ²)
F_f	Fraction exposed skin area that contacts the mouth (unitless)
FT_{sm}	Fraction substance transferred from the skin to mouth (unitless)





Wipe Sample Interpretation

Exposure Assessment




$$ADI_{\text{derm}} = \frac{\sum (SA_i \times Fd_i) \times FT_{ss} \times Cs \times ABS \times EV_{\text{derm}} \times EF \times ED \times 10^{-3}}{BW \times AT}$$



Wipe Sample Interpretation

Exposure Assessment




$$ADI_{inh} = \left(\frac{f_{resp} \times C_s \times 10^4 \times A_s \times R}{V(\lambda_{dep} \times \lambda_a)} \right) \times \frac{IR_{inh} \times ET \times EF \times ED \times 10^{-3}}{BW \times AT}$$



Wipe Sample Interpretation

Safe Wipe Level Calculation





Wipe Sample Interpretation

Example Comparisons

Substance	Source	Safe level (ug/100 cm ²)
Beryllium	DOE	3 and 0.2
	TG 312	4.7
PCB	TSCA EPA	10
	TG312	1.60 and 9.04
	Michaud et al.	7.5
2,3,7,8 TCDD	EPA WTC	0.00002
	TG312	0.0000354
	Michaud et al.	0.00125





Wipe Sample Interpretation

References

Michaud, et al (1994) "PCB and Dioxin Re-Entry Criteria for Building Surfaces and Air", Journal of Exposure Analysis and Environmental Epidemiology, Vol 4, No. 2.

Contaminants of Potential Concern Committee (2003) "World Trade Center Indoor Environment Assessment: Selecting Contaminants of Potential Concern and Setting Health-Based Benchmarks".

USACHPPM, (2009) Technical Guide 312 Health Risk Assessment Methods and Screening Levels for Evaluating Office Worker Exposures to Contaminants on Indoor Surfaces Using Surface Wipe Data.





Wipe Sample Interpretation

References (cont)

Department of Energy, 10 CFR Part 850, Chronic Beryllium Disease Prevention Program; Final Rule

Toxic Substance Control Act, PCB Regulations: 40 CFR Part 761.61, PCB remediation waste.

Nicas, M Et al (2008) “A Study Quantifying the Hand-to-Face Contact Rate and Its Potential Application to Predicting Respiratory Tract Infection”, Journal of Occupational and Environmental Health.



Acknowledgements



- Ms. Hsieng-Ye Chang, MS, JD, PE
DuPont de Nemours
- Ms. Ronie Shackelford, BA, MA
USAPHC (Provisional)



UNCLASSIFIED

Backup Slides



UNCLASSIFIED



Inhalation of Resuspended Surface Particles

$$C_{air} = \frac{f_{resp} \cdot C_s \cdot 10^4 \cdot A_s \cdot R}{V \cdot \lambda_{dep} + V \cdot \lambda_a}$$

C_{air}	Resuspended air concentration (mg/m ³)
C_s	Contaminant surface loading (mg/cm ²)
f_{resp}	Fraction respirable (unitless)
10^4	Units conversion, cm ² to m ²
A_s	Source area (m ²)
V	Room volume (m ³)
R	Resuspension rate (1/hr)
λ_{dep}	Deposition loss rate (1/hr)
λ_a	Air exchange rate (air changes per hour [ACH])





Direct Dermal Contact

$$PD_{dermal} = \left[\sum_{i=1}^n SA_i \cdot Fd_i \right] \cdot FT_{SS} \cdot C_s$$

PD_{dermal}	Potential dermal dose (mg/event)
SA_i	Exposed skin surface area per event (cm^2 /event)
Fd_i	Fraction exposed skin surface area that actually contacts the surface (unitless)
i	Body part in contact with the surface (e.g., hand, forearm)
n	Total number of body parts in contact with the surface
FT_{SS}	Fraction transferred from surface to the skin (unitless)
C_s	Contaminant surface loading (mg/cm^2)

