

## **Deep-Water Ambient Noise Profiling; Marine Sediment Acoustics; and Doppler Geo-Acoustic Spectroscopy**

Michael J. Buckingham  
Marine Physical Laboratory, Scripps Institution of Oceanography  
University of California, San Diego  
9500 Gilman Drive  
La Jolla, CA 92093-0238  
phone: (858) 534-7977 fax: (858) 534-7641 email: [mbuckingham@ucsd.edu](mailto:mbuckingham@ucsd.edu)

Grant Number: N00014-10-1-0092  
<http://extreme.ucsd.edu>

### **LONG-TERM GOALS**

- 1) Deep-water ambient noise profiling Profile the spectral, temporal and spatial properties of broadband (3 Hz – 30 kHz) ambient noise from the surface to the bottom of the deepest ocean. Develop theoretical models of the second-order spatial statistics of the noise.
- 2) Marine sediment acoustics Develop a unified, physics-based model of sound wave and shear wave propagation in saturated, unconsolidated marine sediments.
- 3) Doppler geo-spectroscopy Develop the use of a light aircraft as a sound source for performing inversions for the geo-acoustic parameters of the seabed in shallow water.

### **OBJECTIVES**

- 1) The scientific objective of the deep-water ambient noise research is to measure the second-order spatial statistics of the ambient noise in the deepest oceans as a function of depth, from the sea surface to the seabed. Regions of interest include the Challenger Deep in the Mariana Trench (11 km), the Tonga Trench (9 km), and the Puerto Rico Trench (8 km). Environmental and system data will also be depth-profiled, including temperature, salinity, pressure and sound speed, along with all system motions (translational and rotational). Theoretical modeling of the spectral, spatial and temporal properties of the ambient noise will also be performed.
- 2) The sediment acoustics research is aimed at developing a unified theory of wave propagation in marine sediments in the form of the dispersion relations for the compressional and shear waves. Besides the frequency dependencies of the wave speeds and attenuations, these expressions will also return the dependence of the wave parameters on the mechanical properties of the sediment, namely porosity, density, grain size and overburden pressure.
- 3) The Doppler geo-acoustic spectroscopy technique, in which an aircraft is used as a sound source, is in support of the sediment acoustics research, providing a means of measuring the sound speed in the sediment at frequencies between 80 and 1000 Hz.

# Report Documentation Page

Form Approved  
OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE <b>2012</b>		2. REPORT TYPE <b>N/A</b>		3. DATES COVERED <b>-</b>	
4. TITLE AND SUBTITLE <b>Deep-Water Ambient Noise Profiling; Marine Sediment Acoustics; and Doppler Geo-Acoustic Spectroscopy</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Marine Physical Laboratory, Scripps Institution of Oceanography University of California, San Diego 9500 Gilman Drive La Jolla, CA 92093-0238</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release, distribution unlimited</b>					
13. SUPPLEMENTARY NOTES <b>The original document contains color images.</b>					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

## APPROACH

- 1) Deep-water ambient noise profiling A deep-diving autonomous instrument platform known as Deep Sound has been designed and developed by my research group. Deep Sound consists of a Vitrovex glass sphere housing a microprocessor for system control, along with data acquisition and storage electronics. External to the sphere are several hydrophones (bandwidth 3 Hz – 30 kHz, calibrated to equivalent depths of 12 km), which may be arranged in various vertical and horizontal configurations, and an environmental sensor package (CTD and sound velocimeter). The system is untethered, descending under gravity and, after releasing a drop weight at a pre-assigned depth, returning to the surface under buoyancy. Throughout the descent and ascent, at 0.6 m/s, acoustic and environmental data are continuously recorded. Three beacons (high intensity strobe, radio antenna, and Argos GPS) aid recovery of the system. Numerous fail-safe devices are onboard, intended to ensure that the weight is indeed dropped, thus allowing the system to return to the surface.

In conjunction with the experimental work using Deep Sound, I am developing a series of analytical models for the spatial coherence and cross-correlation properties of ambient noise in the deep ocean. These models address the directionality and bandwidth of the noise as they affect the cross-correlation function. The models will help us interpret the two-point measurements of noise recovered from Deep Sound.

- 2) Marine sediment acoustics My theoretical approach involves the development of the compressional and shear wave dispersion relations, based on inter-granular interactions. In the latest version of the grain-shearing theory, the viscosity of the pore fluid is included in the analysis, which leads to low frequency ( $< 10$  kHz) compressional wave behavior that is in accord with measurements made during the ONR-supported Sediment Acoustics Experiment 1999 (SAX 99). At higher frequencies, above 10 kHz, the effect of pore fluid viscosity is negligible and again the new theory fits the compressional wave data. Thus, the latest version of the theory fits all the available compressional wave and shear wave data from the SAX 99 experiments and elsewhere.
- 3) Doppler geo-acoustic spectroscopy A light aircraft flown at low level over the ocean acts as a sound source, which is used as the basis of an inversion technique for recovering the speed of sound in the seabed. The sound from the aircraft consists of a series of harmonics, typically 80, 160, 240, . Hz. Some of this acoustic energy penetrates the sea surface and reflects off the seabed, acquiring information about the sediment in the process. From recordings made on hydrophones in the water column and/or buried in the seabed, an inversion is performed which returns the phase speed of the compressional wave in the sediment. Once the compressional speed is known, most of the remaining geo-acoustic parameters are estimated using the correlations provided by the grain-shearing theory.

## WORK COMPLETED

Three versions of Deep Sound, designated the Mk. I, Mk. II and Mk. III, have been designed and built. The Mk. I has been deployed a number of times to great depths, approximately 6 km in the Philippine Sea and 9 km in the Mariana Trench, and is now showing signs of wear. Spalling of the glass sphere around the internal equator has occurred as a result of the compression due to the extreme hydrostatic pressure encountered at depth. Mk. I has now been retired and is no longer operational.

Deep Sound Mk. II was deployed in the Mariana Trench in November 2009, where it successfully recorded ambient noise on vertically and horizontally aligned hydrophones from the surface to a depth of 9 km over the acoustic frequency band from 3 Hz to 30 kHz.

Deep Sound Mk. III is the most sophisticated of the three systems, capable of descending to a depth of 11 km. On board is a sound velocimeter, a sing-around instrument which records sound speed directly, for comparison with the computed values from the CTD. This allows us to test the validity of sound speed algorithms for the extreme pressures found at the bottom of ocean trenches.

An attempt was made to deploy Deep Sound Mk. II and III in the Mariana Trench in July 2011, to depths of 9 km and 11 km, respectively. Working with a National Geographic group, a research vessel, the M/V Super Emerald, was chartered out of Saipan (at no cost to us) and used for the four-day deployment. Weather conditions were bad, with heavy seas and storms throughout the time at sea. The M/V Super Emerald was not well suited to the task, and at least one system (not one of ours) was lost. We should have been the last to deploy but decided against putting our systems in the water, given the extreme problems the other groups had experienced.

In early September 2012, Deep Sound Mk. II & III were deployed in the Tonga Trench from the R/V Roger Revelle. The hydrophones on board each of the systems were fitted with newly designed flow shields, intended to suppress the effects of turbulent flow generated by the motion of Deep Sound through the water column. Both Deep Sound systems descended to a depth of approximately 9 km, stayed on the bottom for an extended period, and returned to the surface after about 10 hours, all the while collecting broadband (3 Hz – 30 kHz) ambient noise data, along with environmental and system data.

An invited paper<sup>1</sup> on the Deep Sound has been published in a special issue of the *Journal of the Marine Technology Society* commemorating the Golden Anniversary of the dive of the manned submersible *Trieste* to the bottom of the Challenger Deep. A theoretical paper<sup>2</sup> on the directionality of ambient noise and its effects on the two point (vertical and horizontal) cross-correlation function has been published in the *Journal of the Acoustical Society of America* (JASA). Another paper<sup>3</sup> has been published in JASA in which a theoretical model of a three-dimensional noise field is developed. The model represents noise from a storm, showing a strong peak in the horizontal combined with significant vertical directionality. This is relevant to some of the data that were collected in the Philippine Sea, by Deep Sound Mk. I as an intense storm passed more or less overhead. An analysis of the storm data is described in a paper<sup>4</sup> that is currently under review at JASA. Wind-driven noise, also recorded during one of the deployments of Deep Sound Mk. I in the Philippine Sea, is described in another paper<sup>5</sup> that has been conditionally accepted for publication in JASA. A fifth, theoretical paper<sup>6</sup> has been published in JASA on band-limited noise, and the effects the filtering has on the cross-correlation function.

## RESULTS


Deep Sound Mk. I and II were both deployed to a depth of 9 km in the Mariana Trench in November 2009, where they continuously recorded ambient noise data, along with environmental information, on the descent and the ascent. The spectral level of the noise was found to increase slightly with increasing depth, even through the critical depth, by an amount that depends on frequency.

During an earlier deployment of Mk. I, in the Philippine Sea in May 2009, the sound of several rainstorms was recorded and is clearly visible in the spectrograms. An analysis shows that the rain noise is highly directional, with a strong lobe near the downward vertical<sup>4</sup>. Through an inversion, the storm was tracked across the sea surface, and some verification of the results was provided by ground truth measurements of rainfall rates made on board the research ship, the R/V Kilo Moana, supporting the deployment.

Wind-driven noise was recorded on two vertically aligned hydrophones during one of the deployments of Deep Sound Mk. I in the Philippine Sea in May 2009. The vertical coherence of the noise<sup>5</sup> is in excellent agreement with the theory of Cron & Sherman<sup>7</sup> for surface-generated noise in a semi-infinite, isovelocity ocean. This implies that the noise was essentially all downward traveling, suggesting that reflections from the sea bed, even in the proximity of the bottom, made a negligible contribution to the noise field. A comparison of the measured noise coherence at a depth of 3 km with Cron & Sherman's theory<sup>7</sup> is shown in Fig. 1.

My theoretical treatment of cross-correlation in spatially homogeneous, anisotropic ambient noise fields identifies the conditions necessary for extraction of the Green's function from directional noise<sup>2</sup>, on the assumption that the noise spectrum is white. Several examples of anisotropy are considered in the analysis, including Cron and Sherman's deep-water, downward-traveling ambient noise field<sup>7</sup>. In that particular case, because of the surface reflection, the Green's function depends on the depth of the transducers, whereas the cross-correlation function is independent of absolute position in the water column. It is, therefore, impossible to recover the Green's function from the noise, because the surface-reflected component will always be absent from the cross-correlation function. It is possible, however, to recover the sound speed in the medium from the noise, since the cross-correlation function exhibits sharp features at correlation delay times numerically equal to the travel time between the sensors.

In a related analysis, the effect on the cross-correlation function of band-limiting the ambient noise was investigated theoretically<sup>6</sup>. Horizontal and vertical alignments of the hydrophones were considered in the context of isotropic noise and deep-water noise, as represented by the Cron and Sherman<sup>7</sup> model. The focus of the analysis is on sensor separations that are very much greater than the longest wavelength (associated with the lowest frequency) in the pass-band. It turns out that, if the filter has a low-frequency roll-off, scaling as frequency to the power of  $2n$ , this acts as a differentiator of order  $2n$  on the cross-correlation function. In general, cross-correlation functions of filtered ambient noise do not exhibit delta functions that could be identified with the Green's function, but they may possess sharp features that allow the travel time between the sensors to be determined. From the travel time, it is possible to recover the sound speed in the medium. The mechanism that gives rise to these sharp features in the cross-correlation function is, however, quite different from that responsible for the appearance of delta functions in the derivative of the cross-correlation function of isotropic white noise.



**Fig. 1** Vertical coherence function at depth of 3 km on the ascent, Phillipine Sea. The jagged black line represents data, the smooth red line is Cron and Sherman's theory. a) Real part, b) imaginary part, c) magnitude.

In order to interpret the directional features of the storm noise recorded by Deep Sound Mk. I during the Philippine Sea deployment, a theoretical model of a three-dimensional noise field showing strong horizontal and vertical directionality has been developed<sup>3</sup>. The horizontal component of the noise is represented by a von Mises distribution<sup>8</sup>, taken from directional statistics, whilst the vertical component, associated with wave-breaking, is consistent with a set of dipoles distributed randomly across the sea surface. Expressions for the spatial coherence and cross-correlation of the noise at two horizontally aligned hydrophones are derived. Through a couple of parameters, the intensity and angular position of the storm can be adjusted, allowing the spatial statistics of the noise for various geometrical configurations to be investigated.

## **IMPACT/APPLICATIONS**

Deep Sound Mk. III is modular allowing the hydrophones in the current configuration to be replaced with any other type of sensor, for instance, dissolved oxygen, carbon dioxide or hydrocarbon sensors. Mk. III can even profile the local current vector, since the on board inertial navigation system tracks translational (and rotational) motion due to advection from the current.

My theory of wave propagation in marine sediments<sup>9</sup> has a variety of applications, particularly in regard to acoustic inversions for the geo-acoustic parameters of the seabed. Charles Holland and Ross Chapman are independently using the theory to develop numerical inversion schemes for recovering the properties of the bottom.

Michael Porter is developing a suite of 3-D acoustic propagation models. He is using my analytical models<sup>10, 11</sup> of the penetrable wedge and the conical seamount for comparison with his numerical results.

A conference was held recently on “Basic Science and the Future Warfighter” [ASD(R&E) Basic Science/Labs, Arlington, VA July 30-31 2012]. One of the prominent ideas discussed was the use of low-flying aircraft as sources of sound for underwater acoustics applications, including target detection and bottom characterization, as investigated theoretically and experimentally by my research group at SIO over recent years<sup>12</sup>.

## **TRANSITIONS**

As previously reported.

## **RELATED PROJECTS**

As previously reported.

## **REFERENCES**

1. D. R. Barclay, F. Simonet and M. J. Buckingham, “Deep Sound: a free-falling sensor platform for depth-profiling ambient noise in the deep ocean”, *Mar. Tech. Soc. J.* **43**, 144-150 (2010)

2. M. J. Buckingham, "On the two-point cross-correlation function of anisotropic, spatially homogeneous ambient noise fields in the ocean and its relationship to the Green's function", *J. Acoust. Soc. Am.*, **129**, 3562-3576 (2011)
3. S. C. Walker and M. J. Buckingham, "Spatial coherence and cross-correlation in three-dimensional ambient noise fields in the ocean", *J. Acoust. Soc. Am.*, **131**, 1079-1086 (2012)
4. D. R. Barclay and M. J. Buckingham, "Depth dependence of the power spectral density and vertical directionality of rain noise in the Philippine Sea", *J. Acoust. Soc. Am.*, **in review** (2012)
5. D. R. Barclay and M. J. Buckingham, "Depth dependence of wind-driven, broadband ambient noise in the Philippine Sea", *J. Acoust. Soc. Am.*, **conditionally accepted** (2012)
6. M. J. Buckingham, "Cross-correlation of band-limited ocean ambient noise", *J. Acoust. Soc. Am.*, **131**, 2643-2657 (2012)
7. B. F. Cron and C. H. Sherman, "Spatial correlation functions for various noise models", *J. Acoust. Soc. Am.*, **34**, 1732-1736 (1962); **38**, 885 (1965).
8. M. Evans, *Statistical Distributions*, Wiley, New York (2000)
9. M. J. Buckingham, "On pore-fluid viscosity and the wave properties of saturated granular materials including marine sediments", *J. Acoust. Soc. Am.*, **122**, 1486-1501 (2007)
10. M. J. Buckingham, "Theory of three-dimensional acoustic propagation in a wedgelike ocean with a penetrable bottom", *J. Acoust. Soc. Am.*, **82**, 198-210 (1987)
11. M. J. Buckingham, "Theory of acoustic propagation around a conical seamount", *J. Acoust. Soc. Am.*, **80**, 265-277 (1986)
12. M. J. Buckingham and E. M. Giddens, "Theory of sound propagation from a moving source in a three-layer Pekeris waveguide", *J. Acoust. Soc. Am.*, **120**, 1825-1841 (2006)

## PUBLICATIONS

### *Journal Articles & Chapters in Books*

1. M. J. Buckingham, "Cross-correlation of band-limited ocean ambient noise", *J. Acoust. Soc. Am.*, **131**, 2643-2657 (2012) [published, refereed]
2. D. R. Barclay and M. J. Buckingham, "Depth dependence of wind-driven, broadband ambient noise in the Philippine Sea", *J. Acoust. Soc. Am.*, **in review** (2012) [conditionally accepted, refereed]
3. D. R. Barclay and M. J. Buckingham, "Depth dependence of the power spectral density and vertical directionality of rain noise in the Philippine Sea", *J. Acoust. Soc. Am.*, **in review** (2012) [submitted, refereed]
4. S. C. Walker and M. J. Buckingham, "Spatial coherence and cross-correlation in three-dimensional ambient noise fields in the ocean", *J. Acoust. Soc. Am.*, **131**, 1079-1086 (2012) [published, refereed]
5. M. J. Buckingham, "On the two-point cross-correlation function of anisotropic, spatially homogeneous ambient noise fields in the ocean and its relationship to the Green's function", *J. Acoust. Soc. Am.*, **129**, 3562-3576 (2011) [published, refereed]



6. D. R. Barclay, F. Simonet and M. J. Buckingham, "Deep Sound: a free-falling sensor platform for depth-profiling ambient noise in the deep ocean ", *Mar. Tech. Soc. J.* **43**, 144-150 (2010) [published, refereed]
7. M. J. Buckingham, "Response to 'Comments on "Pore fluid viscosity and the wave properties of saturated granular materials including marine sediments [*J. Acoust. Soc. Am.* **122**, 1486-1501 (2007)]", *J. Acoust. Soc. Am.* **127**, 2099-2102 (2010) [published, refereed]
8. D. R. Barclay and M. J. Buckingham, "On the shapes of natural sand grains", *J. Geophys. Res.* **114**, B02209, doi:10, 1-12 (2009) [published, refereed]
9. M. J. Buckingham, "On the transient solutions of three acoustic wave equations: van Wijngaarden's equation, Stokes' equation and the time-dependent diffusion equation", *J. Acoust. Soc. Am.*, **124**, 1909-1920 (2008) [published, refereed]
10. M. J. Buckingham, "On pore-fluid viscosity and the wave properties of saturated granular materials including marine sediments", *J. Acoust. Soc. Am.*, **122**, 1486-1501 (2007) [published, refereed]
11. M. J. Buckingham and E. M. Giddens, "Theory of sound propagation from a moving source in a three-layer Pekeris waveguide", *J. Acoust. Soc. Am.*, **120**, 1825-1841 (2006) [published, refereed]
12. M. J. Buckingham and E. M. Giddens, "On the acoustic field in a Pekeris waveguide with attenuation in the bottom half-space", *J. Acoust. Soc. Am.*, **119**, 123-142 (2006) [published, refereed]
13. M. J. Buckingham, "Causality, Stokes' wave equation and acoustic pulse propagation in a viscous fluid", *Phys. Rev. E.*, **72**, 026610(9) (2005) [published, refereed].
14. M. J. Buckingham, "Compressional and shear wave properties of marine sediments: comparisons between theory and data", *J. Acoust. Soc. Am.*, **117**, 137-152 (2005) [published, refereed].
15. M. J. Buckingham, "Acoustic remote sensing of the sea bed using propeller noise from a light aircraft," in *Sounds in the Sea: Introduction to Acoustical Oceanography*, edited by H. Medwin, (Cambridge University Press, Cambridge, 2005) pp. 581-597 [published, refereed].
16. P. D. Thorne and M. J. Buckingham, "Measurements of the form function and total scattering cross section for suspensions of sands," *J. Acoust. Soc. Am.*, **116**, 2976-2890 (2004) [published, refereed].
17. M. J. Buckingham, "A three-parameter dispersion relationship for Biot's fast compressional wave in a marine sediment," *J. Acoust. Soc. Am.*, **116**, 769-776 (2004) [published, refereed].
18. M. J. Buckingham, "On the sound field from a moving source in a viscous medium," *J. Acoust. Soc. Am.*, **114**, 3112-3118 (2003) [published, refereed].
19. T. R. Hahn, T. K. Berger, and M. J. Buckingham, "Acoustic resonances in the bubble plume formed by a plunging water jet," *Proc. Roy. Soc. Lond. A*, **459**, 1751-1782 (2003) [published, refereed].
20. M. J. Buckingham, E. M. Giddens, F. Simonet and T. R. Hahn, "Propeller noise from a light aircraft for low-frequency measurements of the speed of sound in a marine sediment," *J. Comp. Acoust.*, **10** (4), 445-464 (2002) [published, refereed].

21. M. J. Buckingham, E. M. Giddens, J. B. Pompa, F. Simonet and T. R. Hahn, "Sound from a light aircraft for underwater acoustics experiments?," *Acta Acust. united with Acust.*, **88** (5), 752-755 (2002) [published, refereed].
22. M. J. Buckingham and M. D. Richardson, "On tone-burst measurements of sound speed and attenuation in sandy marine sediments," *IEEE J. Ocean. Eng.*, **27** (3), 429-453 (2002) [published, refereed].
23. M. J. Buckingham and M. S. Garcés, "Airborne acoustics of explosive volcanic eruptions," *J. Comp. Acoust.*, **9** (3), 1215-1225 (2001) [keynote address, published, refereed].
24. N. G. Lehtinen, S. Adam, G. Gratta, T. K. Berger and M. J. Buckingham, "Sensitivity of an underwater acoustic array to ultra-high energy neutrinos," *Astroparticle Phys.*, **697**, 1-14 (2001) [published, refereed].
25. M. D. Richardson, K. B. Briggs, D. L. Bibee, P. A. Jumars, W. B. Sawyer, D. B. Albert, T. K. Berger, M. J. Buckingham, *et al.*, "Overview of SAX99: environmental considerations," *IEEE J. Ocean. Eng.*, **26** (1), 26-53 (2001) [published, refereed].
26. M. J. Buckingham, "Precision correlations between the geoacoustic parameters of an unconsolidated, sandy marine sediment," *J. Comp. Acoust.*, **9** (1), 101-123 (2001) [published, refereed].
27. M. J. Buckingham, "Wave propagation, stress relaxation, and grain-to-grain shearing in saturated, unconsolidated marine sediments," *J. Acoust. Soc. Am.*, **108**, 2796-2815 (2000) [published, refereed].
28. C. L. Epifanio, J. R. Potter, G. B. Deane, M. L. Readhead, and M. J. Buckingham, "Imaging in the ocean with ambient noise: the ORB experiments," *J. Acoust. Soc. Am.*, **106**, 3211-3225 (1999) [published, refereed].
29. M. J. Buckingham, "Theory of compressional and transverse wave propagation in consolidated porous media," *J. Acoust. Soc. Am.*, **106**, 575-581 (1999) [published, refereed].
30. M. J. Buckingham, "On the phase speed and attenuation of an interface wave in an unconsolidated marine sediment," *J. Acoust. Soc. Am.*, **106**, 1694-1703 (1999) [published, refereed].
31. M. J. Buckingham, "Acoustic daylight imaging in the ocean," in *Handbook on Computer Vision and Applications*, vol. 1, Sensors and Imaging, B. Jähne, H. Haubecker, and P. Geibler, Eds. San Diego: Academic Press, 1999, pp. 415-424 [published, refereed].
32. M. J. Buckingham, "Acoustic pulse propagation in dispersive media," in *New Perspectives on Problems in Classical and Quantum Physics. Part II. Acoustic Propagation and Scattering - Electromagnetic Scattering*, vol. 2, P. P. Delsanto and A. W. Sáenz, Eds. Amsterdam: Gordon and Breach, 1998, pp. 19-34 [published, refereed].
33. M. J. Buckingham, "Theory of compressional and shear waves in fluid-like marine sediments," *J. Acoust. Soc. Am.*, **103**, 288-299 (1998) [published, refereed].
34. G. B. Deane, M. J. Buckingham, and C. T. Tindle, "Vertical coherence of ambient noise in shallow water overlying a fluid seabed," *J. Acoust. Soc. Am.*, **102**, 3413-3424 (1997) [published, refereed].

35. N. M. Carbone, G. B. Deane, and M. J. Buckingham, "Estimating the compressional and shear wave speeds of a shallow-water seabed from the vertical coherence of ambient noise in the water column," *J. Acoust. Soc. Am.*, **103**, 801-813 (1997) [published, refereed].
36. M. J. Buckingham, "Sound speed and void fraction profiles in the sea surface bubble layer," *Appl. Acoust.*, **51**, 225-250 (1997) [published, refereed].
37. M. J. Buckingham and N. M. Carbone, "Source depth and the spatial coherence of ambient noise in the ocean," *J. Acoust. Soc. Am.*, **102**, 2637-2644 (1997) [published, refereed].
38. M. J. Buckingham, "Theory of acoustic attenuation, dispersion, and pulse propagation in unconsolidated granular materials including marine sediments," *J. Acoust. Soc. Am.*, **102**, 2579-2596 (1997) [published, refereed].

## **PATENTS**

As previously reported.

## **HONORS/AWARDS/PRIZES**

As previously reported.