

Naturally Occurring Culturable Aerobic Gut Flora of Adult *Phlebotomus papatasi*, Vector of *Leishmania major* in the Old World

Jaba Mukhopadhyay¹, Henk R. Braig³, Edgar D. Rowton¹, Kashinath Ghosh^{1,2*}

1 Division of Entomology, Walter Reed Army Institute of Research, Silver Spring, Maryland, United States of America, **2** Division of Experimental Therapeutics, Walter Reed Army Institute of Research, Silver Spring, Maryland, United States of America, **3** School of Biological Sciences, Bangor University, Bangor, Wales, United Kingdom

Abstract

Background: Cutaneous leishmaniasis is a neglected, vector-borne parasitic disease and is responsible for persistent, often disfiguring lesions and other associated complications. *Leishmania*, causing zoonotic cutaneous leishmaniasis (ZCL) in the Old World are mainly transmitted by the predominant sand fly vector, *Phlebotomus papatasi*. To date, there is no efficient control measure or vaccine available for this widespread insect-borne infectious disease.

Methodology/Principal Findings: A survey was carried out to study the abundance of different natural gut flora in *P. papatasi*, with the long-term goal of generating a paratransgenic sand fly that can potentially block the development of *Leishmania* in the sand fly gut, thereby preventing transmission of leishmania in endemic disease foci. Sand flies, in particular, *P. papatasi* were captured from different habitats of various parts of the world. Gut microbes were cultured and identified using 16S ribosomal DNA analysis and a phylogenetic tree was constructed. We found variation in the species and abundance of gut flora in flies collected from different habitats. However, a few Gram-positive, nonpathogenic bacteria including *Bacillus flexus* and *B. pumilus* were common in most of the sites examined.

Conclusion/Significance: Our results indicate that there is a wide range of variation of aerobic gut flora inhabiting sand fly guts, which possibly reflect the ecological condition of the habitat where the fly breeds. Also, some species of bacteria (*B. pumilus*, and *B. flexus*) were found from most of the habitats. Important from an applied perspective of dissemination, our results support a link between oviposition induction and adult gut flora.

Citation: Mukhopadhyay J, Braig HR, Rowton ED, Ghosh K (2012) Naturally Occurring Culturable Aerobic Gut Flora of Adult *Phlebotomus papatasi*, Vector of *Leishmania major* in the Old World. PLoS ONE 7(5): e35748. doi:10.1371/journal.pone.0035748

Editor: Markus M. Heimesaat, Charité - Campus Benjamin Franklin, Germany

Received: October 12, 2011; **Accepted:** March 23, 2012; **Published:** May 22, 2012

Copyright: © 2012 Mukhopadhyay et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This study was funded by the Department of Defense through the In-House Laboratory Independent Research program at the Walter Reed Army Institute of Research. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: Kashinath.ghosh@us.army.mil

Introduction

Phlebotomine sand flies (Diptera: Psychodidae) are important vectors of leishmaniasis, Carrion's disease or bartonellosis, and a variety of arboviral diseases [1,2,3]. Not only are novel viruses currently being discovered in sand flies, but also different reservoirs are being identified for pathogens and parasites of human diseases, transmitted by sand flies. The distribution areas of sand flies and the diseases they transmit are also expanding. New viral diseases of humans transmitted by sand flies are being reported as well [4,5,6,7]. From a public health viewpoint, however, their greatest importance is as vectors of leishmaniasis [8,9,10]. The genus *Phlebotomus* in the Old World, and *Lutzomyia* in the New World, include most of the important vectors of human leishmaniasis. This disease complex is widely distributed in tropical and subtropical regions of the Americas, Africa, southern Europe, and central Asia. It is estimated that some 350 million people in the world are at risk of acquiring leishmaniasis and that approximately 12 million people are currently infected [11,12]. After malaria, leishmaniasis is the second most important vector-borne parasitic disease and a leading cause of death. There are

500,000 annual new cases of visceral leishmaniasis (VL) or kala-azar in the world and about one-half of them are in India. Bihar, the most affected state in India witnesses almost 90% of the new cases of VL each year with a 10% mortality rate [13,14].

Cutaneous leishmaniasis is more prevalent throughout the world and causes disfiguration and other associated complications. Cutaneous leishmaniasis and Zoonotic cutaneous leishmaniasis (ZCL), caused by *L. tropica* and *L. major*, respectively, are widely distributed in Turkey, Egypt, Israel, Saudi Arabia and the northern part of India, where mainly *P. sergenti* and *P. papatasi* have been incriminated as the vectors [15]. The Afro-Asian vector of ZCL, *P. papatasi* is widely distributed and is the type species of the genus. The distribution of *P. papatasi* coincides with the distribution of ZCL in most parts of the old world and shows little population differentiation between peridomestic sites and borrows of wild rodents [16,17]. Despite the demonstrated public health importance, relatively little attempt has been undertaken to block the transmission of this disease by this insect vector.

Information on breeding sites of *P. papatasi* is available from several countries [18,19]. In India, immature stages of *P. papatasi*

Report Documentation Page

Form Approved
OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE 22 MAY 2012	2. REPORT TYPE	3. DATES COVERED 00-00-2012 to 00-00-2012			
4. TITLE AND SUBTITLE Naturally Occurring Culturable Aerobic Gut Flora of Adult Phlebotomus papatasi, Vector of Leishmania major in the Old World		5a. CONTRACT NUMBER			
		5b. GRANT NUMBER			
		5c. PROGRAM ELEMENT NUMBER			
6. AUTHOR(S)		5d. PROJECT NUMBER			
		5e. TASK NUMBER			
		5f. WORK UNIT NUMBER			
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Walter Reed Army Institute of Research, Division of Entomology, 503 Robert Grant Ave, Silver Spring, MD, 20910		8. PERFORMING ORGANIZATION REPORT NUMBER			
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)		10. SPONSOR/MONITOR'S ACRONYM(S)			
		11. SPONSOR/MONITOR'S REPORT NUMBER(S)			
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Background: Cutaneous leishmaniasis is a neglected, vector-borne parasitic disease and is responsible for persistent, often disfiguring lesions and other associated complications. Leishmania, causing zoonotic cutaneous leishmaniasis (ZCL) in the Old World are mainly transmitted by the predominant sand fly vector, Phlebotomus papatasi. To date, there is no efficient control measure or vaccine available for this widespread insect-borne infectious disease. Methodology/Principal Findings: A survey was carried out to study the abundance of different natural gut flora in P. papatasi, with the long-term goal of generating a paratransgenic sand fly that can potentially block the development of Leishmania in the sand fly gut, thereby preventing transmission of leishmania in endemic disease foci. Sand flies, in particular, P. papatasi were captured from different habitats of various parts of the world. Gut microbes were cultured and identified using 16S ribosomal DNA analysis and a phylogenetic tree was constructed. We found variation in the species and abundance of gut flora in flies collected from different habitats. However, a few Gram-positive, nonpathogenic bacteria including Bacillus flexus and B. pumilus were common in most of the sites examined. Conclusion/Significance: Our results indicate that there is a wide range of variation of aerobic gut flora inhabiting sand fly guts, which possibly reflect the ecological condition of the habitat where the fly breeds. Also, some species of bacteria (B. pumilus, and B. flexus) were found from most of the habitats. Important from an applied perspective of dissemination, our results support a link between oviposition induction and adult gut flora.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified	Same as Report (SAR)	10	

have been consistently recovered from cattle sheds and human dwellings in urban areas [20,21]. In rural areas, they have been found in various habitats: unused poultry houses made of bricks and clay, manure heaps, caves, embankments, dried-up cesspits and latrines [22]. In Egypt, breeding sites of *P. papatasi* have been found in a similar range of ecotopes [23]. Rabbit holes in peri-domestic areas serve as breeding sites, which reduce the indoor abundance of *P. papatasi* in Tunisia [24]. In the Central Asian Republics of the former Soviet Union and neighboring states, burrows of the desert gerbil (*Rhombomys opimus*) are recognized as breeding sites [16]. Caves and dense vegetation of valleys are important in the Judean desert [25]. Dog shelters are important as breeding sites in peri-urban areas of southern Italy [17]. The ease with which *P. papatasi* adapt to an urban environment can be illustrated with the collection of sand flies near the bed and in the bathroom on the second floor in a house in a big city and in another highly urbanized area in southern Italy [26].

Currently, insecticide application at breeding sites is the method of choice for the vector control vis-à-vis control of disease. This control effort targets adult sand flies to bring down populations in order to reduce transmission. Application of insecticide may be limited due to adverse effects on the environment, human health, and the emergence of insecticide resistance in sand flies [27,28].

Sand flies spend a major part of their life as eggs, larvae and pupae in soil. During the immature stages, they are exposed to a variety of different soil microbes that are available for ingestion. In fact, gravid *P. papatasi* choose oviposition sites by presence of frass and certain soil bacteria [29,30]. Consequently, it is expected that the sand fly gut harbors a variety of microbial flora. The information on the distribution of the gut flora in feral sand fly populations, especially *P. papatasi*, across different habitat is still lacking. There are a few reports available on other species: from colonized *P. duboscqi* [31,32], from natural population of *L. longipalpis* [33], and from feral population of *P. argentipes* [34]. A very preliminary study on PCR fingerprinting of the gut flora from Moroccan *P. papatasi* flies identified just two bacteria [32,35]. There is also a small report on the distribution of gut flora from *P. papatasi* collected in Egypt [32,35]. Adler and Theodor suggested as early as 1929 that the presence of microbes in the midgut might interfere with *Leishmania* infection [36]. Later, Schlein et al. saw a reduction of infection rate of *L. major* in *P. papatasi* under the influence of yeasts and bacteria [37]. There is little doubt that the developing *Leishmania* in a sand fly gut is exposed to gut flora [38]. In an attempt to develop a strategy to block the transmission of leishmania, which has been demonstrated for some other vector-borne disease pathogens [39], we searched for nonpathogenic gut flora that could be genetically manipulated to release an anti-leishmanial substance and then be reintroduced into the sand fly gut through larval breeding habitats. The long-term objective would be to block or partially disrupt the metacyclogenesis of *Leishmania* sp. by the released product of the recombinant bacterium and thereby render the sand fly incapable of transmitting the disease. This will help to prevent further epidemic outbreak of leishmaniasis. A similar approach has been successfully applied in the development of paratransgenic *Rhodnius prolixus*, a vector of Chagas disease in Central America, with the help of genetically transformed *Rhodococcus rhodnii* [40]. A paratransgenic strategy has also been applied to *Glossina morsitans*, the vector of African sleeping sickness [41,42]. Additionally, a viral paratransgenic approach has been used to generate a transgenic *Anopheles gambiae*, a vector of malaria [43]. A paratransgenic control strategy has also been applied to the glossy-winged sharpshooter with the help of genetically marked *Alcaligenes* sp. [44,45]. The use of paratransgenesis is explored in the brine shrimp *Artemia* as a

model for controlling infectious diseases in mariculture and in an increasing number of insect groups such as fleas and termites [46,47,48]. In mosquitoes, symbiotic yeasts are discussed for control purposes [49]. This is only a short step to consider other eukaryotic symbionts of arthropods [50,51,52].

Here we examine the presence and distribution of different aerobic gut microbes of *P. papatasi*, the major vector of ZCL, in different habitats of various geographical parts of the world.

Materials and Methods

Collection of field samples

A large number of live sand flies were collected from India, Turkey and Tunisia. Sand flies were captured mainly from human dwellings, sheep sheds, chick pens, rabbit holes and mixed dwellings using light traps, or an aspirator and a flash light. Oral informed consent was obtained from head of households for indoor aspiration of sand flies and/or property owners for shed and outdoor collections that may have included light traps operated overnight. An explanation, in the local language, of the purposes for the collection, how the specimens would be used, the collection methods and any effects the collecting might have on the residents and/or their animals was provided before consent was obtained. Consents were listed in a written log kept by the collectors. Collected sand flies were released in containers with a plaster of Paris bottom. The containers were placed in individual plastic bags with moist cotton to provide necessary humidity for transportation to the laboratory.

Laboratory colony (control)

We used a laboratory colony of *P. papatasi* originated from field-collected samples from North Sinai, Egypt (PPNS). The colony is maintained at WRAIR following the method of Modi and Rowton [53].

Preparation of the media

Both liquid and solid agar based sterile media were prepared for the gut bacterial culture. Brain Heart Infusion (BHI) agar plates were prepared following the manufacturer's protocol (BD Biosciences; Cat. # 241830) and liquid culture broth was prepared using Terrific Broth Base (Invitrogen, Cat. # 22711-022).

Isolation and preservation of bacterial flora

Field collected sand flies were identified following the description by Lewis [54]. Only female *P. papatasi* was selected for the isolation of gut flora. In a sterile hood, each sand fly was rinsed in 70% ethanol for two minutes, followed by three quick rinses of sterile 1× PBS. Then the fly gut was dissected out and homogenized in about 60 µl of sterile 1× PBS in a sterile microfuge tube. Forty micro liter of the fly sample homogenate was quickly plated on BHI-agar plate, previously labeled with sand fly origin and number. The plates were subsequently placed in a 33±1°C incubator overnight. The remainder of the homogenate was cryopreserved in a -70°C freezer.

Selection and culture of clones

After overnight incubation, two to six colonies (depending on the number of colonies obtained from each fly) were picked up using a sterile toothpick and two copies of each colony were cultured in liquid media. The bacterial cultures were allowed to grow overnight in a shaker at 250 rpm at 33±1°C. One culture was used for isolation of DNA while the other was cryopreserved (using 17% sterile glycerol) in a freezer at -70°C. A relatively high incubation temperature was selected for the isolation of the flora

because we are mostly interested in generating a recombinant bacterial flora that can grow well and withstand a higher temperature when spread in natural breeding places in a tropical climate.

DNA extraction, PCR amplification and identification of the bacteria

Genomic DNA was isolated from individual cultures, using DNeasy Blood & Tissue Kit (Qiagen, Cat. #69581). Two sets of primers were used for amplification of the 16S rDNA: a) 533F- 5'-GT TGC CAG CAG CCG CGG TAA-3' and 1541R- 5'-AAG GAG GTG WTC CAR CC-3' [55,56]; and b) 8F-I 5'-AGA GTT TGA TYM TGG CTC AI-3' and 907R-I 5'-CCG TCA ATT CMT TTG AGT TI-3' [57]. PCR reactions were carried out in a 25 μ l reaction mixture containing 25–50 ng of template DNA, 1 \times PCR buffer (with 2.5 mM MgCl₂, 0.2–1 μ M of each primer, 0.2 mM dNTPs) and 1 Unit of Taq DNA polymerase (Promega, Cat. #M186). The PCR machine was programmed for the following amplification protocol: one cycle at 95°C for four min; 35 cycles for: 95°C (60 sec), 52°C (60 sec) and 72°C (90 sec) and the final extension step of one cycle at 72°C for six minutes. One non-template control was used for each run. PCR products were detected by agarose gel electrophoresis and purified with QIAquick gel extraction kit (Qiagen, Cat#28704). Nucleotide sequence for each amplicon was determined by using BigDye Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems, Cat# 4337455), and 1 U of one of the primers used during PCR amplification. Sequences were blasted and compared with the available sequences at the GenBank database. Isolates were recognized as the same species when their 16S rDNA sequences shared \geq 97% homology with complete 16S rDNA.

Data collection: After identification, the results were tabulated to show the relative abundance of different species of bacteria isolated from sand flies, collected from different locations and habitats.

Phylogenetic analysis

The sequences were manipulated in programs SeaView version 4 and MEGA version 5.05 [58,59]. Alignment of the sequences was based on the secondary structure of their RNAs with the alignment program SINA in the ARB software package using the silva comprehensive ribosomal RNA database version 108 [60]. The alignment was checked by hand. The best evolutionary model among 88 options for the analysis of the alignment was chosen with the help of the program jModeltest version 2 [61,62]: GTR+I+ Γ . Bayesian analysis was performed with the program Mr Bayes version 3.2 [63]. The analysis was carried out with two independent runs with four chains each for 1,000,000 generations of which the first 25% were dismissed. An average standard deviation of split frequencies of 0.0075 was reached, at which point the maximum Potential Scale Reduction Factor for parameter values was 1.002 suggesting convergence. The harmonic mean of the log likelihood of the resulting trees was –9,612.0. The tree was drawn with the help of the program Figtree version 1.3.1.

Results

A total of 107 *P. papatasi* field samples were dissected, of which 43 were collected from Tunisia, 31 originated from Turkey and 33 from India. Of the samples collected, 103 guts were cultured (two guts did not produce any colonies and two others were contaminated during preparation, Table 1). Forty-three female *P. papatasi* from one of our laboratory colonies originating from Egypt (PPNS) were used as control. The number of colonies




Figure 1. Bacterial clones of sand fly gut flora grown in BHI agar plate showing more than 100 colonies from a single *P. papatasi* female gut.

doi:10.1371/journal.pone.0035748.g001

generated from each fly gut varied widely. From some flies, there were as few as three colonies, while in others there were as many as 153 colonies (Figure 1). Two to six colonies from each sand fly gut were selected for further processing and identification of the flora.

The diversity of flora among *P. papatasi* populations, collected from several habitats in three different countries and a laboratory colony is shown in Figure 2. It is evident that there is more variation of the gut flora in the flies collected from animal dwellings of Tunisia and Turkey than in the samples captured from human dwellings in India.

In *P. papatasi* samples from Tunisia, *Bacillus flexus* was the most dominant bacterium irrespective of the collection habitat. Two other bacteria, *B. pumilus* and *B. megaterium* were also quite common. The flora from *P. papatasi* samples collected in Turkey was diversified with a clear dominance (31%) of *B. pumilus*. Other bacteria, including *B. clausii*, *B. cereus*, *B. subtilis* and *Brevibacillus brevis* were also present but at lower frequencies. Aerobic gut microbes in female *P. papatasi* collected from human dwellings of Patna, India, showed less diversity compared to the other two sites; the majority of them (54%) were *B. pumilus*. Four other species were also present in the captured samples but with much lower frequencies (Figure 2). The colonized sand flies from Egypt also showed a relative abundance of *B. pumilus* (30%) with few other microbes. With the exception of two species, *Enterobacter aerogenes* (Enterobacteriaceae, Proteobacteria) and *Plantibacter flavus* (Microbacteriaceae, Actinobacteria), all other bacteria belong to the families Bacillaceae and Paenibacillaceae (Figure 3).

Discussion

The present study is the most comprehensive evaluation of the distribution of intestinal flora of *P. papatasi* to date, as it describes the abundance of the bacterial gut flora from different habitats of three different countries. Our results show that *P. papatasi* harbor a wide selection of gut bacteria. Roughly, half of the detected bacteria are described for the first time from sand flies and some are described for the first time from insects (Table 2). The diversity of microbes from different habitats strongly suggests that the sand fly gut–microbial association is dependent on microbes in the environment in which those sand flies breed and live.

The gut flora in sand flies collected from sheep sheds and rabbit holes in Tunisia and Turkey showed more diversity than other groups. However, no significant differences in the distribution of


Table 1. Description of *P. papatasi* samples collected and screened for gut flora.

Site of collection	Habitat	No. of female flies examined	No of colonies produced	No. of clones examined
Tunisia, SS	Sheep shed	22	514	74
Tunisia, RH	Rabbit hole	21@	447	72
Turkey, SS	Chick/sheep shed - mixed	31#	527	80
Patna, India	Human dwellings	33#,@	518	86
Egypt	Lab colony	43	559	103

#Fly, which did not produce any colony.

@Contaminated sample.

doi:10.1371/journal.pone.0035748.t001

**Figure 2.** Distribution of gut flora of adult *P. papatasi* females.

doi:10.1371/journal.pone.0035748.g002




Figure 3. Bayesian 16S tree of gut flora of adult *P. papatasi* females. Posterior probabilities are given along internodes. The scale bar denotes substitutions per nucleotide for the branch lengths. Species that have been implicated in inducing oviposition behavior are highlighted in red. doi:10.1371/journal.pone.0035748.g003

the microbes were observed in the gut of sand flies collected from sheep shed or rabbit holes in Tunisia. Among the predominant flora observed from the flies collected from these two habitats, *B. cereus* is a potential human pathogenic bacterium [115]. The same is true for *En. aerogenes*, which has also been found to cause infections [116]. An interesting case is *B. circulans* because in the older Russian literature, it was mentioned together with *B. mycoides* as an entomopathogen of the gut of larval fleas [117]. Later, it was also recognized as a gut pathogen of mosquito larvae [118]. More recently, *B. circulans* was investigated as a potential probiotic for juvenile rohu in freshwater fish aquaculture [119]. Among the bacterial flora, *B. megaterium* and *B. flexus* are reported not only as non-pathogenic but also having some beneficial effect as probiotics [68,69].

The sand flies in Tunisia and Turkey were collected from animal shelters including sheep sheds, rabbit holes and poultry pens. Usually, the soil in and around these areas is contaminated by the excreta of the animals and other environmental contaminants making the soil a fertile medium for the growth of coprophilic bacteria. This contamination could explain the diversity of the bacterial flora found in the sand fly gut collected from these habitats. The diversity may be accentuated in places where animal shelters are in close proximity to agricultural land where the use of biofertilizers add more microbes to the nearby animal shelters. One example of this is the presence of *B. megaterium*, which have been found to have a good growth enhancement effect and yield, and have been used as a biofertilizer [120,121].

The diversity of bacterial population is somewhat restricted in the sand flies captured from India. Although other bacteria are present in less frequency, *B. pumilus* is the predominant bacterium found in the sand flies from Patna, India. Here, the majority of *P. papatasi* were obtained from human dwellings which is consistent with the anthropophilic nature of *P. papatasi* [122,123]. Blood-fed sand flies use the loose soil in the dark corners inside the mud houses as the most favorable place for oviposition [124]. Larvae are only exposed to the microbes inside the mud-house but not to the excreta of animals and other environmental contaminants. This may explain the lower diversity of the gut flora isolated from sand flies captured from human dwellings.

An unexpected result is that the Egypt lab colony seems to show a higher or similar diversity of bacterial flora as samples originated

from any of the natural habitats. This observation might be explained by the fact that the sand fly larvae are maintained in the laboratory on a diet composed of rabbit chow and rabbit feces, which are additional resources of gut flora and might have contributed to the bacterial diversity. Blood-fed females defecate gut bacteria along with the remains of the blood meal. Sand fly larvae are coprophagous. Therefore some gut bacteria are vertically transmitted to the next generation.

Bacillus pumilus, one of the most dominant bacteria of all the populations, is a Gram-positive, aerobic, rod-shaped, soil-dwelling bacterium. Like other *Bacillus* species, the spores produced by *B. pumilus* are more resistant than vegetative cells to heat, desiccation, UV radiation, γ -radiation, H_2O_2 , and starvation. This species has been found in extreme environments such as the interior of Sonoran desert basalt and the Mars Odyssey spacecraft [125,126]. The presence of *B. pumilus* in higher numbers in sand flies collected from human dwellings might be significant from the microbiological point of view as it has been shown that *B. pumilus* exhibits strong antifungal and antiviral activity [127,128]. Schlein et al. postulated that some gut bacteria might help to destroy fungi, thereby indirectly helping the development of *Leishmania* in the sand fly gut [37]. It is not clear at this point if *B. pumilus* is engaged in antifungal activity in the sand fly gut at all or if it acts together with other closely related *Bacillus* species or in combination with other gut factors to make the sand flies mycosis free. A fungi-free gut may help *Leishmania* survive which would make sand flies a more competent vector.

In the present study, a large number of *Bacillus* species was identified from *P. papatasi*. A preliminary study reported a different profile of bacteria. Species of *Enterobacter* and *Cronobacter* were isolated in greatest abundance from *P. papatasi* from Egypt by Dillon and others [35]. The authors emphasized that they used a rather selective medium and culture conditions. However, in a previous study on *P. argentipes* from India, we found a higher abundance of *Enterobacteriaceae* [34].

For the New World sand flies, Oliveira et al. found a high percentage of *Staphylococcus* sp. (28%) and *B. thuringiensis* (18%) in *Lutz. longipalpis* samples collected from Lapinha cave, Brazil [129]. They also recorded a relatively low percentage of *En. cloacae* (9%). We believe that these variations in the abundance of different bacteria from feral populations of sand flies are due to the ecological setting of their breeding habitat and species related.

Table 2. Distribution of *P. papatasi* gut bacteria among other hosts.

Bacterial species	other sand fly hosts	other host insects or mites	notes
Firmicutes			
Bacillaceae			
<i>Bacillus flexus</i> §		<i>Macrotermes carbonarius</i> [64]	plants [65], seaweed [66]
<i>Bacillus pumilus</i> §	<i>P. argentipes</i> [34]	<i>Apis mellifera</i> [67]	human and aquaculture probiotic [68,69], entomopathogen [70], strong oviposition inducer for gravid <i>P. papatasi</i> [29]
<i>Bacillus clausii</i> §			human probiotic [68]
<i>Bacillus badius</i> §			soil [71]
<i>Bacillus megaterium</i> §	<i>P. argentipes</i> [34]	<i>Macrotermes carbonarius</i> [64]	aquaculture probiotic [68], entomopathogen [72]
<i>Bacillus cereus</i> §	<i>P. argentipes</i> [34]	<i>Apis mellifera</i> [67]	human and veterinary probiotic [68], symbiont [73], entomopathogen [74], food pathogen [75], oviposition inducer for gravid <i>P. papatasi</i> [29]
<i>Bacillus licheniformis</i> §		<i>Dalbulus maidis</i> [76]	human, veterinary and aquaculture probiotic [68], very strong oviposition inducer for gravid <i>P. papatasi</i> [29]
<i>Bacillus endophyticus</i> §			plants [77]
<i>Bacillus subtilis</i> §	<i>P. argentipes</i> [34]	<i>Dalbulus maidis</i> [76]	human and veterinary probiotic [68]
<i>Bacillus circulans</i> §			entomopathogen [78]
<i>Bacillus [Lysinibacillus] fusiformis</i> §		<i>Apis mellifera</i> [79]	bioremediation [80,81]
<i>Lysinibacillus boronitolerans</i> §			soil [82]
<i>Oceanobacillus</i> sp.§		<i>Chironomus</i> sp. [83]	fermented food [84]
<i>Terribacillus saccharophilus</i> §			soil [85]
Paenibacillaceae			
<i>Brevibacillus brevis</i> §		<i>Malacosoma neustria</i> larvae [86]	plant antifungal [87]; entomo- and human pathogen, <i>B. laterosporus</i> : human probiotic [78]
<i>Brevibacillus reuszeri</i> §			soil, rhizobacterium [88]
<i>Paenibacillus</i> sp§		<i>Apis mellifera</i> [89]	entomopathogens [90]
Staphylococcaceae			
<i>Staphylococcus saprophyticus</i> §	<i>P. argentipes</i> [34]	<i>Musca domestica</i> [91]	very strong oviposition inducer for gravid <i>P. papatasi</i> [29]
unassigned family			
<i>Solibacillus</i> sp.§			forest soil [92]
Proteobacteria			
Enterobacteriaceae			
<i>Enterobacter aerogenes</i> §	<i>P. argentipes</i> [34], <i>L. longipalpis</i> [33]	<i>Apis mellifera</i> [93]	scale insect symbiont [94], human pathogen [95]
<i>Enterobacter cloacae</i> [35]	<i>P. argentipes</i> [34], <i>L. longipalpis</i> [33]		
<i>Cronobacter (Enterobacter) sakazakii</i> [35]		<i>Stomoxys calcitans</i> [96,97]	human pathogen [98]
<i>Erwinia</i> spp. [35]		Hemiptera [99]	phytopathogen [100]
<i>Serratia marcescens</i> [35]	<i>L. longipalpis</i> [33]	<i>Longitarsus</i> spp. [101]	entomo- and human pathogen [102,103]
Moraxellaceae			
<i>Acinetobacter</i> sp. [35]	<i>P. argentipes</i> [34], <i>L. longipalpis</i> [33]	<i>Bactericera cockerelli</i> [103]	human pathogen [104]
Pseudomonadaceae			
<i>Pseudomonas aeruginosa</i> [35]	<i>L. longipalpis</i> [33]	<i>Musca domestica</i> [105]	human pathogen [106]
<i>Pseudomonas</i> spp. [35]	<i>P. argentipes</i> [34]		entomo-, phyto- and human pathogen [106]
Actinobacteria			
Microbacteriaceae			
<i>Plantibacter flavus</i> §			grass [107]
<i>Microbacterium</i> spp. [32]	<i>P. argentipes</i> [34], <i>P. duboscqi</i> [32]	<i>Bemisia tabaci</i> [108]	human pathogens [109]

Table 2. Cont.

Bacterial species	other sand fly hosts	other host insects or mites	notes
Propionibacteriaceae			
<i>Propionibacterium</i> sp. [35]		<i>Psoroptes ovis</i> [110]	human and veterinary probiotic [111,112], <i>P. acne</i> : human pathogen [113]
Chloroflexi			
Chlorobacteria spp.* [32]	<i>P. duboscqi</i> * [32]		filamentous green non-sulfur bacteria [114]

[§]this report;^{*}immature stages only.

doi:10.1371/journal.pone.0035748.t002

The phylogenetic analysis shows strong support for all clades with the exception of *B. badius*. It is very reassuring that with the exception of *B. megaterium*, all species of our phylogenetic analysis using 16SRNA only showed similar relationship to a recent whole-genome phylogenetic analysis of the family Bacillaceae [130]. *Staphylococcus* species often clusters in 16S phylogenies within clades of *Bacillus* species [131]. *Bacillus fusiformis* of the literature cited here should be recognized as a *Lysinibacillus* species [119]. Species that have been observed by Radjame et al. to induce oviposition behavior in gravid *P. papatasi* females do not form a strict clade but cluster in a bigger group among the species recovered in this study [29].

Since there is very little information on the symbiotic association of bacteria with sand flies, gut colonization of bacteria is believed to be dependent on the larval food and the breeding soil. The larvae acquire many soil microbes during their immature stages of development which are believed to survive during the transformation until the adult emergence as reported in *P. duboscqi* by Volf et al. [31] (unpublished observation, Ghosh) However, in nature, adult sand flies may also have the opportunity to ingest microorganism through contaminated sugar meal derived from leaves, fruits or aphid honeydew taken between blood meals. Some sand fly species, in particular *P. papatasi*, may ingest microorganism from the plant cuticle while sucking the plant juice [132]. This explains some of the plant-associated bacteria found in our study.

Radjame et al. found that several soil bacteria significantly enhance the oviposition response of *P. papatasi* females [29]. The most pronounced effect was observed with *B. firmus* (P 0.00001 in cattle sheds), followed by *S. saprophyticus* (0.0003 in termite mounds and 0.002 in human dwellings), and *B. licheniformis* (0.0007 in cattle sheds, 0.003 in termite mounds and 0.0091 in human dwellings).

References

- Killick-Kendrick R (2002) Phlebotomine sand flies: biology and control. In: Farrell JP, ed. Leishmania. Dordrecht: Kluwer Academic. pp 33–43.
- Lane RP (1993) Sandflies (Phlebotominae). In: Lane RP, Crosskey RW, eds. Medical Insects and Arachnids. London: Chapman & Hall. pp 79–119.
- Rutledge LC, Gupta RK (2009) Moth flies and sand flies (Psychodidae). Med Vet Entomol. 2nd ed. London: Academic Press. pp 147–162.
- Depaquit J, Grandadam M, Fouque F, Andry PE, Peyrefitte C (2010) Arthropod-borne viruses transmitted by Phlebotomine sandflies in Europe: a review. Eurosurveillance 15: 40–47.
- Feldmann H (2011) Truly emerging-A new disease caused by a novel virus. N Engl J Med 364: 1561–1563.
- Papa A, Velo E, Bino S (2011) A novel phlebovirus in Albanian sandflies. Clin Microbiol Infect 17: 585–587.
- Yu X-J, Liang M-F, Zhang S-Y, Liu Y, Li J-D, et al. (2011) Fever with thrombocytopenia associated with a novel Bunyavirus in China. N Engl J Med 364: 1523–1532.
- Jhingarani A, Chatterjee M, Madhubala R (2008) Leishmaniasis: Epidemiological trends and diagnosis. In: Myler PJ, Fasel N, eds. Leishmania: After the Genome. Norfolk: Caister Academic Press. pp 1–14.
- Pavli A, Maltezou HC (2010) Leishmaniasis, an emerging infection in travelers. Int J Infect Dis 14: E1032–E1039.
- Ready PD (2010) Leishmaniasis emergence in Europe Eurosurveillance 15: e19505.
- Desjeux P (2004) Leishmaniasis: current situation and new perspectives. Comp Immunol Microbiol Infect Dis 27: 305–318.
- WHO (2011) Leishmaniasis. <http://www.who.int/leishmaniasis/en/>.
- Mondal S, Bhattacharya P, Ali N (2010) Current diagnosis and treatment of visceral leishmaniasis Expert Rev Anti-Infect Ther 8: 919–944.
- Sundar S, Mehta H, Suresh A, Singh SP, Rai M, et al. (2004) Amphotericin B treatment for Indian visceral leishmaniasis: Conventional versus lipid formulations. Clin Infect Dis 38: 377–383.
- Killick-Kendrick R (1990) Phlebotomine vectors of the leishmaniasis—a review. Med Vet Entomol 4: 1–24.
- Parvizi P, Benlarbi M, Ready PD (2003) Mitochondrial and *Wolbachia* markers for the sandfly *Phlebotomus papatasi*: little population differentiation between peridomestic sites and gerbil burrows in Isfahan province, Iran. Med Vet Entomol 17: 351–362.

Importantly, *B. pumilus* also induced oviposition of sand flies in cattle sheds significantly [29]. More studies are needed to find out the ability of *Bacillus* species to induce oviposition behavior under various conditions, especially in human dwellings. Of all the species considered, *B. pumilus* is particularly attractive because it has been recovered from all our study sites.

This study succeeded in identifying several candidate species for paratransgenesis in *P. papatasi*: *B. flexus*, *B. pumilus*, *B. licheniformis*, *B. megaterium* and *B. subtilis*. These bacteria are genetically tractable and trackable and are often used as probiotics. Most importantly, *B. pumilus* and *B. licheniformis* have been proposed as strong oviposition inducers for gravid *P. papatasi* [29]. The latter fact identifies those bacteria as true symbionts and not merely as environmental contaminants, which might be crucial for the dissemination of the bacteria into sand fly populations.

Acknowledgments

The authors would like to thank Dr. Elyes Zhioua and Dr. Ifhem Chelbi for their help in collecting sand flies in Tunisia and to Tobin Rowland for his help with sand fly maintenance at WRAIR. The opinions or assertions contained herein are the private views of the authors and are not to be construed as official, or as reflecting true views of the Department of the Army or the Department of Defense.

Author Contributions

Conceived and designed the experiments: KG JM HRB ER. Performed the experiments: JM KG. Analyzed the data: JM KG HRB. Contributed reagents/materials/analysis tools: JM KG ER HRB. Wrote the paper: JM KG HRB ER.

17. Tarallo VD, Dantas-Torres F, Lia RP, Otranto D (2010) Phlebotomine sand fly population dynamics in a leishmaniasis endemic peri-urban area in southern Italy. *Acta Trop* 116: 227–234.
18. Feliciangeli MD (2004) Natural breeding places of phlebotomine sandflies. *Med Vet Entomol* 18: 71–80.
19. Killick-Kendrick R (1999) The biology and control of phlebotomine sand flies. *Clin Dermatol* 17: 279–289.
20. Dhiman RC, Shetty PS, Dhanda V (1983) Breeding habitats of phlebotomine sandflies in Bihar, India. *Indian J Med Res* 77: 29–32.
21. Pandya AP, Niyogi AK (1980) Ecological studies on immature stage phlebotomid sandflies in Gujarat. *Indian J Med Res* 72: 355–358.
22. Sivagnaname N, Amalraj DD (1997) Breeding habitats of vector sandflies and their control in India. *J Commun Dis* 29: 153–159.
23. Doha S, Kamal H, Shehata M, Helmy N, Kader MA, et al. (1980) The breeding habitats of *Phlebotomus* sand flies (Diptera: Psychodidae) in El Agamy, Alexandria, Egypt. *J Egypt Soc Parasitol* 20: 747–752.
24. Chelbi I, Kaabi B, Derbali M, Ahmed SBH, Dellagi K, et al. (2008) Zoophylaxis: Impact of breeding rabbits around houses on reducing the indoor abundance of *Phlebotomus papatasi*. *Vector-Borne Zoonotic Dis* 8: 741–747.
25. Mueller GC, Kravchenko VD, Rybalov L, Schlein Y (2011) Characteristics of resting and breeding habitats of adult sand flies in the Judean desert. *J Vector Ecol* 36 Suppl.: S195–S205.
26. Dantas-Torres F, Latrofa MS, Otranto D (2010) Occurrence and genetic variability of *Phlebotomus papatasi* in an urban area of southern Italy. *Parasit Vectors* 3: e77.
27. Alexander B, Maroli M (2003) Control of phlebotomine sandflies. *Med Vet Entomol* 17: 1–18.
28. Kishore K, Kumar V, Kesari S, Dinesh DS, Kumar AJ, et al. (2006) Vector control in leishmaniasis. *Indian J Med Res* 123: 467–472.
29. Radjame K, Srinivasan R, Dhanda V (1997) Oviposition response of phlebotomid sandfly *Phlebotomus papatasi* to soil bacteria isolated from natural breeding habitats. *Indian J Exp Biol* 35: 59–61.
30. Wasserberg G, Rowton ED (2011) Sub-additive effect of conspecific eggs and frass on oviposition rate of *Lutzomyia longipalpis* and *Phlebotomus papatasi*. *J Vector Ecol* 36 Suppl.: S138–S143.
31. Volf P, Kiewegova A, Nemeč A (2002) Bacterial colonisation in the gut of *Phlebotomus dubosqi* (Diptera: Psychodidae): transtadial passage and the role of female diet. *Folia Parasitol* 49: 73–77.
32. Guernaoui S, Garcia D, Gazanion E, Ouhdouch Y, Boumezzough A, et al. (2011) Bacterial flora as indicated by PCR-temperature gradient gel electrophoresis (TGGE) of 16S rDNA gene fragments from isolated guts of phlebotomine sand flies (Diptera: Psychodidae). *J Vector Ecol* 36 Suppl. pp S144–S147.
33. Gouveia C, Ansens MD, Zahner V, Rangel EF, de Oliveira SMP (2008) Study on the bacterial midgut microbiota associated to different Brazilian populations of *Lutzomyia longipalpis* (Lutz & Neiva) (Diptera: Psychodidae). *Neotrop Entomol* 37: 597–601.
34. Hillesland H, Read A, Subhadra B, Hurwitz I, McKelvey R, et al. (2008) Identification of aerobic gut bacteria from the kala azar vector, *Phlebotomus argentipes*: A platform for potential paratransgenic manipulation of sand flies. *Am J Trop Med Hyg* 79: 881–886.
35. Dillon RJ, Elkordy E, Shehata M, Lane RP (1996) The prevalence of a microbiota in the digestive tract of *Phlebotomus papatasi*. *Ann Trop Med Parasitol* 90: 669–673.
36. Adler S, Theodor O (1929) Attempts to transmit *Leishmania tropica* by bite: the transmission of *L. tropica* by *Phlebotomus sergenti*. *Ann Trop Med Parasitol* 23: 1–18.
37. Schlein Y, Polacheck I, Yuva IB (1985) Mycoses, bacterial infections and antibacterial activity in sandflies (Psychodidae) and their possible role in the transmission of leishmaniasis. *Parasitology* 90: 57–66.
38. Azambuja P, Garcia ES, Ratcliffe NA (2005) Gut microbiota and parasite transmission by insect vectors. *Trends Parasitol* 21: 568–572.
39. Coutinho-Abreu IV, Zhu KY, Ramalho-Ortígo M (2010) Transgenesis and paratransgenesis to control insect-borne diseases: current status and future challenges. *Parasitol Int* 59: 1–8.
40. Fieck A, Hurwitz I, Kang AS, Durvasula R (2010) *Trypanosoma cruzi*: Synergistic cytotoxicity of multiple amphiphilic anti-microbial peptides to *T. cruzi* and potential bacterial hosts. *Exp Parasitol* 125: 42–347.
41. Aksoy S, Weiss B, Attardo G (2008) Paratransgenesis applied for control of tsetse transmitted sleeping sickness. *Adv Exp Med Biol* 627: 35–48.
42. Pontes M H, Dale C (2011) Lambda Red-mediated genetic modification of the insect endosymbiont *Sodalis glossinidius*. *Appl Environ Microbiol* 77: 1918–1920.
43. Ren X, Hoičzyk E, Rasgon JL (2008) Viral paratransgenesis in the malaria vector *Anopheles gambiae*. *PLoS Pathog* 4: e1000135.
44. Bextine B, Lauzon C, Potter S, Lampe D, Miller TA (2004) Delivery of a genetically marked *Alcaligenes* sp. to the glossy-winged sharpshooter for use in a paratransgenic control strategy. *Curr Microbiol* 48: 327–331.
45. Ramirez JL, Perring TM, Miller TA (2008) Fate of a genetically modified bacterium in foregut of glossy-winged sharpshooter (Hemiptera: Cicadellidae). *J Econ Entomol* 101: 1519–1525.
46. Husseneder C, Collier RE, Bourtzis K, Miller TA (2009) Paratransgenesis in termites. In: Bourtzis K, Miller TA, eds. *Symbiosis*. Boca Raton: CRC Press. pp 361–376.
47. Subhadra B, Hurwitz I, Fieck A, Rao DVS, Rao GS, et al. (2010) Development of paratransgenic *Artemia* as a platform for control of infectious diseases in shrimp mariculture. *J Appl Microbiol* 108: 831–840.
48. Erickson DL, Anderson NE, Cromar LM, Jolley A (2009) Bacterial communities associated with flea vectors of plague. *J Med Entomol* 46: 1532–1536.
49. Ricci I, Mosca M, Valzano M, Damiani C, Scuppa P, et al. (2011) Different mosquito species host *Wickerhamomyces anomalus* (*Pichia anomala*): perspectives on vector-borne diseases symbiotic control. *Antonie Van Leeuwenhoek* 99: 43–50.
50. Lantova L, Ghosh K, Svobodova M, Braig HR, Rowton E, et al. (2010) The life cycle and host specificity of *Psychodiella sergenti* n. sp. and *Ps. tobbi* n. sp. (Protozoa: Apicomplexa) in sand flies *Phlebotomus sergenti* and *Ph. tobbi* (Diptera: Psychodidae). *J Invertebr Pathol* 105: 182–189.
51. Perotti MA, Braig HR (2011) Eukaryotic ectosymbionts of Acari. *J Appl Entomol* 135: 514–523.
52. Votycka J, Lantova L, Ghosh K, Braig H, Volf P (2009) Molecular characterization of gregarines from sand flies (Diptera: Psychodidae) and description of *Psychodiella* n. g. (Apicomplexa: Gregarinida). *J Eukaryot Microbiol* 56: 583–588.
53. Modi GB, Rowton ED (1999) Laboratory maintenance of phlebotomine sand flies. In: Maramorosch K, Mahmood F, eds. *Maintenance of human, animal, and plant pathogen vectors*. Enfield, NH: Science Publishers. pp 109–121.
54. Lewis DJ (1978) Phlebotomine sand flies of the oriental region. *Bull Brit Mus (Nat Hist) Entomol* 37: 217–343.
55. Lane DJ (1991) 16S/23S rRNA sequencing. In: Stackebrandt E, Goodfellow M, eds. *Nucleic Acid Techniques in Bacterial Systematics*. Chichester: Wiley and Sons. pp 115–175.
56. Reysenbach AL, Wickham GS, Pace NR (1994) Phylogenetic analysis of the hyperthermophilic pink filament community in Octopus Spring, Yellowstone National Park. *Appl Environ Microbiol* 60: 2113–2119.
57. Ben-Dov E, Shapiro OH, Siboni N, Kushmaro A (2006) Advantage of using inosine at the 3 termini of 16S rRNA gene universal primers for the study of microbial diversity. *Appl Environ Microbiol* 72: 6902–6906.
58. Gouy M, Guindon S, Gascuel O (2010) SeaView version 4: a multiplatform graphical user interface for sequence alignment and phylogenetic tree building. *Mol Biol Evol* 27: 221–224.
59. Tamura K, Peterson D, Peterson N, Stecher G, Nei M, et al. (2011) MEGA5: Molecular Evolutionary Genetics Analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol* 28: 2731–2739.
60. Pruess E, Quast C, Knüttel K, Fuchs B, Ludwig W, et al. (2007) SILVA: a comprehensive online resource for quality checked and aligned ribosomal RNA sequence data compatible with ARB. *Nucleic Acids Res* 35: 7188–7196.
61. Guindon S, Gascuel O (2003) A simple, fast and accurate method to estimate large phylogenies by maximum-likelihood. *Syst Biol* 52: 696–704.
62. Posada D (2008) jModelTest: Phylogenetic model averaging. *Mol Biol Evol* 25: 1253–1256.
63. Ronquist F, Huelsenbeck JP (2003) MRBAYES 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19: 1572–1574.
64. Tay B-Y, Lokesh BE, Lee C-Y, Sudesh K (2010) Polyhydroxyalkanoate (PHA) accumulating bacteria from the gut of higher termite *Macrotermes carbonarius* (Blattodea: Termitidae). *World J Microbiol Biotechnol* 26: 1015–1024.
65. Sanchez-Gonzalez M, Blanco-Gamez A, Escalante A, Valladares AG, Olvera C, et al. (2011) Isolation and characterization of new facultative alkaliphilic *Bacillus flexus* strains from maize processing waste water (nejayote). *Lett Appl Microbiol* 52: 413–419.
66. Singh RP, Mantri VA, Reddy CRK, Jha B (2011) Isolation of seaweed-associated bacteria and their morphogenesis-inducing capability in axenic cultures of the green alga *Ulva fasciata*. *Aquat Biol* 12: 13–21.
67. Patil PB, Zeng Y, Coursey T, Houston P, Miller I, et al. (2010) Isolation and characterization of a *Nocardiopsis* sp. from honeybee guts. *FEMS Microbiol Lett* 312: 110–118.
68. Cutting SM (2011) *Bacillus* probiotics. *Food Microbiol* 28: 214–220.
69. Mandiki SNM, Milla S, Wang N, Blanchard G, Djonkack T, et al. (2011) Effects of probiotic bacteria on growth parameters and immune defence in Eurasian perch *Perca fluviatilis* L. larvae under intensive culture conditions. *Aquacult Res* 42: 693–703.
70. Molina CA, Cana-Roca JF, Osuna A, Vilchez S (2010) Selection of a *Bacillus pumilus* strain highly active against *Ceratitils capitata* (Wiedemann) larvae. *Appl Environ Microbiol* 76: 1320–1327.
71. Pichinoty F (1984) [Description of the type strain of *Bacillus badius*]. *Ann Microbiol (Paris)* B135: 21–27.
72. Aksoy HM, Ozman-Sullivan SK (2008) Isolation of *Bacillus megaterium* from *Aphis pomi* (Homoptera: Aphididae) and assessment of its pathogenicity. *J Plant Pathol* 90: 449–452.
73. Swiecicka I (2008) Natural occurrence of *Bacillus thuringiensis* and *Bacillus cereus* in eukaryotic organisms: a case for symbiosis. *Biocontrol Sci Technol* 18: 221–239.
74. Chatterjee S, Ghosh TS, Das S (2010) Virulence of *Bacillus cereus* as natural facultative pathogen of *Anopheles subpictus* Grassi (Diptera: Culicidae) larvae in submerged rice-fields and shallow ponds. *Afr J Biotechnol* 9: 6983–6986.
75. Stenfors Arnesen LP, Fagerlund A, Granum PE (2008) From soil to gut: *Bacillus cereus* and its food poisoning toxins. *FEMS Microbiol Rev* 32: 579–606.

76. Toledo AV, Alippi AM, de Remes Lenicov AMM (2011) Growth inhibition of *Beauveria bassiana* by bacteria isolated from the cuticular surface of the corn leafhopper, *Dalbulus maidis* and the planthopper, *Delphacodes kuscheli*, two important vectors of maize pathogens. *J Insect Sci* 11: 1–13.
77. Reva ON, Smirnov VV, Petterson B, Priest FG (2002) *Bacillus endophyticus* sp nov., isolated from the inner tissues of cotton plants (*Gossypium* sp.). *Int J Syst Evol Microbiol* 52: 101–107.
78. Sanders ME, Morelli L, Tompkins TA (2003) Sporeformers as human probiotics: *Bacillus*, *Sporolactobacillus*, and *Brevibacillus*. *Compr Rev Food Sci Food Safety* 2: 101–110.
79. Evans JD, Armstrong TN (2005) Inhibition of the American foulbrood bacterium, *Paenibacillus larvae larvae*, by bacteria isolated from honey bees. *J Apicul Res* 44: 168–171.
80. Lin C, Gan L, Chen Z-L (2010) Biodegradation of naphthalene by strain *Bacillus fusiformis* (BFN). *J Hazard Mater* 182: 771–777.
81. Zhao CQ, Zhang XY, Song D, Zheng LX, Chen WY (2010) Removal of chromium (VI) from tannery wastewater by immobilized *Bacillus fusiformis*. *J Soc Leather Technol Chem* 94: 21–25.
82. Ahmed I, Yokota A, Yamazoe A, Fujiwara T (2007) Proposal of *Lysinibacillus boronitolens* gen. nov. sp. nov., and transfer of *Bacillus fusiformis* to *Lysinibacillus fusiformis* comb. nov. and *Bacillus sphaericus* to *Lysinibacillus sphaericus* comb. nov. *Int J Syst Evol Microbiol* 57: 1117–1125.
83. Raats D, Halpern M (2007) *Oceanobacillus chironomi* sp nov., a halotolerant and facultatively alkaliphilic species isolated from a chironomid egg mass. *Int J Syst Evol Microbiol* 57: 255–259.
84. Whon TW, Jung M-J, Roh SW, Nam Y-D, Park E-J, et al. (2010) *Oceanobacillus kimchii* sp nov isolated from a traditional Korean fermented food. *J Microbiol* 48: 862–866.
85. An S-Y, Asahara M, Goto K, Kasai H, Yokota A (2007) *Terribacillus saccharophilus* gen. nov., sp nov and *Terribacillus halophilus* sp nov., spore-forming bacteria isolated from field soil in Japan. *Int J Syst Evol Microbiol* 57: 51–55.
86. Tozlu E, Dadasoglu F, Kotan R, Tozlu G (2011) Insecticidal effects of some bacteria on *Brachus dentipes* Baudi (Coleoptera: Bruchidae). *Fresenius Environ Bull* 20: 918–923.
87. Chandel S, Allan EJ, Woodward S (2010) Biological control of *Fusarium oxysporum* f. sp. *lycopersici* on tomato by *Brevibacillus brevis*. *J Phytopathol* 158: 470–478.
88. Yildirim E, Karlidag H, Turan M, Dursun A, Goktepe F (2011) Growth, nutrient uptake, and yield promotion of broccoli by plant growth promoting rhizobacteria with manure. *HortScience* 46: 932–936.
89. Genersch E (2010) American foulbrood in honeybees and its causative agent, *Paenibacillus larvae*. *J Invertebr Pathol* 103 Suppl. pp S10–S19.
90. Govindasamy V, Senthilkumar M, Magheshwaran V, Kumar U, Bose P, et al. (2010) *Bacillus* and *Paenibacillus* spp.: Potential PGPR for sustainable agriculture. In: Maheshwari DK, ed. *Plant Growth and Health Promoting Bacteria*. New York: Springer. pp 333–364.
91. Butler JF, Garcia-Maruniak A, Meek F, Maruniak JE (2010) Wild Florida house flies (*Musca domestica*) as carriers of pathogenic bacteria. *Fla Entomol* 93: 218–223.
92. Krishnamurthi S, Chakrabarti T, Stackebrand E (2009) Re-examination of the taxonomic position of *Bacillus silvestris* Rheims et al. 1999 and proposal to transfer it to *Solibacillus* gen. nov. as *Solibacillus silvestris* comb. nov. *Int J Syst Evol Microbiol* 59: 1054–1058.
93. Nada DV, Vukovic V, Nedic N (2010) Pathogenicity of some bacterial species isolated from the bee digestive tract. *Acta Vet* 60: 49–57.
94. Matsuura Y, Koga R, Nikoh N, Meng X-Y, Hanada S, et al. (2009) Huge symbiotic organs in giant scale insects of the genus *Drosicha* (Coccoidea: Monophlebidae) harbor flavobacterial and enterobacterial endosymbionts. *Zoolog Sci* 26: 448–456.
95. Chang E-P, Chiang D-H, Lin M-L, Chen T-L, Wang F-D, et al. (2009) Clinical characteristics and predictors of mortality in patients with *Enterobacter aerogenes* bacteremia. *J Microbiol Immunol Infect* 42: 329–335.
96. Hamilton JV, Lehane MJ, Braig HR (2003) Isolation of *Enterobacter sakazakii* from midgut of *Stomoxys calcitrans*. *Lancet* 9: 1355–1356.
97. Mramba F, Broce AB, Zurek L (2007) Vector competence of stable flies, *Stomoxys calcitrans* L. (Diptera: Muscidae) for *Enterobacter sakazakii*. *J Vector Ecol* 32: 134–139.
98. Healy B, Cooney S, O'Brien S, Iversen C, Whyte P, et al. (2010) *Cronobacter (Enterobacter sakazakii)*: An opportunistic foodborne pathogen. *Foodborne Pathog Dis* 7: 339–350.
99. Caspi-Fluger A, Zchori-Fein E (2010) Do plants and insects share the same symbionts? *Isr J Plant Sci* 58: 113–119.
100. Nadarasah G, Stavrinides J (2011) Insects as alternative hosts for phytopathogenic bacteria. *FEMS Microbiol Rev* 35: 555–575.
101. Kelley ST, Dobler S (2011) Comparative analysis of microbial diversity in *Longitarsus* flea beetles (Coleoptera: Chrysomelidae). *Genetica* 139: 541–550.
102. Sanchez-Contreras M, Vlisidou I (2008) The diversity of insect-bacteria interactions and its applications for disease control. *Biotechnol Genet Eng Rev* 25: 203–243.
103. Weber DJ, Rutala WA, Sickbert-Bennett EE (2007) Outbreaks associated with contaminated antiseptics and disinfectants. *Antimicrob Agents Chemother* 51: 4217–4224.
104. Doughari HJ, Nkaidemi PA, Human IS, Benade S (2011) The ecology, biology and pathogenesis of *Acinetobacter* spp.: An overview. *Microbes Environ* 26: 101–112.
105. Davari B, Kalantar E, Zahirnia A, Moosa-Kazemi SH (2010) Frequency of resistance and susceptible bacteria isolated from houseflies. *Iran J Arthropod-Borne Dis* 4: 50–55.
106. Silby MW, Winstanley C, Godfrey SAC, Levy SB, Jackson RW (2011) *Pseudomonas* genomes: diverse and adaptable. *FEMS Microbiol Rev* 35: 652–680.
107. Behrendt U, Ulrich A, Schumann P, Naumann D, Suzuki K (2002) Diversity of grass-associated Microbacteriaceae isolated from the phyllosphere and litter layer after mulching the sward; polyphasic characterization of *Subtercola pratensis* sp. nov., *Curtobacterium herbarum* sp. nov. and *Plantibacter flavus* gen. nov., sp. nov. *Int J Syst Evol Microbiol* 52: 1441–1454.
108. Indiragandhi P, Yoon C, Yang JO, Cho S, Sa TM, et al. (2010) Microbial communities in the developmental stages of B and Q biotypes of sweetpotato whitefly, *Bemisia tabaci* (Hemiptera: Aleyrodidae). *J Korean Soc Appl Biol Chem* 53: 605–617.
109. Gneiding K, Frodl R, Funke G (2008) Identities of *Microbacterium* spp. encountered in human clinical specimens. *J Clin Microbiol* 46: 3646–3652.
110. Hogg JC, Lehane MJ (2001) Microfloral diversity of cultured and wild strains of *Psoroptes ovis* infesting sheep. *Parasitology* 123: 441–446.
111. Cousin FJ, Mater DDG, Foligne B, Jan G (2011) Dairy *Propionibacteria* as human probiotics: A review of recent evidence. *Dairy Sci Technol* 91: 1–26.
112. Seo JK, Kim S-W, Kim MH, Upadhaya SD, Kam DK, et al. (2010) Direct-fed microbials for ruminant animals. *Asian-Australas J Anim Sci* 23: 1657–1667.
113. Jappe U (2003) Pathological mechanisms of acne with special emphasis on *Propionibacterium acnes* and related therapy. *Acta Derm Venereol* 83: 241–248.
114. Yamada T, Sekiguchi Y (2009) Cultivation of uncultured Chloroflexi subphyla: Significance and ecophysiology of formerly uncultured Chloroflexi 'Subphylum I' with natural and biotechnological relevance. *Microbes Environ* 24: 205–216.
115. Didelot X, Barker M, Falush D, Priest FG (2009) Evolution of pathogenicity in the *Bacillus cereus* group. *Syst Appl Microbiol* 32: 81–90.
116. De Gheldre Y, Maes N, Rost F, De Ryck R, Clevenbergh P, et al. (1997) Molecular epidemiology of an outbreak of multidrug-resistant *Enterobacter aerogenes* infections and in vivo emergence of imipenem resistance. *J Clin Microbiol* 35: 152–160.
117. Vasilev GI, Bazanova LP (1987) [The effect of the entomopathogenic bacteria-*Bacillus myocoides* Flugge and *Bacillus circulans* Jordan-on the gut microflora of larval fleas] In: Cherepanov AI, ed. *Ekologiya i geografiya chlenistonogikh Sibiri* [Ecology and Geography of Arthropods of Siberia]. Novosibirsk: Nauka. pp 208–210.
118. Darriet F, Hougard J-M (2002) An isolate of *Bacillus circulans* toxic to mosquito larvae. *J Am Mosq Control Assoc* 18: 65–67.
119. Ghosh K, Sen SK, Ray AK (2003) Supplementation of an isolated fish gut bacterium, *Bacillus circulans*, in formulated diets for rohu, *Labeo rohita*, fingerlings. *Isr J Aquacult Bamidgch* 55: 13–21.
120. Chakraborty U, Chakraborty B, Basnet M (2006) Plant growth promotion and induction of resistance in *Camellia sinensis* by *Bacillus megaterium*. *J Basic Microbiol* 46: 186–195.
121. de Freitas JR, Banerjee MR, Germida JJ (1997) Phosphate-solubilizing rhizobacteria enhance the growth and yield but not phosphorus uptake of canola (*Brassica napus* L.). *Biol Fertil Soils* 24: 4358–4364.
122. Morsy TA, Aboul ERG, Abdelmawla MM, el Gozamy BM (1993) Counter-immuno-electrophoresis, a modified technique for the identification of blood meals of sandflies collected from Qalyubia Governorate, Egypt. *J Egypt Soc Parasitol* 23: 109–132.
123. Namita M, Joshi V, Bansal SK (1991) Host preference pattern of phlebotomine sandflies of Bikaner city. *Indian J Med Res* 93: 328–329.
124. Singh R, Lal S, Saxena VK (2008) Breeding ecology of visceral leishmaniasis vector sandfly in Bihar state of India. *Acta Trop* 107: 117–120.
125. Benardini JN, Sawyer J, Venkateswaran K, Nicholson WL (2003) Spore, UV and acceleration resistance of endolithic *Bacillus pumilus* and *Bacillus subtilis* isolates obtained from Sonoran desert basalt: implications for lithopanspermia. *Astrobiology* 3: 709–717.
126. Kempf MJ, Chen F, Kern R, Venkateswaran K (2005) Recurrent isolation of hydrogen peroxide-resistant spores of *Bacillus pumilus* from a spacecraft assembly facility. *Astrobiology* 5: 391–405.
127. Bottone EJ, Peluso RW (2003) Production by *Bacillus pumilus* (MSH) of an antifungal compound that is active against *Mucoraceae* and *Aspergillus* species: preliminary report. *J Med Microbiol* 52: 69–74.
128. Naruse N, Tenmyo O, Kobaru S, Kamei H, Miyaki T, et al. (1990) Pumilacidin, a complex of new antiviral antibiotics. Production, isolation, chemical properties, structure and biological activity. *J Antibiot (Tokyo)* 43: 267–280.
129. Oliveira S M, Moraes BA, Gonçalves CA, Giordano-Dias CM, D'Almeida JM, et al. (2000) [Prevalence of microbiota in the digestive tract of wild females of *Lutzomyia longipalpis* Lutz & Neiva, 1912] (Diptera: Psychodidae). *Rev Soc Bras Med Trop* 33: 319–322.
130. Schmidt TR, Scott EJ, II, Dyer DW (2011) Whole-genome phylogenies of the family Bacillaceae and expansion of the sigma factor gene family in the *Bacillus cereus* species-group. *BMC Genomics* 12: e430.
131. Boehme K, Fernandez-No IC, Barros-Velazquez J, Gallardo JM, Canas B, et al. (2011) Rapid species identification of seafood spoilage and pathogenic

Gram-positive bacteria by MALDI-TOF mass fingerprinting. *Electrophoresis* 32: 2951–2965.

132. Schlein Y, Jacobson RL (1994) Mortality of *Leishmania major* in *Phlebotomus papatasi* caused by plant feeding of the sand flies. *Am J Trop Med Hyg* 50: 20–27.