

Seabed Geoacoustic Planning Support for the QPE Uncertainty DRI

Charles W. Holland
The Pennsylvania State University
Applied Research Laboratory
P.O. Box 30, State College, PA 16804-0030
phone: (814) 865-1724 fax (814) 863-8783 email: holland-cw@psu.edu

Grant Number: N00014-07-1-0252

LONG TERM GOALS

Develop capability for quantifying, predicting and exploiting (QPE) the impact of seabed uncertainty on sonar system performance.

OBJECTIVES

The objectives were to contribute to the development of a science plan for the QPE DRI that builds on the recent advances in the understanding of uncertainty.

APPROACH

Assisted in shaping science plan drawing on the open literature, Taiwanese and other QPE colleagues, in close collaboration with members of the Seabed Characterization Working Group including Larry Mayer and Brian Calder (University of New Hampshire), Jim Lynch (WHOI), Bob Miyamoto (APL-UW) as well as Phil Abbot (team leader for Acoustics Working Group).

WORK COMPLETED

- 1) Helped identify/obtain in southern East China Sea, pertinent to DRI goals
 - key results from geologic/geophysical/geoacoustic community
 - key missing elements of seabed characterization
 - methods for quantifying and predicting geoacoustic uncertainty
- 2) Provided guidance on seabed uncertainty and variability issues in order to develop goals and a roadmap for the ensuing 4-year DRI.

RESULTS

Unknown acoustic boundary conditions at the seabed lead to large uncertainties in TL and noise predictions. Spatial and possibly temporal (see features discussed below) variability will increase those uncertainties. A literature search was performed to identify seabed processes and features that might play a significant role in acoustic performance uncertainty. A short summary follows.

Report Documentation Page

*Form Approved
OMB No. 0704-0188*

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE 30 SEP 2007	2. REPORT TYPE	3. DATES COVERED 00-00-2007 to 00-00-2007			
4. TITLE AND SUBTITLE Seabed Geoacoustic Planning Support for the QPE Uncertainty DRI		5a. CONTRACT NUMBER			
		5b. GRANT NUMBER			
		5c. PROGRAM ELEMENT NUMBER			
6. AUTHOR(S)		5d. PROJECT NUMBER			
		5e. TASK NUMBER			
		5f. WORK UNIT NUMBER			
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Pennsylvania State University, Applied Research Laboratory, PO Box 30, State College, PA, 16804-0030		8. PERFORMING ORGANIZATION REPORT NUMBER			
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)		10. SPONSOR/MONITOR'S ACRONYM(S)			
		11. SPONSOR/MONITOR'S REPORT NUMBER(S)			
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified	Same as Report (SAR)	3	

The seabed processes that appear to be important are:

- Erosion, enhanced by seismicity, landslides (sub-aerial and submarine) leading to riverine sediment deposition (e.g., Lanyang Hsi, NE Taiwan). This region is somewhat unique inasmuch as Taiwan supplies ~400 Mtons/year of sediment to the ocean which is ~2% of the total world's sediment discharge, yet Taiwan only accounts for ~0.02% of the earth's exposed surface.
- Sediment transport across the ECS shelf from the Yangtze river around NE tip of Taiwan. The importance and exact path of this transport is under debate. These are fine-grained sediments that eventually end up in the Southern Okinawa Trough.
- Biogenic contributions: Shell material is abundant at the shelf edge and upper slope is generally associated with sandy sediments. Muddy sediments also contain 25%-40% CaCO₃ from planktonic foraminifera and pteropod shells associated with upwelling.
- Rock fragments are common and abundant constituents of all shelf sands, with a relative abundance of sandstone clasts.
- The ECS shelf is strongly shaped by glacio-eustasy, alternating continental and marine sediments corresponding to glaciation and interglaciation are evident.
- The Southern Okinawa Trough (SOT) is shaped by sediment transported by currents from China (Yangtze), Taiwan, and equatorial Pacific. While SOT is only 10% of entire OT, it contains the majority of OT sediments. Mass wasting events are an extremely important in the SOT and greatly contribute to the complexity of the area.
- The Kuroshio Current seems to play a role in the distribution of sediments under/near its path. There are sediment "mixtures" probably associated with loop currents on the shelf and directional changes of the Kuroshio. Beneath the main Kuroshio flow, sediments are non-biogenic fine-grained mud.

There are several seabed features that may have important implications including:

- Tidal induced sand ridges (left from the postglacial sea level rise): tidal currents are high, > 1 m/s in some areas, and 0.75 cm/s at the shelf break. There is strong evidence of large sand ridges on the Chilung shelf that appear to vary over seasonal time scales.
- On outer shelf and slope, presence of deep thermogenic methane leads to mud volcanoes (5-40 m in height; radii ~20-600 m have been observed) and pockmarks; (up to 40m in depth, radii up to 150m). Besides the bathymetric expression, the presence of gas drastically alters seabed physical properties relative to the surrounding sediments. Some of the mud volcanoes are expected to be "active" meaning that they are actively venting methane (e.g., in bubble plumes) and/or oil into the ocean. Very little is known about the temporal scales associated with the methane fluxes.

IMPACT/APPLICATIONS

The study was useful inasmuch as it provides a basis for developing goals and a roadmap for the ensuing 4-year DRI. An important question that will be addressed in the follow-on program is how do uncertainties in bottom geoacoustic properties and morphology affect uncertainty in acoustic performance prediction (propagation, noise, and coherence).

RELATED PROJECTS

none