

# **Bedforms Generated by Turbidity Currents in Continental Margins**

Marcelo H. Garcia  
Department of Civil and Environmental Engineering  
University of Illinois at Urbana-Champaign  
205 North Mathews Avenue  
Urbana, Illinois 61801  
phone: (217) 244-4484, fax: (217) 333-0687, email: [mhgarcia@uiuc.edu](mailto:mhgarcia@uiuc.edu)

Award #: N00014-93-1-0044  
<http://cee.ce.uiuc.edu/hydro/>

## **LONG-TERM GOAL**

Our long-term scientific goal is to develop sound theoretical models for submarine sediment movement that can be used to predict the initiation, spatial development, and time duration of mud flows and turbidity currents. Of particular interest are the characteristics of the sedimentary deposits that these flows generate, in particular their capability to develop bed forms and gullies by means of sediment erosion and deposition in continental margins.

## **OBJECTIVES**

This past year we have concentrated on testing the hypothesis that turbidity currents are capable of forming gullies along sloping beds. In particular, our main objective has been to elucidate the role played by flow instabilities at the front of very wide turbidity flows on the formation of submarine channels. A second objective has been to explore, with the help of laboratory experiments, the ability of turbidity currents to generate ripples, dunes and antidunes, along their path.

## **APPROACH**

Our approach has consisted of theoretical work, laboratory experiments, and numerical modeling. The theoretical work has concentrated on a perturbation analysis of the equations of motion for turbidity currents carrying a conservative component such as clay, with the goal of assessing the conditions for which bed instabilities will grow and bedforms will develop. Laboratory experiments have been directed to study the formation of bedforms by three-dimensional turbidity currents emanating from a line source at the head of a model slope. Numerical experiments have been conducted as well, to determine the influence of bottom slope and initial sediment concentration on the genesis of gullies at field scale.

## **WORK COMPLETED**

A linear stability analysis of the equations of motion for turbidity currents (Parker et al., 1986) flowing over a movable sediment bed was completed. A conservative component (i.e. clay) was added to a sand-laden turbidity current. The main goal was to study the growth or decay of small bed-level perturbations in the flow direction. The conditions, as prescribed by the theoretical analysis, were then used to design and conduct a set of laboratory experiments with channelized turbidity currents, with

# Report Documentation Page

*Form Approved*  
*OMB No. 0704-0188*

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE <b>SEP 2000</b>		2. REPORT TYPE		3. DATES COVERED <b>00-00-2000 to 00-00-2000</b>	
4. TITLE AND SUBTITLE <b>Bedforms Generated by Turbidity Currents in Continental Margins</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Department of Civil and Environmental Engineering,,University of Illinois at Urbana-Champaign,,205 North Mathews Avenue,,Urbana,,IL,61801</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>Same as Report (SAR)</b>	18. NUMBER OF PAGES <b>5</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

the goal of observing the development or lack of bedforms. Preliminary experiments were also completed in the MARGINS tank, to explore the generation of submarine gullies by 3-D turbidity currents. Also, density (saline) currents were produced in order to analyze the effect of these flows on bed deformation. Investigation of front dynamics (front instabilities) for both turbidity and density (saline) currents, has also been performed using image analysis from videotaped experiments.

To complement and guide the laboratory work with turbidity flows from a line source, a numerical model was set up. The model combines the set of equations for turbidity currents proposed by Parker et al. (1986), with a simple, rule-based cellular model proposed by Murray and Paola (1994) for the modeling of braided streams.

## **RESULTS**

So far, no gullies have been observed in the laboratory. Only bedforms such as three-dimensional ripples and dunes, as well as long-wavelength antidune-like sediment waves were observed in the MARGINS tank; all of them generated by turbidity currents and saline currents emanating from a line source.

A variety of scales were found within the patterns of frontal lobes and clefts observed in turbidity and density currents flowing along a wide channel. Figure 1 shows a fractal representation of these scales, obtained by computing the number  $N$  of lobes of a characteristic length  $\lambda$ , for a large number of front images. The fractal exponents (dimension) are found to be slightly different for turbidity and density currents. Also, the life time of some of the relevant scales of lobes for turbidity currents is shown in Figure 2, from where a characteristic velocity scale for frontal processes is derived (slope of the fit), this velocity being similar to the propagation velocity of the front.

Both the theoretical perturbation analysis and the laboratory experiments have shown that clay content in turbidity currents plays an important role in the formation of submarine bedforms.

## **IMPACT/APPLICATION**

Our results have important implications for the recognition of morphological features in the field. Turbidity currents are considered to be the sculptors of many, but not all, submarine canyons. Hence it is reasonable to consider the possibility that gullies could also be the result of turbidity current activity, even though their reproduction in the laboratory has so far being rather elusive (Field et al., 1999). The laboratory experiments have clearly shown the capability of these currents for producing longitudinal small-scale bedforms such as dunes and ripples, as well as long-wavelength antidunes. Such antidunes are thought to be related to interfacial instabilities at the upper boundary of the flow. Understanding the front dynamics is also relevant in order to characterize mixing processes and evolution of frontal instabilities, and their effects on the overall current mechanics. Our understanding of the mechanics of bedforms in continental margins until now has been very limited and we hope that our findings will facilitate both the interpretation of the geologic record as well as the design and placement of submarine structures on stable sediment deposits.

## **REFERENCES**

Field, M.E., Gardner, J.V. and Prior, D.B., 1999. Geometry and significance of stacked gullies on the northern California slope. *Marine Geology, Special Issue, Vol. 154*, pp. 271-286.

Murray, A.B. and Paola, C., 1994. A cellular model of braided rivers. *Nature*, Vol. 371, September, pp. 54-57.

Parker, G., Fukushima, Y. and Pantin, H.M., 1986. Self-accelerating turbidity currents. *J. Fluid Mechanics*, Vol. 171, pp. 145-181.

## **FY 2000 PUBLICATIONS**

Admiraal, D.M., and Garcia, M.H. 2000. Laboratory measurements of suspended sediment concentration using an acoustic concentration profiler (ACP). *Experiments in Fluids*, 28, 116-127.

Fedele, J.J. and Garcia, M.H. 1999. Formation of submarine gullies by turbidity currents. *EOS Transactions, American Geophysical Union, San Francisco, CA., December (invited)*.

Fedele, J.J. and Garcia, M.H. 2000. Generation of gullies on continental slopes by turbidity flows. *EOS Transactions, American Geophysical Union, Ocean Sciences, San Antonio, TX., January (poster)*.

Fedele, J.J. and Garcia, M.H. 2000. Alluvial roughness in streams with dunes: a boundary-layer approach. In *River, Estuarine, and Coastal Morphodynamics*, G. Seminara (editor), to be published by Springer-Verlag.


Gira, E. 2000. Study on the formation of gullies in continental margins by turbidity currents. Undergraduate Thesis, Ecole des Mines de Nancy, France (in French).

Huang, X., and Garcia, M.H., 2000. Pollution of gravel spawning grounds by deposition of suspended sediment. *Journal of Environmental Engineering, ASCE*, October issue.


Huang, X. and Garcia, M.H., 1999. Modeling of non-hydroplaning mudflows on continental slopes. *Marine Geology* 154, 131-142.

Lopez, F. and Garcia, M.H., 2000. Risk of sediment erosion and suspension in turbulent flows. *Journal of Hydraulic Engineering, ASCE*, accepted for publication.

Parsons, J. D. and Garcia, M. H. 2000. Enhanced sediment scavenging due to double-diffusive convection. *Journal of Sedimentary Research*, vol. 70, p. 47-52.



*Figure 1. Fractal dimensions for turbidity and density current fronts (frontal roughness).*



*Figure 2. A characteristic velocity scale for frontal processes (instabilities) in turbidity currents is obtained from the slope of the fit.*