

Theoretical and Numerical Studies of Variability and Predictability in an Unsteady Ocean

Geoffrey K. Vallis

AOS Program, Princeton University, NJ 08544

phone: (609) 258 6176 fax: (609) 258 2850 email: gkv@princeton.edu

Award #: N00014-99-1-0211

<http://www.gfdl.gov/~gkv>

LONG-TERM GOALS

Our long-term goals are to understand the nature of variability within the ocean, in particular that due to the motion of mesoscale eddies and their interaction with and dependence on the general circulation.

OBJECTIVES

Our particular objectives are to understand the nature and predictability of mesoscale eddies in the ocean. This includes the mechanisms of their equilibration, their structure and energetics, and their dependence on the large-scale parameters set by the general circulation.

APPROACH

Our approach is to use a hierarchy of numerical and analytic models of the ocean circulation, of varying complexity. At the simplest level are linear quasi-geostrophic models of baroclinic instability. At the next level are nonlinear, eddy resolving quasi-geostrophic models in idealized domains. Finally, we employ eddy resolving primitive equation models in fairly realistic configuration and domain.

WORK COMPLETED

We have completed a sequence of integrations using a quasigeostrophic model to explore the equilibration properties of mesoscale eddies, with oceanically realistic stratification. In collaboration with GFDL scientists we have configured two primitive equation numerical models to perform eddy resolving integrations of the Southern Ocean, and some preliminary experiments have been performed.

RESULTS

Considerable progress has been made in understanding the mechanisms that determine the scale and equilibration of mesoscale eddies in the ocean. Application of the theory of geostrophic turbulence has led to analytic predictions of the scale and magnitude of such eddies. It has been shown that the effects of non-uniform stratification (i.e., the presence of a thermocline) leads to a changing of the energetic pathways of mesoscale eddies, with the preferential concentration of baroclinic energy at the scale of the first deformation radius, in some contrast to classical theory in which the energy cascades to the gravest horizontal and vertical scales.


Report Documentation Page

Form Approved
OMB No. 0704-0188


Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE SEP 2000	2. REPORT TYPE	3. DATES COVERED 00-00-2000 to 00-00-2000			
4. TITLE AND SUBTITLE Theoretical and Numerical Studies of Variability and Predictability in an Unsteady Ocean		5a. CONTRACT NUMBER			
		5b. GRANT NUMBER			
		5c. PROGRAM ELEMENT NUMBER			
6. AUTHOR(S)		5d. PROJECT NUMBER			
		5e. TASK NUMBER			
		5f. WORK UNIT NUMBER			
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) AOS Program, Princeton University, Princeton, NJ, 08544		8. PERFORMING ORGANIZATION REPORT NUMBER			
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)		10. SPONSOR/MONITOR'S ACRONYM(S)			
		11. SPONSOR/MONITOR'S REPORT NUMBER(S)			
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 4	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Iso-vorticity: uniform stratification



Iso-vorticity: thermocline-like stratification



Different vorticity structures produced by mesoscale eddies in the presence of a uniform stratification (top) and a realistic stratification (bottom). Note the surface trapping in the lower case. From Smith and Vallis (2000, J. Phys. Oceanogr., in press)

IMPACT/APPLICATIONS

The impact of this lies in understanding the nature of mesoscale energetics in the ocean, and its dependence on the large scale parameters of the oceanic general circulation. Results from the more realistic simulations will indicate the predictability of the ocean on the mesoscale and deformation scales.

TRANSITIONS

Our results have not yet appeared in print. But nevertheless they are being used to interpret the results of altimeter measurements and primitive equation numerical models.

RELATED PROJECTS

A related GFDL/NOAA funded project on modeling eddies provides an invaluable insight into eddies in the Southern Ocean, as well as the resources for the extensive primitive equation calculations. In particular, GFDL is providing all the computer time for the experiments we are doing here, at no cost to ONR. We continue to collaborate closely with GFDL scientists.

PUBLICATIONS

Smith, K. S. and G. K. Vallis. Scales and equilibration of mid-ocean eddies: Freely decaying flow. *J. Phys. Oceanogr.* (in press).

Schonbek, M. and Vallis, G. K. Energy Decay of Solutions to the Boussinesq , Primitive and Planetary Geostrophic Equations. *J. Math. Analysis & Applics.* 234, 457-481.

Vallis, G. K. Large-scale circulation and production of stratification: effects of wind, geometry and diffusion. *J. Phys. Oceanogr.*, 30, 933--954.