

# **Shallow Water MCM and ASW Using Off-Board, Autonomous Sensor Networks and Multistatic, Time-Reversal Acoustics**

PI: Henrik Schmidt  
Massachusetts Institute of Technology  
77 Massachusetts Avenue  
Room 5-204  
Cambridge, MA 02139  
Phone: (617) 253-5727 Fax: (617) 253-2350 Email: [henrik@mit.edu](mailto:henrik@mit.edu)

CoPI: David Battle  
Massachusetts Institute of Technology  
77 Massachusetts Avenue  
Room 5-204  
Cambridge, MA 02139  
Phone: (617) 324-1461 Fax: (617) 253-2350 Email: [dbattle@mit.edu](mailto:dbattle@mit.edu)

Award Number: N00014-04-1-0014  
<http://acoustics.mit.edu/SWAMSI/SWAMSI.html>

## **LONG-TERM GOALS**

To achieve robust multi-static detection and classification of mine-like objects using cooperative networks of virtual acoustic sensors.

## **OBJECTIVES**

To utilize high fidelity acoustic modeling of both scatterers and shallow-water environments to better understand and bound the limits of detectability for mine-like objects via autonomous networks of sensors. To conduct a series of realistic experiments using multiple sonar-equipped AUVs in shallow water and then cross-validate the results obtained with high precision modeling and visualization. To better understand the problems of cooperative autonomous vehicle interaction to define the base-line infrastructure requirements for cooperative detection, classification and navigation.

## **APPROACH**

This program couples high accuracy acoustic modeling and visualization with customized AUV technology. The sonar sensing uses the bi-static and multi-static Synthetic Aperture created by the network, in combination with medium frequency (4-24 kHz) wide-beam insonification to provide coverage, bottom penetration and location resolution for concurrent detection, localization and classification of proud and buried targets in SW and VSW. The signal processing effort in SWAMSI is therefore centered around generalizing SAS processing to bi-static and multi-static configurations, including bi-static generalizations of auto-focusing and track-before-detect (TBD) algorithms. Another issue concerns the stability and coherence of surface and seabed multiples and their potential use in advanced medium-frequency sonar concepts.

MIT's acoustic modeling capabilities derive from both the SEALAB suite (VASA Associates) for general shallow water acoustics and FEMLAB (COMSOL Inc) for detailed structural acoustics and

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>30 SEP 2006</b>		2. REPORT TYPE		3. DATES COVERED <b>00-00-2006 to 00-00-2006</b>	
4. TITLE AND SUBTITLE <b>Shallow Water MCM and ASW Using Off-Board, Autonomous Sensor Networks and Multistatic, Time-Reversal Acoustics</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge, MA, 02139</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>Same as Report (SAR)</b>	18. NUMBER OF PAGES <b>7</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

target modeling. SEALAB incorporates the OASES environmental acoustic modeling framework developed at MIT [1,3], which is a widely distributed suite of models covering a variety of ocean waveguide and source/receiver representations. Recent developments are computational modules for full wave theory modeling of mono-static and bi-static target scattering and reverberation in shallow water waveguides. The most recently developed module, OASES-3D provides wave-theory modeling of the full 3-D acoustic environment associated with mono-static and bi-static configurations in SW and VSW with aspect-dependent targets and reverberation features [2,3]. It incorporates environmental acoustic features specifically associated with bi-static sonar concepts in shallow water, including aspect-dependent target models, seabed porosity, and scattering from anisotropic seabed roughness such as sand ripples.

With every major AUV deployment, the Mission Oriented Operating Suite (MOOS) previously created at MIT by research engineer Paul Newman advances in robustness and flexibility, and has been undergoing major upgrades in regard to the behavior-based control using the new IvP-Helm developed by Mike Benjamin of NUWC, who works closely with the MIT team as a Visiting Scientist. Another significant component is the development of a comprehensive simulation testbed, coupling the MOOS-IvP autonomous vehicle simulation environment with the SEALAB high-fidelity acoustic simulator, resulting in a complete, distributed software base for planning, simulating and analyzing multi-vehicle MCM missions.



## **WORK COMPLETED**

### **OASES3D Target Scattering Model**

Work has continued on developing a robust and efficient numerical approach for coupling the target scattering models with general propagation models handling the complex ocean waveguide physics. The virtual source approach developed in FY04 [4] provides an extremely efficient approach to coupling wavenumber integration codes, such as OASES3D, or normal mode codes (CSNAP, KRAKEN) with high fidelity target models such as the FESTA and FEMLAB FEM codes. By using a 3D spectral representation of the waveguide Green's function for the virtual sources, the expanded OASES3D modeling framework now handles complex target shapes and compositions, and in addition handles partially or completely buried targets, incorporating multiple scattering effects. In FY06 we have continued to use this new modeling capability has been applied to quantify the multiple scattering effects, and investigate the effect of target burial depth.

A specific modeling development has been performed in a joint effort between MIT and MPL-SIO, aimed at simulating time-reversal mirror scenarios for target DCL. The modeling concept, illustrated in Fig. 1, uses OASES to simulate the broadband insonification of the target by an array of individual sources in a source-receiver array (SRA). The insonification is then convolved with the target target response using OASES-3D, and the field received on the SRA is simulated. This produces a 'matrix' of Green's functions to and from all elements of the SRA, which can then be applied to perform 'synthetic' iterative time-reversal to enhance the target response over the background reverberation.

$$r_{ij}(t) = h_i(t) * s_{ij}(t) * h_j(t) + n(t)$$




**Figure 1. Principle of synthetic, iterative time reversal modeling.**

## EVA'06 Experiment


A major effort was undertaken in FY06 in collaboration with NURC on the planning of the EVA'06 experiment in October 2006 at Marciana Marina, Italy. The objective of this experiment is to provide an extensive 3-dimensional data set for bi-static scattering from proud, and partially and completely buried complex targets. Thus, the experiment will use the TOPAS parametric source for controlled insonification of the targets, in combination with an extensive receiver suite, including a small bi-static array synthetic aperture below the R/V Leonardo, and a dome array deployed by NRL-Stennis allowing for 3-D measurement of the scattering in the near-field of the targets. In addition to providing a unique data set for validation of the new target scattering models developed jointly between MIT and NURC, the experiment will also explore the possibility of using synthetic-aperture, time-reversal acoustics for target classification.

## RESULTS



***Figure 2 Spectral response enhancement achieved by synthetic, iterative time-reversal for a flush-buried spherical target, using a vertical; source-receiver array 200 m from the target, buried in 100 m water.***

In collaboration with MPL-SIO (Karim Sabra and W.A. Kuperman) an extensive simulation study has been performed regarding the potential of using synthetic, iterative time-reversal for enhancing the resonances of manmade, buried targets. The study has been performed using the new OASES-3D TR modeling framework described above, and the preliminary results are encouraging. Thus, Figure shows the performance of the approach for a flush-buried, 1 m diameter, spherical shell (GOATS sphere). The red symbols show the frequency response of the spherical shell, compared with the response of a spherical cavity of the same size, shown in blue, and showing no resonance structure. The study showed that the enhancement for this scenario will allow detection of the flush-buried sphere at signal-to-reverberation ratios as low as 0 dB.



**Figure 3. Back-propagated field focusing on flush-buried spherical target. Left frame shows the result for a spherical cavity, where the focusing highlights the specular scattering point. The frame to the right shows the equivalent result for a spherical shell (GOATS sphere), where the strong focusing on the lower part of the shell confirms the hypothesis [9] that the high-frequency scattering from a flush-buried spherical shell is dominated by the circumferential flexural wave, excited by the evanescent incident field, and re-radiated supersonically into the sediment and back into the water column at super-critical angles.**

The study also confirmed an earlier hypothesis regarding the strong high-frequency scattering anomaly observed for a flush-buried sphere in the GOATS experiments, suggesting that it is due to strong coupling at sub-critical incidence into the flexural circumferential, flexural waves in the shell, which are supersonic above the coincidence frequency, and therefore re-radiate from the lower portion of the shell, through the sediment and back to the water column, at supercritical angles [9]. Thus, the new TR model was used to back-propagate the field received on the SRA, and as seen in Fig. 3 the focusing at the lower part of the sphere is evident, thus confirming the hypothesis, which also has been confirmed independently in [10] by analysing bistatic, horizontal array data collected during the GOATS'98 experiment.

## IMPACT/APPLICATIONS

The long-term impact of this effort is the development of new sonar concepts for VSW MCM, which take optimum advantage of mobility, autonomy and adaptivity. For example, bi-static and multi-static, medium-frequency sonar configurations are being explored for completely or partially proud or buried mines in shallow water, with the traditional high-resolution acoustic imaging being replaced by a 3-D acoustic field characterization as a combined detection and classification paradigm, exploring spatial and temporal characteristics which uniquely define the target and its environment.

## TRANSITIONS

The virtual source modeling approach developed under this project has been transitioned to NURC as part of the OASE3D target modeling framework. Here it is coupled to the FEMLAB finite element framework to allow modeling of complex elastic targets. It has also been transitioned to NUWC (J. Blottman), CSS (D. Burnett), and WSU (Marston) for the same purpose. It is currently

being integrated with the MIT-MCM simulation framework developed under GOATS (N00014-05-1-0255) for simulating autonomous, adaptive target classification [7,8].

## RELATED PROJECTS

This effort is closely related to the GOATS project, initiated as the GOATS'2000 Joint Research Project (JRP) with the NATO Undersea Research Centre (NURC), and continued at MIT under the GOATS'2005 grant (N00014-05-1-0255), funded jointly by ONR codes 321OA (Livingston), 321OE (Swan), and 321TS (Commander). The collaboration with NURC, is continued under the Hybrid Target Modeling and Focused Acoustic Field (FAF) Joint Research Projects (JRP).

## REFERENCES

1. Schmidt, H. "OASES V 2.7 User's Guide", Massachusetts Institute of Technology, 2000.
2. Schmidt, H. and Lee, J.Y., "Physics of 3-D scattering from rippled seabeds and buried targets in shallow water," *Journal of the Acoustical Society of America*, 105(3), 1605-1617, 1999.
3. Schmidt, H. "Three-dimensional scattering from buried targets in shallow water," *In: Bovio, E., Schmidt, H., editors. GOATS 2000, SACLANTCEN CP-46*, 2002.
4. Schmidt, H. "Virtual source approach to scattering from partially buried elastic targets", *In High Frequency Acoustics*, M.B. Porter, M. Siderius, W.A. Kuperman, Eds., American Institute of Physics, 2004. [pdf](#)
5. Zampolli, M, Jensen, F.B., and Tesei, A. "A Finite element and hybrid modeling tools for the detection and classification of buried objects in shallow water", *In Boundary Influences in High Frequency, Shallow Water Acoustics*, University of Bath Press, Bath, UK, 2005.
6. Schmidt, H., Maguer, A, Bovio, E., Fox, W.L.J., LePage, K., Pace, N.G., Hollett, R., Guerrini, P., Sletner, P.A., Michelozzi, E., Moran, B. and Grieve, R. "GOATS'98 – AUV Network Sonar Concepts for Shallow Water Mine Countermeasures," *SACLANT Undersea Research Centre*, Report SR-302, 1998. [pdf](#)
7. M. Montanari, J.R. Edwards, and H. Schmidt. "Autonomous Underwater Vehicle-based Concurrent Detection and Classification Concept Using Higher-order Spectral Analysis," *IEEE Journal of Oceanic Engineering*, 31(1), 188-199, 2006
8. Eickstedt, D.P., "Adaptive Sampling in Autonomous Marine Sensor Networks", PhD Thesis, Massachusetts Institute of Technology, June 2006.
9. Schmidt, H. "Bistatic scattering from buried targets in shallow water." *In: Bovio, E., Tyce, R., Schmidt, H., editors. GOATS 2000, SACLANTCEN CP-46*, pp. 361-375. 2001.
10. Lucifredi, I. and Schmidt, H., "Sub-critical Scattering from Buried Elastic Shells". *J. Acous.l Soc. Am*, 2006 (In Press).

## **PUBLICATIONS**

Eickstedt, D.P., “Adaptive Sampling in Autonomous Marine Sensor Networks”, PhD Thesis, Massachusetts Institute of Technology, June 2006.

M. Montanari, J.R. Edwards, and H. Schmidt. “Autonomous Underwater Vehicle-based Concurrent Detection and Classification Concept Using Higher-order Spectral Analysis,” *IEEE Journal of Oceanic Engineering*, 31(1), 188-199, 2006

Irena Lucifredi and Henrik Schmidt , “Sub-critical Scattering from Buried Elastic Shells”. *J. Acoustical Society of America*, 2006 (In Press).

## **HONORS/AWARDS/PRIZES**

Prof. Henrik Schmidt has been awarded the *Pioneers of Underwater Acoustics Medal* of the Acoustical Society of America, to be presented at the 150th Meeting of the ASA, Oct. 17-21, 2005 in Minneapolis, MN.