

Predictability and Dynamics of Geophysical Fluid Flows

Roger M. Samelson

College of Oceanic and Atmospheric Sciences, Oregon State University

104 COAS Admin Bldg, Corvallis, OR 97331-5503 USA

phone: (541) 737-4752 fax: (541) 737-2064 email: rsamelson@coas.oregonstate.edu

Award Number: N00014-06-1-0369 (continuation of N00014-98-1-0813)

<http://www.coas.oregonstate.edu/~faculty/samelson.html>

LONG-TERM GOALS

The long-term goal of the research in this project is to improve our ability to predict environmental conditions using dynamical models.

OBJECTIVES

The central objective of the research in this project is to understand the mathematical and physical connections between the bred-growing-mode and singular vector techniques recently developed for numerical weather prediction, the Lyapunov vectors and exponents of dynamical systems theory, and instability theories of geophysical fluid dynamics. The intent is to gain insight into fundamental mathematical and physical aspects of predictability in unstable (irregular, chaotic) continuous systems.

APPROACH

A combination of analytical and numerical methods are being used to study a variety of mathematical models of geophysical fluid flows.

WORK COMPLETED

Graduate student Christopher Wolfe, who has been supported by this grant, defended his thesis on this topic in September 2006 (Wolfe, 2006). A new, efficient method for recovering Lyapunov vectors from singular vectors has been discovered and tested (Wolfe and Samelson, 2006b). A singular vector analysis of disturbances to the strongly nonlinear baroclinic oscillation has been completed and is described in a manuscript that will soon be submitted (Wolfe and Samelson, 2006c). Under the previous grant (N00014-98-1-0813), the Floquet analysis of linear disturbances to a strongly nonlinear baroclinic wave-mean oscillation in a high-dimensional geophysical fluid model was carried out (Wolfe and Samelson, 2006a).

RESULTS

Under the previous grant, the stability of a time-periodic baroclinic wave-mean oscillation in a two-layer quasi-geostrophic model was examined by computing a full set of time-dependent normal modes (Floquet vectors) for the oscillation (Wolfe and Samelson, 2006a). The Floquet vectors fell into two classes with direct physical interpretations: wave dynamical (WD) modes and damped-advective (DA) modes. The dynamics of the WD modes reflects the dynamics of the wave-mean oscillation, and these

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 2006		2. REPORT TYPE		3. DATES COVERED 00-00-2006 to 00-00-2006	
4. TITLE AND SUBTITLE Predictability and Dynamics of Geophysical Fluid Flows				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Oregon State University, College of Oceanic and Atmospheric Sciences, 104 COAS Admin Bldg, Corvallis, OR, 97331				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 5	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

modes are analogous to the normal modes of steady parallel flow. The DA modes represent instead a generalized solution to the damped advection problem. The leading Floquet exponents provide useful approximations to the leading Lyapunov exponents of the associated chaotic solution.

A new method for recovering Lyapunov vectors from asymptotic singular vectors has been discovered and tested (Wolfe and Samelson, 2006b). Lyapunov vectors are the fundamental time-dependent modes of instability of a flow with arbitrary time-dependence, and reduce to standard normal modes or Floquet modes, respectively, for steady or time-periodic basic flows. Standard methods require $N + 1$ singular vectors for the computation of any Lyapunov vector except the first, where N is the number of degrees of freedom of the model, which in ocean or atmosphere models is typically of order 105-107. The new method requires only $2n$ singular vectors for the computation of n Lyapunov vectors. Since the number n of leading Lyapunov vectors of interest is typically of order 10-102, this is a substantial savings; from a practical point of view, it makes calculations possible that previously were impossible. This method has been tested successfully on a weakly nonlinear model with 8 degrees of freedom (Figure 1) and on a strongly nonlinear model with 3840 degrees of freedom (Figure 2).

The analysis of the relationship between singular vectors and Floquet vectors in the context of a nonlinear baroclinic wave-mean oscillation was completed (Wolfe and Samelson, 2006c). It was found that the singular vectors divide into two dynamical classes which are related to those of the Floquet vectors. Singular vectors in the WD class were found to asymptotically approach constant linear combinations of Floquet vectors. The most rapidly decaying singular vectors projected strongly onto the most rapid decaying Floquet vectors. In contrast, the leading singular vectors projected strongly onto the leading adjoint Floquet vectors. Examination of trajectories near the basic cycle showed that the leading Floquet vectors were geometrically tangent to the local attractor, and so efficiently represented error variance associated with misplacement of the state on the attractor (Figure 3), while the leading initial singular vectors pointed off the local attractor, and so represented this error variance inefficiently (Figure 4). This illustrates that, for the standard linear and perfect-model hypotheses, the time-dependent normal modes (Lyapunov vectors) are a more appropriate basis for ensemble generation than singular vectors.

IMPACT/APPLICATIONS

The primary potential future impact of these results is on the design and use of ensemble forecasting techniques for the prediction of oceanic and atmospheric conditions.

RELATED PROJECTS

This project is a continuation of research that formed part of the ONR Predictability DRI. The research on dynamics and predictability is related to the NSF ITR project “Spatio-Temporal Complexity and Nonlinear Dynamics of Coastal Ocean Flows” (R. Samelson, J. Allen, and G. Egbert, Co-PIs).




Figure 1: One minus the pattern correlation between the second Lyapunov vector and the tangent vector to the nonlinear trajectory of the weakly nonlinear Phillips model vs. time t along the basic cycle. A value of unity indicates orthogonality while a value of zero indicates collinearity.




Figure 2. One minus the pattern correlation between the recovered Floquet vectors and the true Floquet vectors for three different optimization intervals (crosses, pluses, circles) vs. mode number. A value of unity indicates orthogonality while a value of zero indicates collinearity. Pattern correlations $< 10^{-8}$ are plotted at 10^{-8} to show the upper values more clearly.




Figure 3. Fraction of the initial error variance explained by the leading Floquet vectors (vertical axis) vs. maximum error bound. Lines give bin averages and errorbars (plotted on the $n = 5$ line) give bin standard deviations. The bin size is 0.005. The number of Floquet vectors n used to construct the leading subspace increases vertically, starting with $n = 1$ and ending with $n = 10$.




Figure 4. Fraction of the initial error variance explained by the leading singular vectors (vertical axis) vs. maximum error bound. The bin size is 0.005. The number of singular vectors n used to construct the leading subspace increases vertically, starting with $n = 1$ and ending with $n = 10$. All values are less than 0.05.

REFERENCES

Wolfe, C. L., and Samelson, R. M., 2006a. Normal mode analysis of a baroclinic wave-mean oscillation. *Journal of the Atmospheric Sciences*.

PUBLICATIONS

Wolfe, C. L., 2006. Quantifying linear disturbance growth in periodic and aperiodic systems. Ph.D. thesis, Oregon State University.

Wolfe, C. L., and R. M. Samelson, 2006b. Recovering Lyapunov vectors from singular vectors. *Tellus* [submitted]

Wolfe, C. L., and Samelson, R. M., 2006c. Singular vectors and time-dependent normal modes of a baroclinic wave-mean oscillation. *Journal of the Atmospheric Sciences*. [to be submitted]