

Award Number: W81XWH-11-1-0152

TITLE: Vitamin D Pathway Status and the Identification of Target Genes in the Mouse Mammary Gland

PRINCIPAL INVESTIGATOR: Dr. Donald Matthews

CONTRACTING ORGANIZATION: State University of New York at Albany, Rensselaer, New York, 12144

REPORT DATE: November 2014

TYPE OF REPORT: Final Report

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. **PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.**

1. REPORT DATE November 2014		2. REPORT TYPE Final Report		3. DATES COVERED 1 Jan 2011 – 31 Nov 2014	
4. TITLE AND SUBTITLE Vitamin D Pathway Status and the Identification of Target Genes in the Mouse Mammary Gland				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-11-1-0152	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Donald Matthews M. Sc., JoEllen Welsh PhD E-Mail: dmatthews@albany.edu				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) State University of New York at Albany Rensselaer, New York, 12144				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Mammary gland samples were isolated from wild type, vitamin D receptor knockout (VDRKO) and 1 α hydroxylase knockout (1 α KO) female mice for whole mounts and paraffin embedding (inguinal) and for RNA and protein isolation (thoracic). Time points collected included 6-10wk old nulliparous, 9 and 16 days pregnancy, 5 and 10 days lactation and 3 and 6 days involution. All whole mounts were completed and showed increased branching during pregnancy in the VDRKO glands relative to wild type and 1 α KO glands. Paraffin embedded involution samples were stained with hematoxylin and eosin and showed an apparent decrease in alveolar breakdown in the first few days of involution in VDRKO and 1 α KO glands compared to wild type controls. Organ culture studies show that treatment of wild type and 1 α KO glands with 1,25 dihydroxyvitamin D decreases proliferation and retards tertiary branching. This is not the case in VDRKO glands. Gene expression analysis via microarray and qPCR provides only a glimpse into the complexities of the signaling involved in this process and is being complimented by ongoing protein analysis. These results suggest that 1 α KO mammary glands do not develop exclusively similar to VDRKO or wild type glands which verifies our need to complete the remainder of our studies to determine if the VDR is acting to control mammary gland development through the vitamin D pathway or through some other ligand or possibly without a ligand.					
15. SUBJECT TERMS Breast cancer, prevention, vitamin D, transgenic mice					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON USAMRMC
a. REPORT	b. ABSTRACT	c. THIS PAGE			19b. TELEPHONE NUMBER (include area code)
Unclassified	Unclassified	Unclassified	Unclassified	25	

Table of Contents

	<u>Page</u>
Introduction.....	1
Body.....	1-14
Key Research Accomplishments.....	15-16
Reportable Outcomes.....	16
Conclusion.....	16-17
References.....	18
Appendices.....	19-20
Annual Training Summary.....	21-22

Introduction

Epidemiological studies have shown a significant increase in cancer incidence in women who are vitamin D deficient (1;2). It was also shown that there is a correlation between breast cancer death rate and distance from the equator, where lesser exposure of direct sunlight leads to lower endogenous vitamin D levels. These studies suggest that optimal vitamin D status may contribute to prevention of breast cancer in human populations. The processes of proliferation, differentiation and apoptosis of epithelial cells in the mammary gland are carefully regulated by the rise and fall of several hormones which control the budding and extension of ductal branching, alveolar development and regression of these structures at distinct developmental stages. Elevation of these signals occurs at puberty to cause the first major growth of the ductal network. Aberrant growth of mammary gland epithelial tissue may occur as a result of failure of cellular safeguards through familial or somatic mutation, which can lead to the onset of cancer. 1,25-dihydroxyvitamin D (1,25D) is a key regulator of calcium homeostasis but is also a cellular growth regulator in numerous tissues including prostate, skin, colon and breast (3). In the body, vitamin D is converted to 25-hydroxyvitamin D (25D) which is then activated by the 1-alpha-hydroxylase (1 α OHase) enzyme to 1,25 D, which is the high affinity ligand for the transcriptional regulator vitamin D nuclear receptor (VDR) (4). The function of the VDR in calcium regulation has been studied extensively although many questions still remain as to its role in cellular proliferation, differentiation and apoptosis in individual tissues. Mechanisms of 1,25 D-VDR effects include inhibition of cyclin D1 (5;6), induction of cell cycle inhibitors p21 and p27 (7) and growth inhibitor TGF- β (8). 1,25 D promotes differentiation in breast cancer cells by upregulating β -casein (9;10). Also, induction of apoptosis has been observed in response to 1,25 D treatment in a number of cancer cell lines (including breast) due to generation of reactive oxygen species and subsequent release of the apoptosis-inducing cytochrome c (11). Downregulation of anti-apoptotic factor Bcl-2 and upregulation of pro-apoptotic factor Bax has also been observed in 1,25 D treated breast cancer cells *in vitro*. Few additional binding partners for the VDR have been established. β -catenin binds to the VDR while unliganded (without 1,25 D) (9). Prolonged activation of β -catenin can induce hair follicle tumors while loss of function of the VDR leads to alopecia (loss of hair) (12). This alopecia is not present in mice lacking 1 α OHase enzyme suggesting it may be driven by promiscuous or unliganded VDR activity. The studies detailed here will help to determine how the vitamin D pathway affects growth within the mammary gland and whether the increased growth previously observed in mammary glands lacking the VDR is due to VDR specifically or the dysfunction of the vitamin D pathway. By using transgenic mouse models, these studies will provide valuable mammary gland contextual data that cell culture cannot provide.

Body

***In vivo* analysis of mammary gland development**

(Subtask 1) Scheduling the pairings of our breeding colonies (to be completed by months 1-9) took longer than anticipated as we had the unexpected issue of smaller than normal pup size in all of our colonies (figure 1). This was a concern since mammary gland development during puberty is linked to post-natal growth (13). After analysis of the growth of these pups, it was determined that we would

use the larger females at a later age for our study. Once this issue was addressed, the breeding and collection of samples went as anticipated and was finished at the beginning of month 12. We took mammary glands at the following time points: 4-10 week old nulliparous (puberty), 9 and 16 days pregnancy, 5 and 10 days post partum (lactation) and 3 and 6 days involution. This was done for all colonies (WT, VDRKO, 1 α HaseKO (1 α KO)).




Figure 1. Weights of female weanlings were more variable and overall lower than anticipated.

(Subtask 2) Whole mounts were made from the right inguinal mammary glands of each of these mice (months 1-9), using a carmine alum stain. These stains were completed by the end of month 12. (Subtask 3) The ductal outgrowth of these glands was measured (months 3-5) and completed by the end of month 12 (figure 2). (Subtask 4) Terminal end bud formation was analyzed in pregnant mammary glands (months 4-6) and completed by the end of month 12 (figure 3).




Figure 2. Whole mounts of mammary glands from 7wk old (A) WT, (B) 1 α KO, (C) VDRKO female mice. Ductal outgrowth in VDRKO mice were shown to be higher than in WT controls.




Figure 3. Whole mounts of mammary glands from 9 days (top) and 16 days (bottom) pregnant (A,D) WT, (B,E) 1 α KO, (C,F) VDRKO female mice. Tertiary branching is more extensive in the VDRKO glands compared to the WT and 1 α KO glands.

(Subtask 5) All left inguinal glands from each mouse were fixed in a 4% formalin solution overnight, processed in our tissue processor and embedded in blocks of paraffin wax (months 1-9). This was completed by the end of month 12. All blocks were sectioned at 5 microns and stained with hematoxylin and eosin (H&E) (figure 4). (Subtask 6) H&E stained sections for pregnancy and involution samples were analyzed for epithelial cell density (months 6-10) and was done so by the end of month 14 (figure 5).




Figure 4. Hematoxylin and eosin (H&E) stained sections from (A,D) WT, (B,E) 1 α KO, (C,F) VDRKO glands at 3 days involution (top) and 6 days involution (bottom). VDRKO and 1 α KO mammary glands at 3 days involution consist of persistent larger alveoli while their wild type counterparts present much smaller alveoli, suggesting decreased sensitivity to post weaning signaling in those glands. This difference does not appear to be present in glands 6 days post weaning.




Figure 5. Quantification of hematoxylin and eosin (H&E) stained sections from WT, VDRKO and 1 α KO glands at 9 and 16 days pregnancy (Timed Mating (TM)) and 3 and 6 days involution illustrates the growth of epithelial tissue during pregnancy and the regression of epithelial cell structures during involution. These data show that the VDRKO glands possess more epithelial tissue during 9 and 16 days pregnancy (p-values 0.044 and 0.036) and 3 days involution (p=0.002) There is no difference in the amount of epithelial tissue in the 1 α KO mammary glands compared to the WT glands, suggesting similar control of epithelial cell growth and regression in these glands. (asterisks indicate statistical difference to the WT group at that time point calculated by student's t-test).

(Subtask 7) Assessment of proliferation was performed using Ki67 immunofluorescent staining of paraffin embedded sections (months 30-42). We have restricted this analysis to the animals at 9 and 16 days of pregnancy as this is the most important phase during the pregnancy cycle in terms of epithelial cell proliferation. We observed no difference between the Ki67 positive cells of each strain on the 9th day of pregnancy. At 16 days pregnancy the percentage of positive Ki67 cells doubled in both WT and 1 α KO glands, however the VDRKO inguinal glands did not demonstrate the same increase and was significantly lower than that of the other two strains (figure 6). (Subtask 8) Assessment of apoptosis using TUNEL staining of sections (months 8-10) has been completed. We looked specifically at apoptosis in involution samples, as this is when apoptosis is most relevant. A protocol change was implemented for the BrdU and TUNEL staining of sections. We believed we could achieve a more accurate quantification of positive cells with a fluorescent-tagged antibody rather than the colorimetric approach. We have collected the data for the TUNEL assay quantification which is illustrated in figure 7. A 2-fold increase in apoptotic signaling in our two knockout models compared to the wild type controls was observed at 3 days involution. This was not seen at 6 days involution as no statistical difference was shown.




Figure 6. Quantification of fluorescent Ki67-stained sections from WT, VDRKO and 1αKO glands at 9 and 16 days pregnancy illustrates the proliferation of epithelial cells during pregnancy. The data shows an increase in proliferation from 9 to 16 days pregnancy in both WT and 1αKO glands that is not present in VDRKO glands, suggesting there are similarities in proliferatory signaling between the WT and 1αKO glands that is different from that of glands lacking VDR. (p=0.0005). (letters indicate statistical difference calculated by ANOVA)




Figure 7. Quantification of fluorescent TUNEL-stained sections from WT, VDRKO and 1αKO glands at 3 and 6 days involution illustrates the regression of epithelial cell structures during involution as a measure of apoptosis. The data shows that there is an increased amount of apoptosis in the VDRKO and 1αKO glands during the 3rd day of involution (p-values 0.005 and 0.003), suggesting there are similarities in apoptotic signaling in these glands. (asterisks indicate statistical difference to the WT group at that time point calculated by student's t-test).

(Subtask 9) Isolation of RNA from thoracic mammary glands (months 10-12) has been completed and qPCR for each stage of pregnancy was performed. An increase in VDR expression was observed in the 1 α KO glands on the 9th day of pregnancy. (Subtask 10) This coincided with an increase of p21 and cyclin D1 expression (figure 8). In glands lacking the VDR, an observed increase in cyclin D1 mRNA and wild type levels of p21, demonstrates the complex signaling of the vitamin D pathway. At 16 days pregnancy, VDR expression increased 2-3 fold compared to levels at 9 days pregnancy but both p21 and cyclin D1 levels decrease and result in similar expression in all three models. At 5 days lactation, VDR levels in 1 α KO mammary glands were elevated, as was the marker of differentiation TGF- β but not β -casein (figure 9). TGF- β expression was also elevated in VDRKO glands but β -casein was not statistically increased. 5 days later, VDR and TGF- β expression was again increased in 1 α KO samples compared to their wild type counterparts, this time with the addition of β -casein (figure 10). To analyze apoptotic signaling during involution, we chose to observe pro-apoptotic bax and anti-apoptotic bcl2 expression levels. At 3 days involution, none of the genes measured in our 2 knockout models (VDR, cyp24, bax, bcl2) had levels significantly different than that of the wild type glands other than the obvious lack of VDR in the VDRKO group. Interestingly, in the 1 α KO glands, both bax and bcl2 levels were elevated at 6 days involution as well as cyp24 (figure 11) compared to that of the wild type controls.




Figure 8. Gene expression analysis of WT, VDRKO and 1 α KO glands at 9 days pregnancy shows an increase of VDR levels in the 1 α KO glands ($p=0.004$) which coincides with increased expression of p21 ($p=0.0002$) and cyclin D1 mRNA ($p=0.0001$). Glands lacking the VDR still demonstrate an increase in cyclin D1 expression at this time ($p=0.003$) but lack an increase in p21 shown in the 1 α KO glands. Genes normalized to epithelial marker cytokeratin 18. (asterisks indicate statistical difference to the WT group for that gene calculated by student's t-test)




Figure 9. Gene expression analysis of WT, VDRKO and 1αKO glands at 5 days lactation shows an increase of VDR levels in the 1αKO glands ($p=0.007$) which coincides with increased expression of TGF-β mRNA ($p=0.010$). Glands lacking the VDR still demonstrate an increase in TGF-β ($p=0.010$). An increase in β-casein levels was found in 1αKO glands ($p=0.038$). (asterisks indicate statistical difference to the WT group for that gene calculated by student's t-test)




Figure 10. Gene expression analysis of WT, VDRKO and 1αKO glands at 10 days lactation shows an increase of VDR levels in the 1αKO glands ($p=0.008$) which coincides with increased expression of TGF-β ($p=0.006$) and β-casein mRNA ($p=0.031$). Glands lacking the VDR show an increase in TGF-β mRNA ($p=0.013$) but no statistical difference in β-casein levels. (asterisks indicate statistical difference to the WT group for that gene calculated by student's t-test)




Figure 11. Gene expression analysis of WT, VDRKO and 1αKO glands at 6 days involution shows no increase of VDR levels in the 1αKO glands but has an increase in the vitamin D pathway self-regulating cyp24 gene mRNA ($p=0.032$). 1αKO glands also have elevated levels of pro and anti-apoptotic genes, bax ($p=0.007$) and bcl2 ($p<0.001$). Glands lacking the VDR show no statistical difference in cyp24, bax or bcl2 levels. (asterisks indicate statistical difference to the WT group for that gene calculated by student's t-test)

(Subtask 11) Isolation of protein from thoracic mammary glands has been completed (month 12). (Subtask 12) Immunohistochemistry was implemented in the place of western blot analysis due to the heterogeneous nature of the mammary gland. The importance of analyzing proteins in epithelial cells specifically can be more easily studied using fluorescence microscopy and imaging software. Immunofluorescent stains were conducted for VDR, and for cyclin D1 and p21 to further analyze control of proliferation during pregnancy in our models (month 30). We were not able to find an effective antibody to analyze TGF-β1 as we had hoped. VDR protein expression during lactation does not appear to increase in 1αKO epithelial cells compared to WT cells as the mRNA analysis illustrates. Even though most epithelial cells are positive for VDR at 5 days lactation, the 1αKO glands possess a slightly smaller percentage of positive cells than do WT glands (79.8% vs 89.0% respectively, $p=0.044$). At 9 days pregnancy, there appears to be slightly fewer p21 and Cyclin D1 positive epithelial cells in the 1αKO glands compared to WT glands but due to a large amount of biological variability, the differences are not significant ($p=0.099$ and $p=0.057$ respectively). Even though we see a large number of p21 positive cells in all strains, similar to that of cyclin D1 at this time point, the higher intensity of the cyclin D1 staining implies a much higher concentration of cyclin D1 than p21. These results are unsurprising due to the necessity for proliferation of epithelial cells and elevated ductal branching during this stage of pregnancy.

(Subtask 13) Annual results were presented in our departmental seminar. (Subtask 14) The manuscript for *in vivo* experiments is still in progress.

Ex vivo analysis of mammary gland development and treatment with vitamin D metabolites

(Subtask 1) During the delay in the initial setup of the mouse pairings for task 1, we optimized the organ culture conditions (month 13) earlier than expected. (Subtask 2) We maintained hormone pellet implantation for 2 weeks and cultured the glands for 7 days (figure 12) (Subtask 3) while treating with 100nM 1,25-dihydroxyvitamin D, 500nM 25-hydroxyvitamin D or ethanol control.




Figure 12. Whole mounts of organ cultured mammary glands illustrating the effects of hormone supplementation *ex vivo*. Wild type mice were used ((A,D) control at day 0) and glands were grown in culture for (B,E) 1 day and (C,F) 7 days in media containing a hormone cocktail. Top pictures were taken at 3.5x magnification. Bottom pictures were taken at 16x magnification.

(Subtask 4) Whole mount observations (months 15-30) of *ex vivo* mammary glands concluded that treatment of wild type and 1 α KO glands with 1,25D impeded the ability of the ductal branching network to form a large proportion of their secondary and tertiary branches (figure 13). Treatment with 100nM 1,25D did not have a dramatic impact on VDRKO mammary gland branching. Also, treatment with 500nM 25D did not appear to have an effect on the branching of any of the three models. (Subtask 5) Ductal outgrowth analysis demonstrated the impact of 1,25D on the different genotypes, resulting in the significantly reduced branching extension of the wild type and 1 α KO ductal networks compared to the VDRKO (figure 14). In addition, it was determined that the lymph nodes within the inguinal mammary glands of VDRKO mice are larger than those of wild type or 1 α KO glands (figure 15). (Subtask 6) Terminal end bud formation quantitative analysis does not appear to be feasible with the presently implemented technology. The use of ImageJ software to quantify any changes in epithelial tissue area within parts of the whole mounted glands proved too difficult even with the use of dynamic thresholding. Use of the Metamorph software to more specifically quantify tertiary branching instead of overall epithelial cell density is currently being tested as a replacement. (months 16-31).




Figure 13. Mammary glands were treated in culture with 500nM 25D, 100nM 1,25D or ethanol control. Shown are wild type glands treated with (A) ethanol and (B) 1,25D. Treatment with the 1,25D prevented the secondary and tertiary branching structures (arrows) that would normally form upon signaling from the hormone cocktail in culture.




Figure 14. Ductal outgrowth analysis of WT, VDRKO and 1 α KO mammary glands grown in culture and treated with 500nM 25D, 100nM 1,25D or ethanol control. Treatment of glands with the active metabolite of vitamin D, 1,25D elicited a difference in the extension of the ductal branching network into the mammary fat pad between the VDRKO model and the 1 α KO and WT models ($p=0.001$). (letters indicate statistical difference to the WT group for that treatment calculated by ANOVA)




Figure 15. Measurements of the lymph node area of WT, VDRKO and 1αKO mammary glands grown in culture. Lymph nodes present in the inguinal glands of VDRKO mice are consistently larger than those from WT or 1αKO glands ($p=0.0003$). (letters indicate statistical difference to the WT group calculated by ANOVA)

(Subtask 7) Inguinal glands were formalin fixed, dehydrated and embedded in paraffin wax (months 15-30). These blocks were then sectioned and H&E stained. Analysis of H&E stained sections (months 16-31) to determine the amount of epithelial cells present in *ex vivo* glands demonstrated no differences between strains. Treatment of wild type glands with 25D shows a decrease in epithelial tissue area but we believe this to be an artifact of the tissue sectioning process (figure 16). (Subtask 8) Epithelial cell content in H&E stained sections were quantitated and suggest there was no difference in presence of epithelial cells due to treatment. Additional sections were subjected to immunohistochemical analysis for p21 as a downstream target of vitamin D signaling (months 16-42). Unfortunately, the variability in the samples proved too much for any statistical significance to be shown. Perhaps a larger sample size would alleviate this issue. (Subtask 9) Proliferating cells measured by the presence of Ki67 was used to determine the level of proliferation in each of our models and treatments. Treatment of inguinal glands in culture with 1,25D resulted in a decrease in Ki67 positive WT and 1αKO epithelial cells while VDRKO cells were unaffected by treatment (figure 17)

(Subtask 10) Immunofluorescent TUNEL assays were performed to elucidate the presence of apoptotic cells that resulted from vitamin D treatment (months 18-42). Statistical analysis showed no increase in apoptosis with 1,25D treatment. This was likely due to a high level of apoptosis observed in some control samples. A change in organ culture protocol will be necessary to prevent excessive apoptosis in control samples.




Figure 16. Analysis of epithelial cell area in H&E stained mammary gland sections from each strain of mouse treated with either 25D or 1,25D (or ethanol control). A statistically significant decrease was observed in the wild type mice with 25D treatment ($p=0.008$) but is likely due to sections not being in the same plane as the rest of the samples. (asterisks indicate statistical difference to the WT control group calculated by ANOVA)




Figure 17. Immunofluorescent analysis of Ki67 expression in WT, VDRKO and 1αKO glands in organ culture. The data show a decrease in Ki67 positive cells in both WT and 1αKO epithelial cells ($p=0.043$ and $p=0.049$ respectively). (asterisks indicate statistical difference to the control group for that strain of mouse, calculated by student's t-test)

(Subtasks 11-12) We have isolated RNA from all of our thoracic gland samples and have used qPCR to analyze a number of genes. No differences were found in a number of different genes analyzed, including cyclinD1, Bax, Bcl2 and Cyp24. Basal levels of p21 were found to be slightly elevated in control KO glands (VDRKO and 1 α KO) but were unaffected by treatment with either 25D or 1,25D (figure 18). Further gene expression analysis will be performed for additional targets in the future. (months 16-31). (Subtasks 13-14) Microarray analysis showed changes in gene expression upon treatment of WT glands with 1,25D in culture. The significantly modulated genes of interest are listed in Table 1. Protein was harvested from these thoracic glands (months 30-42) but due to further consideration of the heterogeneous nature of the mammary gland, immunohistochemistry was substituted for western blot analysis to analyze protein expression changes and was detailed in Subtask 8 (months 30-42).




Figure 18. Gene expression analysis of WT, VDRKO and 1 α KO glands in organ culture shows a basal increase of p21 levels in the VDRKO and 1 α KO glands ($p=0.002$ and $p=0.023$ respectively). (asterisks indicate statistical difference to the WT group for that gene calculated by student's t-test)

Table 1. Mouse exon expression analysis of WT 1,25D treated vs control mammary glands in organ culture. Listed are the statistically significant changes in expression of both canonical calcium regulatory genes and other possible genes of interest. (Fold change cutoff was set at 1.5-fold)

Calcium Regulatory Genes		
<u>Fold Change</u>	<u>Gene Symbol</u>	<u>Gene description</u>
1.6	Camk2a	calcium/calmodulin-dependent protein kinase II alpha
-1.51	Clca4	chloride channel calcium activated 4
-1.52	S100a4	S100 calcium binding protein A4
-1.55	Camta1	calmodulin binding transcription activator 1
-1.62	Calca	calcitonin/calcitonin-related polypeptide, alpha
-1.62	Casq2	calsequestrin 2
-1.88	Calml3	calmodulin-like 3
Other Genes of Interest		
<u>Fold Change</u>	<u>Gene Symbol</u>	<u>Gene Description</u>
1.72	Cyp24a1	cytochrome P450, family 24, subfamily a, polypeptide 1
1.63	Cited1	Cbp/p300-interacting transactivator w/ Glu/Asp-rich carboxy-terminal domain 1
1.56	Steap2	six transmembrane epithelial antigen of prostate 2
1.5	Lpin3	lipin 3
-2.61	Csn1s2b	casein alpha s2-like B

(Subtask 15) I will be presenting my new results in addition to the results from the ongoing organ culture studies in our annual departmental seminar in February 2015. I will also be defending my thesis shortly thereafter (Subtask 16) Upon completion of each task, a manuscript will be prepared and submitted for publication.

Key Research Accomplishments

Task 1

- An increase in ductal growth was observed in VDRKO glands during puberty, compared to WT and 1 α KO glands
- An increase in ductal branching was observed in VDRKO glands during pregnancy, compared to WT and 1 α KO glands
- A delay in involution at 3 days was observed in VDRKO and 1 α KO glands, demonstrated by persistent alveoli compared to WT glands
- Invasion of mammary gland tissue by stromal cells at 3 days involution appeared more extensive in VDRKO glands compared to WT and 1 α KO glands
- An increase in apoptotic cells at 3 days involution was observed in VDRKO and 1 α KO glands compared to WT glands
- VDR mRNA was elevated in 1 α KO glands during all stages except involution compared to VDRKO and WT glands
- Bax and Bcl2 mRNA was elevated in 1 α KO glands at 6 days involution compared to WT glands. VDRKO may have increased Bax mRNA levels at this time point (as described previously) but sample size may be too small
- An increase in cyclin D1 mRNA levels at 9 days pregnancy was observed in VDRKO and 1 α KO glands compared to WT glands. Interestingly, p21 mRNA levels in 1 α KO glands were also elevated during this time point compared to WT and VDRKO glands.
- An increase in TFG- β 1 mRNA at 5 days lactation was observed in VDRKO and 1 α KO glands compared to WT glands. An increase in β -casein mRNA was also observed in 1 α KO glands at 5 and 10 days lactation compared to WT glands
- At 5 days lactation, the number of VDR positive cells decreased slightly in 1 α KO glands compared to WT glands
- At 5 days pregnancy, both p21 and cyclin D1 positive cells appeared to decrease in 1 α KO glands, albeit not quite significantly (p=0.098 and p=0.056 respectively)
- The percentage of Ki67 positive cells increased from 9 to 16 days pregnancy in WT and 1 α KO glands but not VDRKO glands

Task 2

- Treatment of WT and 1 α KO glands with 1,25D in culture decreased ductal outgrowth and branching compared to VDRKO glands
- There was no effect on ductal morphology in any strain upon treatment with 25D
- VDRKO glands possessed larger lymph nodes than that of WT or 1 α KO glands
- There were no changes in epithelial tissue area within the glands of any strain in culture upon treatment with 25D or 1,25D
- Microarray analysis showed that treatment of WT glands with 1,25D induced changes in both calcium regulatory genes and non-canonical vitamin D pathway genes.

- qPCR analysis demonstrated that KO glands both showed a basal increase in p21 levels compared to WT glands.
- Treatment of WT and 1 α KO glands with 1,25D in culture decreased Ki67 positive cells to near zero. VDRKO glands were unaffected
- TUNEL analysis eluded to a possible increase in apoptosis upon treatment with 1,25D in all strains but variability in control samples were too high to reach statistical significance
- Analysis of p21 protein expression in organ culture proved difficult due to a high level of variability among the samples

Reportable Outcomes

Matthews DG et al. Vitamin D pathway status and the identification of target genes in the mouse mammary gland. AACR Special Conference: Advances in Breast Cancer Research. San Francisco, CA, Oct 12-15, 2011.

Matthews DG et al. Characterization of mammary gland development in vitamin D pathway ablated mice. 15th Vitamin D meeting. Houston Texas, June 20-22, 2012.

Conclusion

VDR mRNA levels were consistently higher throughout pregnancy and lactation in 1 α KO glands, perhaps as a mechanism to attempt to overcome a lack of available active vitamin D. Contrary to this is the decrease in VDR positive cells in the 1 α KO glands during lactation. This discrepancy may be due to the effectiveness of translation, a possible increase in overall VDR protein but decrease in the number of expressing epithelial cells or it could be a fundamental difference between thoracic and inguinal glands which a few previous studies have detected subtle differences. Analysis of whole mounts and H&E stained sections of 9 and 16 day pregnancy samples verify an increase in the presence of epithelial cells due to accelerated ductal branching in VDRKO glands compared to wild type and 1 α KO glands. This correlates with elevated cyclin D1 mRNA levels and suggests that the VDR acts independently of 1,25 dihydroxyvitamin D to mediate these results. Protein analysis shows no change in the percentage of cells containing cyclin D1. Interestingly, 1 α KO glands display elevated mRNA levels of both pro and anti-proliferative genes cyclin D1 and p21 at 9 days pregnancy, something not observed in the other two models. This may be due to an alternate function of the 1 α -hydroxylase enzyme, adding another layer of complexity to vitamin D pathway studies. The percentage of cells containing p21 protein remains unaffected by either transgenic knockout. During lactation, we observed an increase in TGF- β mRNA in both knockout models which indicates a stronger push towards differentiation in these glands and also illustrates some similarity between our vitamin D pathway aberrant models. Paraffin-embedded involution samples which were sectioned and H&E stained, illustrated a delayed apoptotic response in the 1 α KO glands, similar to what was previously observed in VDRKO glands in terms of alveolar

deconstruction 3 days into involution in some animals. However, we do observe an elevated level of apoptosis in both knockout models at this time point, which does not appear to be the result of bax/bcl2 regulation, possibly due to a lag in apoptotic signaling in these glands.

Organ culture studies have confirmed a more extensive branching network in VDRKO glands treated with 1,25D compared to the wild type and 1 α KO models, suggesting that 1 α KO glands are able to utilize vitamin D metabolites downstream of the ablated activating enzyme or possibly alternative VDR ligands which may include 25D or other vitamin D like steroids. Whole mount analysis has also demonstrated that treatment of glands with 1,25D will prevent a significant amount of the secondary and tertiary branching observed in the control glands under hormonal stimulation. Not surprisingly, immunohistochemistry demonstrated a decrease in proliferatory marker Ki67 upon 1,25D treatment of WT and 1 α KO glands but not VDRKO glands although an expected increase in apoptosis did not occur in any of our models. This could be due to our organ culture methodology leading to elevated levels of apoptosis in a large number of control samples. This may be alleviated using small pieces of tissue instead of the entire gland. It was our intention to observe the gland as a whole for more reproducible and accurate comparisons, however this may be to the detriment of the overall health of the organ in culture and possibly enhancing the issues of variability. An increase in sample size would also help limit the effects of biological variability. In addition, although analyzing the localization of the protein is interesting and insightful, knowing the total protein content would be a more effective validation of the qPCR data. We should not ignore our original assertion that taking total glandular protein for western blot analysis would lead to inaccurate results due to the number of other cell types but instead modify our existing protocol to include the digestion of each gland and specific isolation of epithelial cells which has been done in recent published studies. From here we intend to further examine the role of vitamin D pathway proteins on ductal branching, deconstruction of epithelial cell structures and associated signaling by analyzing the effects of the loss of vitamin D proteins within the stromal compartment of the mammary gland. This will allow us to determine the role of the vitamin D pathway within the adipose cells within the mammary tissue specifically and their effects on epithelial cells and overall mammary gland growth and development. This is important to ascertain as the stromal-epithelial interactions are complex and vital to the normal development and function of the gland. The completion of these studies will give us a better idea of how vitamin D affects epithelial cell growth in the mammary gland and provide insight into the correlation between vitamin D deficiency and an increase in breast cancer incidence.

References

- 1** Abbas S et al. Dietary vitamin D and calcium intake and premenopausal breast cancer risk in a German case-control study. *Nutr Cancer* 2007;59(1):54-61
- 2** Dizdar O et al. Vitamin D intake and breast cancer risk in postmenopausal women. *Arch Intern Med* 2007; 167(22):2532.
- 3** Zinser G et al. Vitamin D(3) receptor ablation alters mammary gland morphogenesis. *Development* 2002; 129(13):3067-3076.
- 4** Jensen T et al. The vitamin D3 receptor and retinoid X receptors in psoriatic skin: the receptor levels correlate with the receptor binding to DNA. *Br J Dermatol* 1998;138(2):225-228.
- 5** Shah S et al. The molecular basis of vitamin D receptor and beta-catenin crossregulation. *Mol Cell* 2006;21(6):799-809.
- 6** Schedlich LJ et al. Insulin-like growth factor binding protein-5 interacts with the vitamin D receptor and modulates the vitamin D response in osteoblasts. *Mol Endocrinol* 2007; 21(10):2378-2390.
- 7** Flanagan L et al. Efficacy of Vitamin D compounds to modulate estrogen receptor negative breast cancer growth and invasion. *J Steroid Biochem Mol Biol* 2003; 84(2-3):181-192.
- 8** Capiati DA et al. Inhibition of serum-stimulated mitogen activated protein kinase by 1alpha,25(OH)₂-vitamin D₃ in MCF-7 breast cancer cells. *J Cell Biochem* 2004;93(2):384-397.
- 9** Palmer HG et al. Vitamin D(3) promotes the differentiation of colon carcinoma cells by the induction of E-cadherin and the inhibition of beta-catenin signaling. *J Cell Biol* 2001; 154(2):369-387.
- 10** Wang Q et al. 1,25-dihydroxyvitamin D₃ and retinoic acid analogues induce differentiation in breast cancer cells with function- and cell-specific additive effects. *Breast Cancer Res Treat* 2001; 67(2):157-168.
- 11** Welsh J et al. Role of apoptosis in the growth inhibitory effects of vitamin D in MCF-7 cells. *Adv Exp Med Biol* 1995; 375:45-52.
- 12** Palmer HG et al. The vitamin D receptor is a Wnt effector that controls hair follicle differentiation and specifies tumor type in adult epidermis. *PLoS ONE* 2008; 3(1):e1483.
- 13** Anderson RR. Relation between mammary gland development and body weight. *Journal of Dairy Science* 1976; 59(8):1518-1521.

APPENDIX A

Copy of Abstract for AACR Advances in Breast Cancer Research Meeting (Oct 12-15, 2011)

Vitamin D pathway status and the identification of target genes in the mouse mammary gland

Donald G Matthews, Teresa M Lloyd-Coronado, Elizabeth Stevens, JoEllen Welsh

State University of New York at Albany, Cancer Research Center

Multiple epidemiological studies have demonstrated correlations between vitamin D status and incidence of cancers, including colon and breast. Our long term goal is to understand the mechanisms by which the vitamin D signaling pathway impacts on breast cancer. Upon absorption or endogenous synthesis, vitamin D is metabolized into 25-hydroxyvitamin D (25D) and 1,25-dihydroxyvitamin D (1,25D) in multiple tissues. 1,25D is a high-affinity ligand for the nuclear vitamin D receptor (VDR) that alters gene expression and inhibits growth of normal and transformed mammary cells. However, 25D and other steroid-like compounds such as bile acids, can also activate VDR *in vitro*. Although studies with VDR knockout (KO) mice have demonstrated that VDR affects proliferation, differentiation and apoptosis in the mammary gland, the specific ligands that trigger VDR signaling have yet to be identified. Generation of 1,25D from 25D is mediated by CYP27B1 which is present and developmentally regulated in the mammary gland. Studies with the VDRKO mouse have established that its phenotype is different from mice lacking CYP27B1, supporting the concept that other VDR ligands exist and/or the VDR may act in the absence of ligand. We are using animal models to determine whether VDR exerts 1,25D-independent effects on growth, apoptosis or differentiation of mammary epithelial cells *in vivo*. Glands from VDRKO and CYP27B1KO mice during puberty, pregnancy, lactation, involution and aging have been collected for analysis of gross morphology, histology, and genomic profiling in comparison to age-matched control mice. Our data indicate that mammary glands from VDRKO mice display accelerated branching, enhanced sensitivity to estrogen and progesterone and altered gene expression during puberty and early pregnancy. Surprisingly, initial studies indicate that similar changes are not observed in glands from CYP27B1KO mice, suggesting that other VDR ligands may substitute for 1,25D in control of mammary cell turnover *in vivo*. Alternatively, our results are consistent with the concept that unoccupied VDR may function in the mammary gland. To further explore the role of VDR and CYP27B1 in the glandular epithelial compartment, mice with mammary epithelial-specific deletion of VDR or CYP27B1 were generated by crossing MMTV-cre mice with mice carrying floxed alleles of VDR or CYP27B1. Quantitation of VDR and CYP27B1 expression in the mammary gland, and characterization of the phenotype of these cre-flox mice is in progress. Collectively, these studies have identified novel roles for the VDR and its ligands in mammary gland development that provide insight into the relationship between vitamin D status and breast cancer.

Copy of Abstract for 15th Vitamin D Workshop (June 20-22, 2012)

CHARACTERIZATION OF MAMMARY GLAND DEVELOPMENT IN VITAMIN D PATHWAY ABLATED MICE.

D.G. Matthews, T.M. Lloyd-Coronado, P. Gavett, E. Stevens, J. Welsh Department of Biomedical Sciences, State University of New York at Albany, Rensselaer, NY, USA 12144

Multiple epidemiological studies have demonstrated correlations between vitamin D status and incidence of cancers, including colon and breast. Our long term goal is to understand the mechanisms by which the vitamin D signaling pathway impacts on breast cancer. Generation of the high-affinity vitamin D receptor (VDR) ligand 1,25-dihydroxyvitamin D (1,25D) from 25-hydroxyvitamin D (25D) is mediated by CYP27B1 which is present and developmentally regulated in multiple tissues including the mammary gland. Studies with the VDR knockout (VDRKO) mouse have established that its phenotype is different from mice lacking CYP27B1, supporting the concept that other VDR ligands exist and/or the VDR may act in the absence of ligand. We are using animal models to determine whether VDR exerts 1,25D-independent effects on growth, apoptosis or differentiation of mammary epithelial cells *in vivo*. Glands from VDRKO and CYP27B1KO mice during puberty, pregnancy, lactation, involution and aging have been collected for analysis of gross morphology, histology, and genomic profiling in comparison to age-matched control mice. Our data indicate that mammary glands from VDRKO mice display accelerated branching, enhanced sensitivity to estrogen and progesterone and altered gene expression during puberty and early pregnancy. Surprisingly, initial studies indicate that similar changes are not observed in glands from CYP27B1KO mice although gene expression changes associated with cell cycle regulation (p21, cyclin D1), differentiation (TGF- β , β -casein) and apoptosis (Bax) have been noted. This suggests that other VDR ligands may substitute for 1,25D in control of mammary cell turnover *in vivo*. Alternatively, our results are consistent with the concept that unoccupied VDR may function in the mammary gland. Collectively, these studies have identified novel roles for the VDR and its ligands in mammary gland development that provide insight into the relationship between vitamin D status and breast cancer. Funded by DOD Predoctoral Fellowship Award #W81XWH-11-10152

Annual Training Summary

2011

April 6, 2011 – Topics in Cancer, Journal Club Presentation – Gave 1hr presentation on relevant and current cancer research. Paper presented – Trimboli AJ et al. *Pten* in Stromal Fibroblasts Suppresses Mammary Epithelial Tumors. *Nature* 2009;461(7267):1084-91.

October 13, 2011 – Poster Presentation – Presented current data to interested parties for 2hrs. Matthews DG et al. Vitamin D pathway status and the identification of target genes in the mouse mammary gland. AACR Advances in Breast Cancer Research: Genetics, Biology, and Clinical Applications meeting, San Francisco California.

November 2, 2011 – Topics in Cancer, Journal Club Presentation – Gave 1hr presentation on relevant and current cancer research. Paper presented – Lyons TR et al. Postpartum mammary gland involution drives progression of ductal carcinoma *in situ* through collagen and COX-2. *Nature Medicine* 2011;17(9):1109-15.

November 5, 2011 – Annual Departmental Seminar – Gave 1hr presentation on current research progress. Matthews DG. Vitamin D pathway status and the identification of target genes in the mouse mammary gland.

2012

March 27, 2012 – Topics in Cancer, Journal Club Presentation – Gave 1hr presentation on relevant and current cancer research. Paper presented - Png KJ et al. A microRNA regulon that mediates endothelial recruitment and metastasis by cancer cells. *Nature* 2011;481(7380):190-4.

May 18, 2012 – Annual Departmental Seminar – Gave 1hr presentation on current research progress. Matthews DG. Vitamin D pathway status and the identification of target genes in the mouse mammary gland.

October 3, 2012 – Topics in Cancer, Journal Club Presentation – Gave 1hr presentation on relevant and current cancer research. Paper presented - Li J et al. Treatment of breast cancer stem cells with oncolytic herpes simplex virus. *Cancer Gene Therapy* 2012;19(10):707-14.

June 21, 2012 – Poster Presentation – Presented current data to interested parties for 2hrs. Matthews DG et al. Vitamin D pathway status and the identification of target genes in the mouse mammary gland. 15th Workshop on Vitamin D, Houston Texas.

2013

February 27, 2013 – Topics in Cancer, Journal Club Presentation – Gave 1hr presentation on relevant and current cancer research. Paper presented – Yin G et al. Constitutive proteasomal degradation of TWIST-1 in epithelial-ovarian cancer stem cells impacts differentiation and metastatic potential. *Oncogene* 2013; 32, 39-49.

October 16, 2013 - Topics in Cancer, Journal Club Presentation – Gave 1hr presentation on relevant and current cancer research. Paper presented – Werbeck JL et al. Tumor Microenvironment Regulates Metastasis and Metastasis Genes of Mouse MMTV-PymT Mammary Cancer Cells In Vivo. *Veterinary Pathology* 2013 Oct 3, Epub.

2014

April 23, 2014 – Topics in Cancer, Journal Club Presentation – Gave 1hr presentation on relevant and current cancer research. Paper presented – Volinia S and Croce CM. Prognostic microRNA/mRNA signature from the integrated analysis of patients with invasive breast cancer. *PNAS* 2013 April 30; 110, 7413-7417.

December 9, 2014 - Topics in Cancer, Journal Club Presentation – Gave 1hr presentation on relevant and current cancer research. Paper presented – Ning H et al. Conversion of Adipose-Derived Stem Cells into Natural Killer-Like Cells with Anti-Tumor Activities in Nude Mice. *PLOS One* 2014 August; 9, epub.