

C4d Deposits on the Surface of RBCs in Trauma Patients and Interferes With Their Function*

Takashi Muroya, MD¹; Lakshmi Kannan, PhD¹; Ionita C. Ghiran, MD¹;
Sergey S. Shevkopyas, PhD²; Ziv Paz, MD, MPH¹; Maria Tsokos, MD¹;
Jurandir J. Dalle Lucca, MD, PhD³; Nathan I. Shapiro, MD, MPH¹; George C. Tsokos, MD¹

Objective: Complement system is activated in patients with trauma. Although complement activation is presumed to contribute to organ damage and constitutional symptoms, little is known about the involved mechanisms. Because complement components may deposit on RBCs, we asked whether complement deposits on the surface of RBC in trauma and whether such deposition alters RBC function.

Design: A prospective experimental study.

Setting: Research laboratory.

Subjects: Blood samples collected from 42 trauma patients and 21 healthy donors.

Intervention: None.

Measurements and Main Results: RBC and sera were collected from trauma patients and control donors. RBCs from trauma patients ($n = 40$) were found to display significantly higher amounts of C4d on their surface by flow cytometry compared with RBCs from control ($n = 17$) ($p < 0.01$). Increased amounts of iC3b were found in

trauma sera ($n = 27$) (vs 12 controls, $p < 0.01$) by enzyme-linked immunosorbent assay. Incubation of RBC from universal donors (type O, Rh negative) with trauma sera ($n = 10$) promoted C4d deposition on their surface (vs six controls, $p < 0.05$). Complement-decorated RBC ($n = 6$) displayed limited their deformability (vs six controls, $p < 0.05$) in two-dimensional microchannel arrays. Incubation of RBC with trauma sera ($n = 10$) promoted the phosphorylation of band 3, a cytoskeletal protein important for the function of the RBC membrane (vs eight controls, $p < 0.05$), and also accelerated calcium influx ($n = 9$) and enhanced nitric oxide production ($n = 12$) (vs four and eight controls respectively, $p < 0.05$) in flow cytometry.

Conclusions: Our study found the presence of extensive complement activation in trauma patients and presents new evidence in support of the hypothesis that complement activation products deposit on the surface of RBC. Such deposition could limit RBC deformability and promote the production of nitric oxide. Our findings suggest that RBC in trauma patients malfunctions, which may explain organ damage and constitutional symptoms that is not accounted for otherwise by previously known pathophysiologic mechanisms. (*Crit Care Med* 2014; 42:e364–e372)

Key Words: C4d; complement; deformability; nitric oxide; red blood cell; trauma

*See also p. 1323.

¹Department of Medicine, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, MA.

²Department of Biomedical Engineering, Tulane University, New Orleans, LA.

³Institute of Surgical Research, San Antonio, TX.

This work was performed at Beth Israel Deaconess Medical Center, Harvard Medical School.

Supported, in part, by grant W81XWH-12-1-0001 from Medical Research and Materiel Command, Fort Detrick, MD.

Dr. Ghiran was supported by the National Institutes of Health (NIH) (R01HL096795). He and his institution received support for article research from the NIH (HL096795). Dr. Shevkopyas was supported by a 2012 NIH Director's Transformative Research Award (R01HL117329). He received support for article research from NIH and served as a board member for the Cumberland Data Safety Monitoring Board. His institution received grant support from the NIH, Department of Defense, Thermo-Fisher, Rapid Pathogen Screening, and Cheetah Medical. Dr. George C. Tsokos and his institution received support for article research from the United States Medical Research and Materiel Command (W81XWH-12-1-0001). The remaining authors have disclosed that they do not have any potential conflicts of interest.

For information regarding this article, E-mail: gtsokos@bidmc.harvard.edu
Copyright © 2014 by the Society of Critical Care Medicine and Lippincott Williams & Wilkins

DOI: 10.1097/CCM.0000000000000231

The clinical significance of trauma cannot be underestimated with over 10 million car accidents occurring and over 35 thousand people dying each year in the United States (1). Trauma resulting from accidents or unintentional injuries is the first leading cause of death for those under 50 years old and accounts for one out of about every 20 deaths in the United States (2).

Although trauma usually involves a certain anatomical injury involving extremities or the torso and is followed by obvious pathophysiologic events usually linked to blood loss, it is invariably associated with manifestations from organs not directly affected by injury. The origin of these secondary manifestations is poorly understood. There is information suggesting the presence of an acute inflammatory response in patients experiencing trauma. Specifically, there is evidence that the complement

Report Documentation Page

*Form Approved
OMB No. 0704-0188*

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE 01 MAY 2014		2. REPORT TYPE N/A		3. DATES COVERED -	
4. TITLE AND SUBTITLE C4d Deposits on the Surface of RBCs in Trauma Patients and Interferes With Their Function*				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Muroya T., Kannan L., Ghiran I. C., Shevkoplyas S. S., Paz Z., Tsokos M., Dalle Lucca J. J., Shapiro N. I., Tsokos G. C.,				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) United States Army Institute of Surgical Research, JBSA Fort Sam Houston, TX				8. PERFORMING ORGANIZATION REPORT NUMBER	
				10. SPONSOR/MONITOR'S ACRONYM(S)	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 9	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

system is activated and mediates early posttraumatic inflammatory response (3–5). In addition, the levels of proinflammatory cytokines, including tumor necrosis factor- α , interleukin (IL)-1, and IL-6, are increased in trauma (6, 7). It is suggested that complement activation and proinflammatory cytokine storm occurring in patients with trauma collectively choreograph systemic inflammatory response syndrome associated with acute respiratory distress syndrome and multiple organ failure (8).

In other situations where complement is profusely activated, such as in systemic lupus erythematosus (SLE), complement fragments deposit on the surface of RBCs, which limits their deformability while promoting nitric oxide (NO) production (9). It has been reported that RBC from patients suffering from hemorrhagic shock displays reduced deformability, shows altered morphological appearance, and contributes to the reduction of blood flow to distant organs (10, 11).

Based on the evolving studies, we hypothesized that complement activation affects RBC function in trauma patients. In this communication, we demonstrate that complement becomes activated in trauma patients and split products are deposited on the surface of RBC limiting their ability to pass through capillary-size microchannels and increasing their capacity to produce NO. Our findings suggest that the inhibition of complement can serve as an adjunct treatment in trauma patients.

MATERIALS AND METHODS

Participants

This prospective experimental study was approved by the Institutional Review Board of Beth Israel Deaconess Medical Center and was carried out as anonymized wasted samples under a waiver of consent. A total of 42 patients who visited the emergency department at Beth Israel Deaconess Medical Center for trauma from July 2012 to March 2013 were enrolled nonconsecutively. Trauma patients were 18 years and older and had been admitted for bone fracture, organ injury, or head trauma. They received a blood test as required for their care. Patients carrying the diagnosis of cancer, infection, or autoimmune diseases were excluded. None of the patients had received a transfusion prior to sampling. Healthy donors ($n = 21$) were recruited from the Division of Rheumatology at Beth Israel Deaconess Medical Center after written informed consent. They did not have an apparent illness. RBCs were collected from 40 of 42 trauma patients (two samples were destroyed during the sample collection process) and 17 healthy donors, and sera were collected from 27 patients and 13 healthy donors. Patients' clinical data were recorded, including age, sex, ethnicity, mechanism of injury, and existing injury. Injury Severity Score (ISS) was also calculated.

Antibodies and Reagents

Primary antibodies were anti-C4d monoclonal antibody (A213; Quidel, San Diego, CA) and IgG1k-isotype control (401402; BioLegend, San Diego, CA). Secondary antibody was Alexa Fluor 488-conjugated rabbit anti-mouse IgG (A11059; Life Technologies, Grand Island, NY). Other reagents were as follows: Hanks' balanced salt solution with Ca^{2+} and Mg^{2+}

(HBSS⁺⁺) (14025; Life Technologies), IgG-free bovine serum albumin (BSA) (A9085; Sigma-Aldrich, St Louis, MO), 4-amino-5-methylamino-2',7'-difluorofluorescein diacetate (D23844; Life Technologies), Flio-4 AM (F14217; Life Technologies), and eosin-5-maleimide (E118; Life Technologies).

Measurement of Serum iC3b Levels

The concentration of iC3b present in serum was measured using the MicroVue iC3b Enzyme Immunoassay kit (A006; Quidel).

RBC Preparation and Incubation With Sera

Heparinized blood or serum of trauma patients was obtained from the emergency department after clinical examination, and control blood or serum was obtained from healthy adult volunteers. The sample collection was performed in accordance with guidelines of the Institutional Review Board of Beth Israel Deaconess Medical Center after informed consent in accordance with the Declaration of Helsinki. Blood was collected in heparin-lithium tubes for RBC and serum in serum-separator tubes. The serum was stored at -80°C . Heparinized blood was washed in HBSS⁺⁺ before use.

For serum studies, on the day of the experiment, the serum samples were thawed from various trauma patients and healthy donors and were used for different measurements under the same conditions. Blood from healthy universal donors (type O, Rh negative) was obtained in heparin-lithium tube and washed with HBSS⁺⁺. One microliter of RBC pellet was resuspended in 80 μL of HBSS⁺⁺, incubated with 20 μL of serum from trauma patients or normal donors for 15 minutes at 37°C , and then washed in HBSS⁺⁺ with 0.5% IgG-free BSA (0.5% BSA/HBSS⁺⁺).

Flow Cytometry

RBCs were incubated for 20 minutes with a primary antibody in 0.5% BSA/HBSS⁺⁺ at room temperature and washed and incubated for another 20 minutes with a secondary antibody directed against the primary antibody at a dilution recommended by the manufacturer. RBCs were then washed and analyzed by FAC-Scan (BD Biosciences, San Jose, CA). At least 10,000 events in each sample were acquired and recorded and analyzed by FlowJo (version 7.6.5 software; Tree Star, Ashland, OR).

Eosin-5-Maleimide Staining

RBCs from healthy universal donors were incubated in the presence of Ca^{2+} and Mg^{2+} with 20% trauma or control sera for 5 minutes at 37°C and then incubated for 15 minutes with eosin-5-maleimide at a final concentration of 0.1 mg/mL. RBCs then were washed three times with HBSS⁺⁺ and 0.5% BSA/HBSS⁺⁺ and analyzed by flow cytometry.

Analysis of Intracellular Calcium Influx

RBCs from healthy universal donors were preloaded with Fluo-4 acetoxymethyl ester (AM) for 15 minutes at 37°C , then washed to remove any dye that is nonspecifically bound with RBC surface, and resuspended in HBSS⁺⁺. Loaded RBCs were incubated at room temperature for an additional 10 minutes to allow complete de-esterification of intracellular AM esters.

Fluorescence levels of RBC were acquired by FACScan for establishing the baseline of intracellular Ca^{2+} concentration. After 60 seconds, 100 μ L of control or trauma serum was quickly added to the RBC (20% of final concentration), and the change of the fluorescence intensity that indicates intracellular Ca^{2+} concentration was recorded for an additional 3 minutes.

Measurement of NO Production

RBCs from healthy universal donors preloaded with DAF-FM diacetate in BSA-free HBSS⁺⁺ for 30 minutes at 37°C were washed to remove excess and uncleaved intracellular DAF-FM diacetate probe. Cells were then resuspended in HBSS⁺⁺ with 20% trauma or control serum for 15 minutes at 37°C. RBCs were washed and the fluorescence intensity associated with intracellular NO production was recorded by FACScan using the fluorescence-1 channel.

Measurement of RBC Deformability Using Two-Dimensional Filters

To measure the ability of RBC to undergo capillary-like deformations, we used a two-dimensional microchannel array as previously described (12) that uses the hydrostatic pressure to drive the RBC through the microchannel array. For each experiment, the device was calibrated using as reference the zero pressure difference corresponded to the absence of movement of RBC within the device. RBCs (8 μ L, hematocrit 20%) were loaded into the inlet reservoir and driven into the capillary-like area by lowering the waste reservoir tubing. Once RBC entered the capillary-like area, the waste reservoir tubing was raised at a height that allowed RBC to pass through the 25- μ m length of the capillary in about 3 seconds. The cells were recorded using a 40 \times 0.75 Ph2 Plan Fluorite objective on a TE300 Nikon-inverted microscope (Nikon, Melville, NY), using a Retiga Exi (QImaging, Surrey, Canada) charge couple device camera controlled with iVision 4.7 software (BioVison, Exton, PA) at a rate of 10 frames/s. Movies representing 600 frames were then analyzed off-line frame by frame, and the results were expressed as the number of seconds from entry to egress (9). RBCs that displayed unusual shapes (e.g., echinocytes and acanthocytes) were excluded from the measurement.

Statistical Analysis

SPSS version 17.0 for Windows (SPSS, Chicago, IL) was used for all statistical analysis. Mann-Whitney *U* test was used for between group analysis and values were showed as medians (interquartile range) unless otherwise stated. Correlation between the ISS and the amount of C4d deposition on trauma RBC was assessed by Spearman correlation coefficient. Paired *t* test analysis was performed wherever necessary. A *p* value of less than 0.05 was considered statistically significant.

RESULTS

Patient Characteristics

A total of 42 patients were included in this study (Table 1). The patient group consisted of 33 men and 9 women, with mean age of 47 years. Most of the patients presented with blunt injury resulting from motor vehicle crash (*n* = 19) or fall (*n* = 16). The median value of ISS was 17 (9–25).

TABLE 1. Demographic and Clinical Characteristics of Research Subjects

Demographic Data	<i>n</i> = 42
Age, mean (\pm sd), yr	46.5 (\pm 19.6)
Sex (male/female)	33/9
Ethnicity, <i>n</i> (%)	
White	37 (88)
Black	2 (5)
Hispanic	1 (2)
Other	2 (5)
Mechanism of trauma, <i>n</i> (%)	
Motor vehicle crash	19 (45)
Fall	16 (38)
Pedestrian struck	1 (2)
Other	6 (14)
Organ injury, <i>n</i> (%)	
Bone fracture	33 (79)
Head	9 (21)
Lung	7 (17)
Spleen	5 (12)
Intestine	3 (7)
Liver	0 (0)
Kidney	0 (0)
Other	1 (2)
Type of bone fracture, <i>n</i>	
Femur	11
Tibia or fibula	10
Spine	9
Rib	9
Radius or ulna	6
Humerus	3
Hip	1
Other	21
Operative procedure, <i>n</i>	
Orthopedic surgery	23
Thoracic surgery	4
Injury Severity Score, mean (\pm sd)	19.6 (\pm 12.6)

Deposition of Complement-Activated Product C4d on RBC From Trauma Patients

To determine the extent of complement deposition on the surface of RBC of trauma patients, we used 40 samples from trauma patients and 17 samples from healthy donors. RBC

TABLE 2. Patient Demographics of C4d Deposition on Trauma RBC

Variable	Trauma Patients (n = 40)	Healthy Controls (n = 17)	p
Age, median (interquartile range), yr	47 (31–50)	36 (34–38)	0.32 ^a
Ethnicity, white, n (%)	36 (90)	10 (59)	0.011 ^b
Sex, male, n (%)	31 (78)	10 (59)	0.13 ^b
Mean fluorescence intensity of C4d level, median (interquartile range)	565 (443–886)	386 (309–489)	< 0.01 ^a

^aMann-Whitney U test.

^bFisher exact test.

from trauma patients displayed significant C4d deposition on their membrane as compared with RBC from normal healthy donor (565 [443–886] vs 386 [309–489], $p < 0.01$) (Table 2). A representative tracing is shown in Figure 1A and cumulative data in Figure 1B. We did not find a correlation between the ISS and the amount of C4d deposited on the surface of RBC (data not shown).

It has been reported previously that complement becomes activated in patients who suffer various forms of trauma (4, 13). To confirm that complement activation occurred in our trauma patients, we measured the iC3b levels in the sera

obtained from trauma patients and healthy donors. Indeed, iC3b levels were significantly higher in the sera of trauma patients ($n = 27$) compared with sera from healthy donors ($n = 12$) (10.7 [8.2–14.0] vs 6.6 [5.3–9.4], $p < 0.01$) (Fig. 1C).

C4d Deposition on Healthy RBC Incubated With Sera From Trauma Patients

To evaluate the effect of complement deposition on various aspects of RBC function, we first established a method to show that incubation of normal RBC from universal donors (type O, Rh negative) with sera from trauma patients results




Figure 1. Increased C4d deposition on the surface of membrane of RBCs from trauma. **A**, Deposition of C4d on the surface of RBCs from trauma patients and healthy donors was measured by flow cytometry. Representative of experiments is shown. **B**, Data of C4d deposition on RBCs from all samples were expressed as mean fluorescence intensity, and cumulative data from trauma patients ($n = 40$) or normal healthy donors ($n = 17$) are shown as box plots. Each box shows the 25th and 75th percentiles. Lines outside the boxes show the lowest and the highest values. Lines inside the boxes show the median. **C**, Serum levels of C3b fragments from trauma patients ($n = 27$) and from normal healthy donors ($n = 12$) were assayed as a measure of activation of complement. Horizontal lines indicate the median.

in complement deposition on the surface of RBC. Indeed, as shown in **Figure 2A** (representative experiment) and **Figure 2B** (cumulative data), incubation of normal RBC with sera from trauma patients ($n = 10$) resulted in a significant C4d deposition on the membrane of RBC as compared with sera from healthy controls ($n = 6$) (289 [282–300] vs 267 [261–272], $p < 0.05$). Removal of Ca^{2+} by adding 10 mM EDTA to the serum decreased the deposition of C4d on the surface of RBC (**Fig. 2C**), suggesting that activation of complement through the classical pathway accounts for the deposition of complement on RBC (14).

Decreased RBC Membrane Deformability by Trauma Sera

It has been previously reported that complement deposition on RBC can render their membrane less deformable, limiting the ability of RBC to pass through capillaries (15). We evaluated RBC membrane deformability by using a microfluidic device as described in *Materials and Methods* section (**Fig. 3A**) and measured the time it took RBC to pass through the microchannels. RBCs from a universal donor were treated either with sera from normal individuals ($n = 6$) or trauma patients ($n = 6$). Individual dot blots are presented in **Figure 3B** and cumulative data in **Figure 3C**. Our results show that exposure

of RBC to sera from trauma patients limits their ability to pass through capillary-size microchannels (2.03 [1.96–2.25] vs 1.85 [1.81–1.90], $p < 0.05$).

Phosphorylation Status of Band 3

RBC membrane deformability is related to the phosphorylation status of membrane proteins, such as β -spectrin or band 3 (16, 17). To determine whether band 3 becomes phosphorylated in RBC exposed to trauma sera, we treated RBC exposed to control ($n = 8$) or trauma sera ($n = 10$) with eosin-5-maleimide, a reagent which binds to lysine 430 on the extracellular loop of band 3 and is known to indicate tyrosine phosphorylation levels (17). Binding of eosin-5-maleimide to RBC has been used previously to estimate the phosphorylation levels of band 3 (9, 17), and here, we similarly used eosin-5-maleimide to estimate the tyrosine phosphorylation levels of band 3. As shown in **Figure 4**, band 3 of RBC was phosphorylated after incubation with trauma sera compared with incubation with normal sera (89.3 [88.3–92.3] vs 84.8 [83.0–85.5], $p < 0.05$).

Promotion of Ca^{2+} Influx

Tyrosine phosphorylation level of band 3 is known to correlate with intracellular Ca^{2+} concentration (18, 19). We asked






Figure 2. Increased C4d deposition on healthy RBC membranes incubated with trauma sera. **A**, Deposition of C4d on the surface of healthy RBCs (type O, Rh negative) incubated with sera from trauma patients or healthy donors was measured by flow cytometry. Representative data are shown. **B**, Data of C4d deposition on RBCs from all samples were expressed as mean fluorescence intensity, and cumulative data from trauma patients ($n = 10$) or healthy donors ($n = 6$) are shown as box plots. Each box shows the 25th and 75th percentiles. Lines outside the boxes show the lowest and the highest values. Lines inside the boxes show the median. **C**, Deposition of C4d on the surface of healthy RBCs (type O, Rh negative) incubated with sera from trauma patients with or without 10mM EDTA. Representative data are shown.






Figure 3. Decreased RBC membrane deformability after incubation with trauma sera. **A**, A composite series of images illustrating the passage of a single RBC through a capillary-size microchannel of the two-dimensional filter device. **B**, Healthy donor's (type O, Rh negative) RBCs deformability were measured by using the two-dimensional filter device after incubation with sera from trauma patients or healthy donors. Decreased deformability of each RBC was associated with increased transit time (i.e., the time it took the cell to pass through the capillary microchannels of two-dimensional filter). Data are shown as dot plots of six experiments with each circle showing the passage time for one RBC. Horizontal lines show the mean. Representative data are shown. **C**, Cumulative data from trauma patients ($n = 6$) or healthy donors ($n = 6$) are shown. Data are expressed as box plots. Each box shows the 25th and 75th percentiles. Lines outside the boxes show the lowest and the highest values. Lines inside the boxes show the median. D = healthy donor, Tr = trauma patient.

whether exposure of RBC to trauma sera results in increased Ca^{2+} entry. As shown in **Figure 5A** (representative tracings) and **Figure 5B** (cumulative data), the addition of trauma sera ($n = 9$) to Fluo-4 AM-loaded RBCs resulted in increased Ca^{2+} concentrations into the RBC compared with RBC to which normal sera ($n = 4$) were added (1.08 [1.05–1.12] vs 1.03 [1.02–1.04], $p < 0.05$).

Induction of the NO Production

Recently, it was reported that human RBCs express an active and functional endothelial-type NO synthase (eNOS), which is present in the membrane and in the cytoplasm (20). In RBC, mechanical stress (shear stress) stimulates NO-generating mechanisms and Ca^{2+} is important in this process (21). Having observed that exposure of RBC to trauma serum results in increased intracellular Ca^{2+} concentrations, we asked whether it also results in an increased NO production. Our results showed that exposure of RBC to trauma sera ($n = 12$) promoted NO production compared with exposure to normal sera ($n = 8$) (543 [527–576] vs 453 [425–501], $p < 0.05$) (**Fig. 6A**, representative tracing, and **Fig. 6B**, cumulative data). Increased NO production results from the deposition of complement on the surface of RBC because heat inactivation of sera (55°C for 30 min) eliminated the ability of trauma sera to increase NO production by RBC (**Fig. 6C**) (597 ± 59 vs 521 ± 21, mean ± SD, paired t test, $p < 0.05$).

DISCUSSION

Complement becomes activated in patients suffering various types of trauma, and this was confirmed in our patients by demonstrating increased levels of iC3b in their sera.




Figure 4. Phosphorylation status of band 3 in RBCs incubated with trauma sera. Phosphorylation of band 3 in healthy RBCs (type O, Rh negative) measured by flow cytometry after incubation with sera from trauma patients or healthy donors, using eosin-5-maleimide staining. Band 3 phosphorylation was expressed as mean fluorescence intensity of eosin-5-maleimide fluorescence, and cumulative data from trauma patients ($n = 10$) or healthy donors ($n = 8$) are shown as box plots. Each box shows the 25th and 75th percentiles. Lines outside the boxes show the lowest and the highest values. Lines inside the boxes show the median.




Figure 5. Ca^{2+} influx into RBC after addition of trauma sera. **A**, Time course of serum-induced changes in intracellular Ca^{2+} of trauma patients (Tr4 and Tr5) or a healthy control (N3) are shown. Arrow indicates the time when trauma or control sera were added to RBC preloaded with Fluo-4 AM, 1 min after start of measurement by flow cytometry. Vertical axis indicates relative intracellular Ca^{2+} concentration, which is estimated from mean fluorescence intensity of Fluo-4 AM. Representative data are shown. **B**, Cumulative results of Ca^{2+} influx in RBCs to which trauma sera ($n=9$) or normal sera ($n=4$) were added. Vertical axis shows the ratio of mean fluorescence intensity of Fluo-4 AM before and after adding the serum. Horizontal lines indicate the median.

Uncontrolled complement activation, common in a wide variety of conditions both chronic and acute, has multiple deleterious effects on various circulating blood cells and on organ function primarily through the action of inflammatory split products (anaphylatoxins) such as C3a and C5a (22, 23). To date, there are little data regarding the effects of the noninflammatory complement by-products, mainly C3b and C4b generated during complement activation on circulating cell function, specifically RBC. In this study, we show that C4d, a split product of C4, deposits on the surface of RBC, thus limiting their ability to pass through narrow microchannels and enhancing NO production. These functional effects are likely due to the fact that C4d deposition on the surface of RBC promotes a significant RBC Ca^{2+} influx that mediates phosphorylation of band 3.

C4d is the stable remnant of C4, and it has been shown to remain on the surface of RBC in patients with SLE for a prolonged period of time (24, 25). In our study, we demonstrated increased deposition of C4d on the surface of RBC from patients with various degrees and types of trauma; yet, we failed to demonstrate a direct correlation between ISS determined clinically and levels of C4d deposition on the surface of RBC. One possible explanation we were not able to establish the correlation is the logistics involved in handling trauma patients and the collection of blood or sera samples. The time that lapsed between trauma and the collection of samples could not be controlled, which could have affected the overall amounts of C4d that had accumulated on the surface of RBC.

Following the two key observations—the increased complement activation in sera and increased deposition of complement fragment C4d on RBC in trauma patients—we chose to

test whether the increase in complement activation and deposition leads to RBC dysfunction. Should binding of complement on the surface of RBC have a functional effect, it should be occurring at early time points. To explore this, RBCs from universal donors were incubated with sera from trauma patients who showed enhanced deposition of C4d on RBC. As noted, exposure of RBC to trauma serum resulted in limitation of their ability to pass through capillaries and increased phosphorylation of band 3, Ca^{2+} entry, and NO production. The alterations in RBC function in trauma patients could be due to deposition of complement fragments on the RBC membranes as others and we have shown both in vitro and ex vivo using RBC from normal individuals (15) and patients with SLE (9).

The functional consequences of the deposition of complement on the surface of RBC cannot be overstated. In order for RBCs to perform their main function, that is to deliver oxygen to tissues, they must pass through narrower ($< 8 \mu\text{m}$ in diameter) capillaries and to do that they have to be able to deform (26, 27). If deposition of C4d renders them difficult to deform and reach tissues, then hypoxia will ensue. During normal conditions, RBC-generated NO is a key factor in the local regulation of vasomotor tone and microvascular flow resistance, by interacting directly with endothelial cells and indirectly with vascular smooth muscles (28). In addition, NO derived from eNOS present in RBC also directly regulates and maintains RBC deformability (20, 29). In our study, we demonstrated that under trauma conditions, increased complement deposition decreases RBC deformability and increases NO production. According to the existing literature, under normal conditions, increase in NO production improves microvascular blood flow. However, in our studies, although the NO production is high in trauma patients,




Figure 6. Nitric oxide (NO) production from RBCs induced by trauma serum. **A**, NO production from healthy RBCs (type O, Rh negative) measured by flow cytometry after incubation with sera from trauma patients or healthy donors by using DAF-FM diacetate. Representative data are shown. **B**, NO production from RBCs was expressed as mean fluorescence intensity of DAF-FM diacetate fluorescence, and cumulative data from trauma patients ($n = 12$) or healthy donors ($n = 8$) are shown as box plots. Each box shows the 25th and 75th percentiles. Lines outside the boxes show the lowest and the highest values. Lines inside the boxes show the median. **C**, Inactivation of complement in sera from trauma patients eliminates the ability to trigger NO production. To inactivate complement, sera were incubated at 55°C for 30 min. RBCs were incubated with sera from trauma patients before or after inactivation of complement. Individual sera with/without inactivation of complement are shown. Straight lines are used to connect individual points to help visualize how different they are in each sample ($n = 4$).

we observe a decrease in RBC deformability. We speculate that this may be due to C4d deposition on RBC in trauma patients that may have a prominent effect on RBC deformability compared with NO effect on RBC deformability. The increased production of NO by C4d-decorated RBC in the microcirculation may further alter epithelial and smooth muscle cell function and influence the clinical picture in trauma patients.

Our studies have certain limitations including the fact that the studied cohort was quite diverse. Although we limited the effect of important comorbidities by excluding patients with significant comorbidities from the study, the data about plausible confounding variables, such as smoking, substance abuse, and medications, were not collected. In addition, we were unable to recruit exact age-, ethnicity-, and sex-matched healthy individuals, and hence, these variables show differences between trauma and control groups. Further studies need to be done in clinically homogenous trauma patients compared with matched controls. The second limitation is that although we made every effort to minimize the time between the collection of the blood sample and the performance of the experiments, we could not control the time between the occurrence of traumatic injury and the time the patient was admitted to the emergency department. As indicated above, during that time, variable amounts of C4d could

have been deposited on the surface of RBC. Yet, the studies in which RBC from universal donors were exposed to trauma serum collected and stored properly lend validity to our conclusions. A proper prospective study (e.g., collection of serial blood samples from trauma patients during the first hours or days after admission) should be more informative. Third, for the serum-based experiments, instead of whole study sample, we used sera from trauma patients who showed increased C4d deposition on RBC. In this study, our goal was to narrow down our experimental design from an epidemiologic standpoint to a proof-of-concept study to further understand the molecular mechanisms.

CONCLUSIONS

We conclude that C4d decorates the surface of RBC and possibly limits their ability to deform and pass through capillary-size microchannels and increases the production of NO. Thus, it may contribute to trauma-associated morbidity and mortality. As a next step, we plan to conduct a prospective study with larger population, which includes clinically homogenous trauma patients compared with matched controls. We believe that our present and future findings might suggest modalities that limit complement activation in trauma patients to be considered for clinical trials.

ACKNOWLEDGMENT

We thank Clinical Research Assistants of the Beth Israel Deaconess Medical Center Emergency Department for collecting blood samples.

REFERENCES

1. U.S. Census Bureau: *Statistical Abstract of the United States: 2012 (131th edition)*. Washington, DC, U.S. Department of Commerce, 2011
2. Hoyert DL, Xu J: Deaths: Preliminary data for 2011. *Natl Vital Stat Rep* 2012; 61:1–51
3. Neher MD, Weckbach S, Flierl MA, et al: Molecular mechanisms of inflammation and tissue injury after major trauma—Is complement the "bad guy"? *J Biomed Sci* 2011; 18:90
4. Fosse E, Pillgram-Larsen J, Svennevig JL, et al: Complement activation in injured patients occurs immediately and is dependent on the severity of the trauma. *Injury* 1998; 29:509–514
5. Szebeni J, Baranyi L, Savay S, et al: Complement activation during hemorrhagic shock and resuscitation in swine. *Shock* 2003; 20:347–355
6. Marik PE, Flemmer M: The immune response to surgery and trauma: Implications for treatment. *J Trauma Acute Care Surg* 2012; 73:801–808
7. Hirsiger S, Simmen HP, Werner CM, et al: Danger signals activating the immune response after trauma. *Mediators Inflamm* 2012; 2012:315941
8. Rittirsch D, Redl H, Huber-Lang M: Role of complement in multiorgan failure. *Clin Dev Immunol* 2012; 2012:962927
9. Ghiran IC, Zeidel ML, Shevkopyas SS, et al: Systemic lupus erythematosus serum deposits C4d on red blood cells, decreases red blood cell membrane deformability, and promotes nitric oxide production. *Arthritis Rheum* 2011; 63:503–512
10. Machiedo GW, Zaets SB, Berezina TL, et al: Trauma-hemorrhagic shock-induced red blood cell damage leads to decreased microcirculatory blood flow. *Crit Care Med* 2009; 37:1000–1010
11. Berezina TL, Zaets SB, Machiedo GW: Alterations of red blood cell shape in patients with severe trauma. *J Trauma* 2004; 57:82–87
12. Glodek AM, Mirchev R, Golan DE, et al: Ligation of complement receptor 1 increases erythrocyte membrane deformability. *Blood* 2010; 116:6063–6071
13. Ganter MT, Brohi K, Cohen MJ, et al: Role of the alternative pathway in the early complement activation following major trauma. *Shock* 2007; 28:29–34
14. Major B, Kardos J, Kékesi KA, et al: Calcium-dependent conformational flexibility of a CUB domain controls activation of the complement serine protease C1r. *J Biol Chem* 2010; 285:11863–11869
15. Karnchanaphanurach P, Mirchev R, Ghiran I, et al: C3b deposition on human erythrocytes induces the formation of a membrane skeleton-linked protein complex. *J Clin Invest* 2009; 119:788–801
16. Manno S, Takakuwa Y, Nagao K, et al: Modulation of erythrocyte membrane mechanical function by beta-spectrin phosphorylation and dephosphorylation. *J Biol Chem* 1995; 270:5659–5665
17. Condon MR, Feketova E, Machiedo GW, et al: Augmented erythrocyte band-3 phosphorylation in septic mice. *Biochim Biophys Acta* 2007; 1772:580–586
18. Minetti G, Piccinini G, Balduini C, et al: Tyrosine phosphorylation of band 3 protein in Ca²⁺/A23187-treated human erythrocytes. *Biochem J* 1996; 320:445–450
19. Zipser Y, Piade A, Barbul A, et al: Ca²⁺ promotes erythrocyte band 3 tyrosine phosphorylation via dissociation of phosphotyrosine phosphatase from band 3. *Biochem J* 2002; 368:137–144
20. Kleinbongard P, Schulz R, Rassaf T, et al: Red blood cells express a functional endothelial nitric oxide synthase. *Blood* 2006; 107:2943–2951
21. Ulker P, Meiselman HJ, Baskurt OK: Nitric oxide generation in red blood cells induced by mechanical stress. *Clin Hemorheol Microcirc* 2010; 45:169–175
22. Baldwin WM 3rd, Kasper EK, Zachary AA, et al: Beyond C4d: Other complement-related diagnostic approaches to antibody-mediated rejection. *Am J Transplant* 2004; 4:311–318
23. Fleming SD, Phillips LM, Lambris JD, et al: Complement component C5a mediates hemorrhage-induced intestinal damage. *J Surg Res* 2008; 150:196–203
24. Manzi S, Navratil JS, Ruffing MJ, et al: Measurement of erythrocyte C4d and complement receptor 1 in systemic lupus erythematosus. *Arthritis Rheum* 2004; 50:3596–3604
25. Kao AH, Navratil JS, Ruffing MJ, et al: Erythrocyte C3d and C4d for monitoring disease activity in systemic lupus erythematosus. *Arthritis Rheum* 2010; 62:837–844
26. Diez-Silva M, Dao M, Han J, et al: Shape and biomechanical characteristics of human red blood cells in health and disease. *MRS Bull* 2010; 35:382–388
27. Mohandas N, Gallagher PG: Red cell membrane: Past, present, and future. *Blood* 2008; 112:3939–3948
28. Baskurt OK, Ulker P, Meiselman HJ: Nitric oxide, erythrocytes and exercise. *Clin Hemorheol Microcirc* 2011; 49:175–181
29. Bor-Kucukatay M, Wenby RB, Meiselman HJ, et al: Effects of nitric oxide on red blood cell deformability. *Am J Physiol Heart Circ Physiol* 2003; 284:H1577–H1584