

Turbulence and Internal Tides on the Continental Slope

Eric Kunze

Applied Physics Laboratory, University of Washington

1013 NE 40th

Seattle, WA 98105-6698

Phone: 206-543-8467 Fax: 206-543-6785 e-mail: kunze@ocean.washington.edu

Award #: N000149410038

<http://www.ocean.washington.edu/people/faculty/kunze/kunze.html>

LONG-TERM GOALS

My interest is in oceanic processes that contribute to stirring and mixing in order to understand their impact on larger scales. This includes phenomena ranging from the meso- (10 km) to the microscale (1 cm), with an emphasis on their interactions, including internal waves, tides, potential-vorticity-carrying finestructure (vortical mode), turbulence and double diffusion.

OBJECTIVES

My recent focus has been on how meso- and finescale flow fields interact with complex topography such as seamounts, canyons, ridges and the continental slope. Mixing in the stratified ocean interior is too weak to close the meridional thermohaline circulation (Ledwell *et al.* 1998). I am exploring whether topographically-enhanced turbulent mixing might be sufficient to do so, and determining what mechanisms are responsible for its generation.

APPROACH

During May 1998, I collected data off the Virginia coast in collaboration with Drs. John Toole and Ray Schmitt and Kurt Polzin (WHOI). This observational program (TWIST – Turbulence and Waves over Irregular Sloping Topography) was designed to characterize the internal wave and turbulence climates above a continental slope with 2-3 km wavelength corrugations running down the slope. Interaction of the corrugations with subinertial alongslope flows associated with topographic Rossby waves was thought to be a likely mechanism for internal lee wave generation. I conducted surveys with expendable current profilers (XCPs) and expendable CTDs (XCTDs) to obtain 3-D snapshots of velocity (u , v), temperature T , salinity S and vertical displacement ξ along and across the corrugations. These measurements complement moored array profile time-series (Toole) and fine- and microstructure profiling with the High-Resolution Profiler (Polzin). Under my supervision, postdoctoral researcher Dr. Jonathan Nash (now an assistant professor at Oregon State University) analysed the data and interpreted it in terms of simplified internal wave models.

Microstructure profiling revealed eddy diffusivities in the stratified bottom boundary layer more than one hundred times open-ocean levels between the 1000- and 1200-m isobaths; *identical* values are found using the Gregg-Henyey finescale parameterization for internal-wave-generated turbulence (Gregg 1989). Our goal was to determine the mechanisms responsible for this enhanced mixing.

Report Documentation Page

Form Approved
OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE 30 SEP 2003		2. REPORT TYPE		3. DATES COVERED 00-00-2003 to 00-00-2003	
4. TITLE AND SUBTITLE Turbulence and Internal Tides on the Continental Slope				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Applied Physics Laboratory, University of Washington,,1013 NE 40th,,Seattle,,WA, 98105				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT My interest is in oceanic processes that contribute to stirring and mixing in order to understand their impact on larger scales. This includes phenomena ranging from the meso- (10 km) to the microscale (1 cm), with an emphasis on their interactions, including internal waves, tides, potential-vorticity-carrying finestructure (vortical mode), turbulence and double diffusion.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

WORK COMPLETED

Analysis of the data is complete. A manuscript has been submitted to *J. Phys. Oceanogr.* (J.D. Nash, E. Kunze, J.M. Toole and R.W. Schmitt “Internal Tide Reflection and Turbulent Mixing on the Continental Slope”) and is presently under revision. We examined the finescale data in the expendable surveys and the three moored profile time-series, focussing on the latter half of the record when an along-isobath semidiurnal energy-flux was observed and the expendable surveys provided spatial coverage. A decomposition by vertical scale of the semidiurnal signal reveals the dominant physics. A 2-D ray-tracing model following Eriksen (1982) reproduces many features of the observations.

RESULTS

In contrast to expectations, we found no evidence for internal lee waves based on the absence of corrugation-scale finestructure in the water column. While the flow in most of the water column was in excess of 10 cm s^{-1} , the near-bottom flow was too weak to generate waves

$[U < N\alpha / (k_x \sqrt{1 + f^2 / N^2 \alpha^2}) \sim 15 \text{ cm s}^{-1}$, where α is the slope and k_x the wavelength of the corrugations]. We also found no evidence for internal tide generation at the shelf break, consistent with numerical simulations of the region (Legg 2003). Fluctuations during the last half of the sampling period are dominated by semidiurnal oscillations. Analysis indicates:

- The near-bottom mean flow is *around* not *over* the topographic corrugations consistent with it being too weak to generate lee waves;
- The vertically-integrated internal-wave energy-flux is northward parallel to the continental slope at $O(1 \text{ kW m}^{-1})$;
- Cross- exceeds along-isobath velocity variance;
- Shear is intensified in the bottom 300 m and is predominantly semidiurnal in the latter half of the sampling period when the expendable surveys took place;
- Near-bottom dissipation and shear is enhanced between the 1000- and 1200-sm (WKB stretched meters) isobaths offshore of supercritical (M_2) across-slope bathymetry (right panel of Fig. 1).




Figure 1: Across-slope sections along a corrugation ridge. Data are in a WKB-normalized format. Upper panels show vertically-integrated across-isobath energy-fluxes. There is convergence of low-mode onshore energy-flux (left) and divergence of high-wavenumber offshore flux which takes the form of an intense near-bottom beam emanating between the 1000- and 1200-sm (WKB stretched meters) isobaths (middle). Turbulent eddy diffusivities inferred from the Gregg-Henyey scaling (right) are enhanced only offshore of the near-critical, 1000-sm isobath and coincide with the high-wavenumber beam.

We hypothesize that enhanced semidiurnal shear and mixing is associated with reflection of a low-mode semidiurnal internal wave from near- and supercritical bathymetry. In support of this:

- While the net energy-flux is northward along isobaths, cross-isobath velocities exceed along-isobath velocities, and there is an abrupt cross-slope phase shift near the 1000-m isobath;
- Local generation of internal tides directly from barotropic semidiurnal motions is too weak to account for the observed shear or mixing (Legg 2003);
- A breakdown by vertical scale reveals that there is a cross-slope *convergence* of mode-one energy-flux approximately balanced by cross-slope *divergence* of high-wavenumber energy-flux plus turbulent dissipation (Fig. 1). Theory indicates that an *along-isobath* energy-flux is to be expected within one-quarter wavelength of the slope for the superposition of an incident incoming wave and reflected outgoing *across-isobath* wave;
- Beams of high-wavenumber flux radiate offshore from near the 1000-sm isobath. They coincide with high shear and turbulence signals (central panel of Fig. 1). Well-mixed bottom boundary layers more than 10-m thick are only observed offshore of supercritical topography.

To understand whether the observed semidiurnal shear could result from reflection of a low-mode internal wave off the continental slope, a simple ray-tracing simulation was performed. Bathymetry is assumed two-dimensional, ignoring alongslope corrugations. Boundary conditions consist of an onshore-propagating mode-one internal tide of 2 cm s^{-1} amplitude prescribed 150 km offshore.

Information travels along rays, obeying Eriksen (1982) reflection laws at the surface and bottom, until leaving the domain. The simulation reproduces most of the features of the observations (Fig. 2).




Figure 2: Energy-flux sections from a WKB ray-tracing model following Eriksen (1982). The left panel shows the shoreward flux, the central panel the offshore flux and the right panel the net flux. The box in the right panel denotes the area shown in Fig. 1. Note the near-bottom high-wavenumber variance between the 1000- and 1200-sm isobaths.

The ray-tracing simulations reproduce a 200-m thick layer of high-wavenumber variance on the slope between the 1000- and 1200-sm isobaths (Fig. 2), consistent with the observations (Fig. 1). The model predicts shears exceeding the stratification ($Ri > 1$). Parker MacCready (UW) is running numerical simulations with the Hallberg Isopycnal Model for comparison with the ray-tracing results.

IMPACT/APPLICATION

Our observations and model results suggest that the shear and mixing is associated with reflection of the low-mode internal tide with little dependence on the alongslope corrugations that originally drew our attention to the TWIST site. Elevated turbulence is also reported on the Oregon continental slope (Moum *et al.* 2002). Analysis of historical mooring data suggests that a similar critical reflection of a deep-ocean low-mode internal tide may be responsible. This suggests that continental slopes may be sinks for internal tides generated in the deep ocean. This contrasts with the earlier paradigm of internal tide generation at the shelf break (Prinsenber *et al.* 1974; Baines 1982) radiating semidiurnal internal waves into the deep ocean and onto the shelf. In laboratory experiments of internal waves reflecting off sloping topography, McPhee-Shaw and Kunze (2002) found that the exchange of fluid between the turbulent boundary layer and the quiescent interior was controlled by the turbulent mixing at the boundary which could in turn be diagnosed by the convergence of across-slope internal wave energy-flux. This bears on both boundary-interior fluid exchange and intermediate nepheloid production.

TRANSITIONS

The energy-flux estimation technique is being used on other projects by the PI and others to examine internal tide energy budgets in Monterey Submarine Canyon (Kunze *et al.* 2002), across Mendocino Escarpment (Althaus *et al.* 2003), along the Hawaiian Ridge (HOME; Rudnick *et al.* 2003) and in the world ocean using historical current-meter records (Alford 2003).

RELATED PROJECTS

Larger fluxes are found radiating away from Mendocino Escarpment and the Hawaiian Ridge, as well as into the mouth of Monterey Submarine Canyon. Turbulence levels above the Virginia continental slope are comparable to those observed over seamounts, escarpments and ridges. Motivated by the Oregon slope microstructure measurements (Moum *et al.* 2002), Drs. E. Kunze, J. Nash and M. Alford have submitted a proposal to NSF to make finestructure measurements on the Oregon continental slope to determine if the same mechanism may be responsible. Existing evidence suggests that, except in abyssal waters below 4000-m depth, topographically-enhanced mixing, although 100-1000 times typical interior values (e.g., Kunze and Toole 1997), is not large enough to close the meridional thermohaline overturning cell as envisioned by Munk and Wunsch (1998). This would leave outcrop mixing as the only viable candidate for waters of 1-3 km depth in the temperate and tropical oceans (Sloyan and Rintoul 2001; Toggweiler and Samuels 1998).

REFERENCES

- M.H. Alford, 2003: Energy available for ocean mixing redistributed by long-range propagation of internal waves. *Nature*, **423**, 159-162.
- A.M. Althaus, E. Kunze and T.B. Sanford, 2003: Internal tide radiation from Mendocino Escarpment. *J. Phys. Oceanogr.*, **33**, 1510-1527.
- P.G. Baines, 1982: On internal tide generation models. *Deep-Sea Res.*, **29**, 307-338.
- C.C. Eriksen, 1982: Observations of internal wave reflection off sloping bottoms. *J. Geophys. Res.*, **87**, 525-538.
- M.C. Gregg, 1989: Scaling turbulent dissipation in the thermocline. *J. Geophys. Res.*, **94**, 9686-9698.
- E. Kunze and J. M. Toole, 1997: Tidally-driven vorticity, diurnal shear and turbulence atop Fieberling Seamount. *J. Phys. Oceanogr.*, **27**, 2663-2693.
- E. Kunze, L.K. Rosenfeld, G.S. Carter and M.C. Gregg, 2002: Internal waves in Monterey Submarine Canyon. *J. Phys. Oceanogr.*, **32**, 1890-1913
- J. R. Ledwell, A. J. Watson and C. S. Law, 1998: Mixing of a tracer in the pycnocline. *J. Geophys. Res.*, **103**, 21,499-21,529.
- S. Legg, 2003: Internal tide generated on a corrugated continental slope. *J. Phys. Oceanogr.*, submitted.
- E.E. McPhee-Shaw and E. Kunze, 2002: Boundary-layer intrusions from a sloping bottom: A mechanism for generating intermediate nepheloid layers. *J. Geophys. Res.*, **107**, DOI: 1029/2001JC000801.
- J.N. Moum, D.R. Caldwell, J.D. Nash and G.D. Gunderson, 2002: Observations of boundary mixing over the continental slope. *J. Phys. Oceanogr.*, **32**, 2113-2130.

W. Munk and C. Wunsch, 1998: Abyssal recipes II: Energetics of tidal and wind mixing. *Deep-Sea Res.*, **45**, 1977-2010.

S.J. Prinsenber, W.L. Wilmot and M. Rattray Jr., 1974: Generation and dissipation of coastal internal tides. *Deep-Sea Res.*, **21**, 263-281.

D.L. Rudnick, T. Boyd, R. Brainard, G.S. Carter, G.D. Egbert, M.C. Gregg, P.E. Holloway, J. Klymak, E. Kunze, C.M. Lee, M.D. Levine, D.S. Luther, J. Martin, M.A. Merrifield, J.N. Moum, J.D. Nash, R. Pinkel, L. Rainville, T.B. Sanford and J. Sherman, 2002: From tides to mixing along the Hawaiian Ridge. *Science*, **301**, 355-357.

B. M. Sloyan and S. R. Rintoul, 2001: The Southern Ocean limb of the global deep overturning circulation. *J. Phys. Oceanogr.*, **31**, 143-173.

J. R. Toggweiler and B. Samuels, 1998: On the ocean's large-scale circulation near the limit of no vertical mixing. *J. Phys. Oceanogr.*, **28**, 1832-1852.

PUBLICATIONS

E. Kunze, 2003: A review of salt-fingering theory. *Prog. Oceanogr.*, **56**, 399-417.

K. L. Polzin, E. Kunze, J. M. Toole and R. W. Schmitt, 2003: The partition of finescale energy into internal waves and geostrophic motions. *J. Phys. Oceanogr.*, **33**, 234-248.

E. E. McPhee-Shaw and E. Kunze, 2002: Boundary-layer intrusions from a sloping bottom: A mechanism for generating intermediate nepheloid layers. *J. Geophys. Res.*, **107**, DOI: 1029/2001JC000801.

J.D. Nash, E. Kunze, J.M. Toole and R.W. Schmitt, 2003: Internal Tide Reflection and Turbulent Mixing on the Continental Slope. *J. Phys. Oceanogr.*, submitted.