

RESEARCH HIGHLIGHT

Asynchronous inflammation and myogenic cell migration limit muscle tissue regeneration mediated by a cellular scaffolds

Koyal Garg^{1,2}, Catherine L. Ward¹, Benjamin T. Corona¹

¹United States Army Institute of Surgical Research, Extremity Trauma and Regenerative Medicine

²University of Illinois, Urbana Champaign, Beckman Institute of Advanced Science and Technology

Correspondence: Benjamin T. Corona

E-mail: benjamin.t.corona.vol@mail.mil or corona.benjamin.t@gmail.com

Received: January 14, 2015

Published online: February 11, 2015

Volumetric muscle loss (VML) following orthopaedic trauma results in chronic loss of strength and can contribute to disability. Tissue engineering and regenerative medicine approaches to regenerate the lost skeletal muscle and improve functional outcomes are currently under development. At the forefront of these efforts, decellularized extracellular matrices (ECMs) have reached clinical testing and provide the foundation for other approaches that include stem/progenitor cell delivery. ECMs have been demonstrated to possess many qualities to initiate regeneration, to include stem cell chemotaxis and pro-regenerative macrophage polarization. However, the majority of observations indicate that ECM-repair of VML does not promote appreciable muscle fiber regeneration. In a recent study, ECM-repair of VML was compared to classical muscle fiber regeneration (Garg et al., 2014, Cell & Tissue Research) mediated by autologous minced grafts. The most salient findings of this study were: 1) Satellite cells did not migrate into the scaffold beyond ~0.5 mm from the remaining host tissue, although other migratory stem cells (Sca-1⁺) were observed throughout the scaffold; 2) Macrophage migration to the scaffold was over two-times that observed with muscle grafts, but they appeared to be less active, as gene expression of pro- and anti-inflammatory cytokines (TNF- α , IL-12, IL-4, IL-10, VEGF, and TGF- β 1) was significantly reduced in scaffold-repaired muscles; And, 3) scaffolds did not promote appreciable muscle fiber regeneration. Collectively, these data suggest that the events following ECM transplantation in VML are either incongruous or asynchronous with classical muscle fiber regeneration.

To cite this article: Koyal Garg, et al. Asynchronous inflammation and myogenic cell migration limit muscle tissue regeneration mediated by a cellular scaffolds. Inflamm Cell Signal 2015; 2: e530. doi: 10.14800/ics.530.

Orthopaedic trauma in civilian and military populations usually involves severe skeletal muscle injury. Either directly (e.g., blast injury) or indirectly (e.g., soft tissue evacuation secondary to crush), the traumatized musculature often presents volumetric muscle loss (VML) [1]; which mammalian skeletal muscle is not capable of successfully regenerating [2-4]. There is currently no effective therapy used in the clinic that promotes *de novo* muscle tissue regeneration following VML. To this end, persistent strength deficits and limb dysfunction are currently anticipated as part of the sequelae of this injury [5]. Multiple efforts are ongoing to develop tissue engineering and regenerative medicine

therapies for the repair of VML- the intention of the therapies is to restore strength to the injured musculature by regenerating appreciable and functional muscle tissue. At the forefront of these efforts are decellularized extracellular matrix scaffolds (ECMs) (Table 1), which are being investigated in the clinic for VML [6]. While ECMs represent a logical therapy for VML, by-and-large the capacity of ECMs to promote *de novo* skeletal muscle regeneration appears to be limited [3, 6-8]. The purpose of this report is to compare the putative mechanism of ECM-mediated regeneration with that of classical muscle repair and regeneration.

Report Documentation Page

Form Approved
OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE 01 FEB 2015		2. REPORT TYPE N/A		3. DATES COVERED -	
4. TITLE AND SUBTITLE Asynchronous inflammation and myogenic cell migration limit muscle tissue regeneration mediated by acellular scaffolds				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Garg K., Ward C. L., Corona B. T.,				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) United States Army Institute of Surgical Research, JBSA Fort Sam Houston, TX 78234				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT SAR	18. NUMBER OF PAGES 5	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Table 1. Selected studies of VML repair (Listed chronologically)

	VML Model	VML Size	Species	Therapy	Primary Histological Outcome
Brown et al. ^[8]	Abdominal Wall	1cm x 1cm	Rats	Rat Muscle Autograft	Fibrosis and inflammation
				Rat mECM	Fibrosis and small islands of myofibers
				Porcine Muscle Graft	Fibrosis with inflammation
				Porcine UBM	Fibrosis and small islands of myofibers
Turner et al. ^[36]	Gastrocnemius	N/A	Dogs	Porcine SIS	Myofiber regeneration throughout defect area
Merritt et al. ^[13, 49]	Gastrocnemius	~225 mg	Rats	Rat mECM	Limited ingrowth of myofibers
				Rat mECM + MSCs	Improved but limited myofiber regeneration
Mase et al. ^[4]	Quadriceps	N/A	Human	Porcine SIS	N/A : CT showed unknown tissue deposition
Machingal et al. ^[50]	Latissimus dorsi	~ 25 mg	Mice	Porcine BAM + MDCs	Desmin+ cells throughout defect & fibrosis
				Porcine BAM	Fibrosis & diminished myofiber regeneration
Sicari et al. ^[14]	Quadriceps	4 x 4 x 3 mm	Mice	Porcine SIS	Fibrosis & small islands of myofibers
Corona et al. ^[3]	Tibialis anterior	10x7x3 mm	Rats	Rat mECM	Fibrosis with limited muscle regeneration
				Rat mECM + MSCs	Fibrosis with limited muscle regeneration
Corona et al. ^[2]	Tibialis anterior	10x7x3 mm	Rats	Rat Minced Muscle Grafts	Myofiber regeneration throughout defect area
Sicari et al. ^[6]	Quadriceps	~ 75% loss	Mice	Porcine UBM	Fibrosis & limited ingrowth of muscle fibers
	Lower Extremity	N/A	Human	Porcine UBM	Fibrosis & small islands of myofibers
Ma et al. ^[12]	Abdominal Wall	1.2 x 1.2 cm	Rat	Porcine SIS	Fibrosis with no muscle fiber regeneration
Garg et al. ^[16]	Tibialis Anterior	10x7x3 mm	Rat	Rat Minced Muscle Grafts	Myofiber regeneration throughout defect area
				Rat Muscle Scaffold	Fibrosis throughout defect area

mECM (muscle ECM); UBM (urinary bladder matrix ECM); SIS (small intestine submucosa ECM); MSC (mesenchymal stem cell); CT (computed tomography); BAM (bladder acellular matrix ECM); MDCs (muscle-derived cells)

The precise mechanism underlying ECM-mediated regeneration in skeletal muscle has not been completely elucidated. Following transplantation in a fresh VML wound bed, ECMs are infiltrated with mononuclear cells within hours^[9]. Thereafter, ECMs substantially degrade over the ensuing month^[10, 11]. Within this time, ECMs are thought to promote skeletal muscle regeneration *in vivo* by orchestrating the following events through an undefined spatiotemporal pattern (*see review*:^[9]): **1)** promotion of a vascular bed, **2)** chemotaxis for nearby resident and circulating progenitor and stem cells, **3)** provision of matrix bound growth factors and cryptic peptides that direct functions of migrating cells, and **4)** induction of pro-regenerative macrophage polarization *in vitro*. Ultimately, it is thought that the degrading ECM creates a pro-regenerative environment that promotes myogenic differentiation of migrating stem and progenitor cells, which then effectively regenerate functional muscle fibers.

Despite the regenerative attributes of ECMs, many studies present evidence at prolonged time points (months) that ECMs do not promote appreciable muscle fiber regeneration *in vivo*, particularly in regions of the defect further removed from the remaining tissue bed- instead fibrotic tissue is remodeled^[3, 6, 7, 12-15]. These observations have been made among various labs in different species and VML models using a variety of ECMs derived from autologous, syngeneic, or xenogeneic tissues, to include bladder, small intestine, and muscle (**Table 1**). That being said, fibrotic tissue deposition following ECM transplantation may have under-appreciated

therapeutic benefits, to include improving strength via improved force transmission (as opposed to production)^[3] and protecting the remaining muscle mass from chronic overload-induced injury^[2, 3].

Recently we observed restricted and limited muscle fiber regeneration after autologous devitalized muscle scaffold repair of VML in the rat TA muscle two months post-injury^[16]. Given this fairly consistent observation (**Table 1**), we posited that while ECM preparation may explain subtle variations in regenerative outcomes^[17], it is more likely that the underlying mechanism of ECM-remodeling in a VML defect does not adequately mimic that of adult endogenous adult muscle fiber regeneration. To investigate this possibility, we further interrogated at an acute time point (2 weeks post-injury) the inflammatory and myogenic response to the muscle scaffold^[16], which relies solely on host cell migration for regeneration^[18]. Vital minced grafts (1 mm³ pieces of tissue), which also deliver satellite cells among other cellular and trophic factors resident in whole skeletal muscle, were used as a positive control of successful muscle regeneration^[2, 18-20]. In the following sections, the findings of this study are highlighted and discussed in the context of events necessary for classical muscle repair and regeneration.

Satellite cells (Pax7⁺) are a skeletal muscle specific stem cell population that is indispensable for muscle regeneration following injury^[21]. A combination of signals from the host muscle fibers, circulating cells, interstitial cells (such as macrophages) and muscle resident stem cells can influence

the quiescence, activation and proliferation of the satellite cells [22]. It has been shown that, over 100 new myofibers containing thousands of myonuclei can be generated from as little as seven activated satellite cells associated with one transplanted myofiber [23]. The loss of satellite cell pool due to aging or disease (such as duchenne muscular dystrophy) results in impaired regeneration, increased atrophy and fibrosis of skeletal muscle [24-27]. It has also been reported that genetic elimination of satellite cells results in a complete blockade of regenerative myogenesis following cardiotoxin injury [28, 29]. Other host cell populations are unable to compensate for the loss of regenerative potential and muscle regeneration can only be rescued by replenishing the satellite cell pool [29]. In our study, Pax7⁺ cells and small regenerating myofibers (myosin⁺) were observed in the muscle scaffolds in close proximity to the remaining musculature, but were absent in the defect beyond 0.5 mm from the remaining muscle mass. In contrast, minced graft transplantation presented muscle fiber regeneration throughout the defect with associated satellite cell (Pax7⁺) co-localization. Given the importance of satellite cells to muscle regeneration, the restricted satellite cell migration into the muscle scaffold appear to be a significant limitation to ECM-mediated muscle fiber regeneration.

Besides satellite cells, several other cell types such as bone marrow derived progenitors [30], pericytes [31, 32], mesoangioblasts [33], interstitial cells (PW1⁺/Pax7) [34], perivascular stem cells [35], CD133⁺ progenitors [36] have been shown to be myogenic. And, these alternative myogenic stem cells have been identified in remodeling ECMs [6, 36]. Overall, the contribution of these alternative myogenic cells towards regeneration is believed to be lower compared to satellite cells, but important never the less [30, 31]. We speculate that the therapeutic potential of these cells is limited to an even greater extent in a VML injury model, due to the stark absence of key myogenic cues from the lost myofiber remnants and satellite cells. In the highlighted study, Sca-1⁺ cells migrated throughout muscle scaffold, and importantly in regions removed from the remaining muscle mass that were unpopulated by Pax7⁺ cells. The presence of Sca-1⁺ cells indicate that the muscle scaffold was conducive to stem and progenitor cells migration, but signify that their presence is not sufficient for *de novo* muscle fiber regeneration. Further studies are needed to assess if these potentially alternative myogenic cells require interaction with adult myofibers or activated satellite cells [37] to induce myogenesis.

Following injury, the type of the inflammatory response and the significance of transition from pro- to an anti-inflammatory phenotype is widely recognized as a critical component to regeneration. Through the phases of

repair and regeneration, resident and recruited macrophages among other immune cells exhibit complex and hybrid phenotypes that are believed to be successively activated and timely subsided. While, pro-inflammatory activity is important for the migration and activation of myogenic precursor cells [38-40], an anti-inflammatory response stimulates muscle precursor cell differentiation and myotube formation [7, 38, 40, 41]. An extended and unrestricted pro-inflammatory (M1) macrophage response or an early and premature anti-inflammatory (M2) response has been shown to result in unsuccessful regenerative outcomes [19-23]. In our study, we generally characterized the inflammatory response of muscles repaired with muscle scaffolds and vital grafts at two weeks post-injury. The vital grafts supported a mixed up-regulation of pro- and anti-inflammatory markers *in vivo* that corresponded with improved regenerative outcomes. The overall inflammatory response appeared to be functionally attenuated in scaffold-repaired muscles, which corresponded with aberrant regeneration and fibrotic tissue deposition. Therefore, we speculate that both functionally active populations of pro- and anti-inflammatory cells (e.g., M1 and M2 macrophage phenotypes) are important for *de novo* regeneration of VML injured skeletal muscle. In support of this speculation, other studies have also demonstrated that coordinated efforts by both M1 and M2 macrophage phenotypes are necessary for scaffold vascularization [42] and remodeling [43]. A recent study by Novak *et al.*, reported that macrophage populations present after muscle laceration injury, did not conform to the canonical M1/M2 classification. Furthermore, they demonstrated that activated donor M1 macrophages reduced fibrosis and enhanced myofiber regeneration but non-activated macrophages had no effect [44]. Collectively these data suggest that synchronized efforts of functionally active pro- and anti-inflammatory cells (e.g. M1 and M2 macrophage phenotypes) are required for effective regeneration.

Following recoverable injuries, factors released from injured muscle fibers induce an intricate spatiotemporal communication between the innate immune response and myogenic progenitor cell activity (i.e., primary satellite cells) that promotes repair and regeneration of the injured tissue. Pro-inflammatory M1 macrophages produce cytokines such as TNF- α and IL-6 that promote satellite cell activation and proliferation but not differentiation [45]. Anti-inflammatory M2 macrophages promote myoblast differentiation and fusion and myotube maturation by releasing cytokines such as interleukin (IL)-4 and IL-10 [38, 41]. In addition to secreted myogenic factors, human macrophages have also been shown to deliver anti-apoptotic signals to myogenic cells through direct cell-cell contact [46]. Evidence of two-way communication between myogenic cells and macrophages was demonstrated by our *in vitro* study in which secreted

factors from vital grafts were able to induce arginase expression (indicative of an M2-like phenotype) in macrophages. Therefore, disruption of either the inflammatory response or myogenic cell activity can significantly impact the regenerative outcomes. For instance, restricting monocyte/macrophage tissue invasion by disruption of CCL2/CCR2 signaling has been shown to impair regeneration and functional recovery^[47, 48]. Likewise, ablation of satellite cells using an inducible Pax7 knockout model abrogated regeneration after toxin injury^[28, 29]. The findings of our paper indicate that vital minced grafts comprised of myogenic cells in their native extracellular environment allow for interaction with the ensuing inflammatory response and promote effective muscle tissue regeneration. In contrast, transplantation of a devitalized scaffold results in a stark absence of satellite cells in distant regions of the defect and a diminished activity of pro- and anti-inflammatory cells. These observations present a deviation from the normal spatiotemporal events of muscle regeneration that is likely a major contributor to aberrant regenerative outcomes observed with ECMs. As the field continues to develop approaches to regenerate *de novo* muscle tissue after VML, ECMs and other scaffolds will likely be fundamental to these efforts. However, pursuant to improved regenerative outcomes with ECM-repair of VML is the development of strategies to further align the events of ECM remodeling with the intricate spatiotemporal events that underlie adult skeletal muscle regeneration.

Acknowledgments

The opinions or assertions contained herein are the private views of the authors and are not to be construed as official or as reflecting the views of the Department of the Army or the Department of Defense. This work was supported by the Combat Casualty Care Research Program, Medical Research and Materiel Command.

References

- Grogan BF, Hsu JR. Volumetric muscle loss. *J Am Acad Orthop Surg* 2011; 19 Suppl 1:S35-37.
- Corona BT, Garg K, Ward CL, Mcdaniel JS, Walters TJ, Rathbone CR. Autologous minced muscle grafts: a tissue engineering therapy for the volumetric loss of skeletal muscle. *Am J Physiol Cell Physiol* 2013; 305:C761-775.
- Corona BT, Wu X, Ward CL, Mcdaniel JS, Rathbone CR, Walters TJ. The promotion of a functional fibrosis in skeletal muscle with volumetric muscle loss injury following the transplantation of muscle-ECM. *Biomaterials* 2013; 34:3324-3335.
- Mase VJ, Jr., Hsu JR, Wolf SE, Wenke JC, Baer DG, Owens J, et al. Clinical application of an acellular biologic scaffold for surgical repair of a large, traumatic quadriceps femoris muscle defect. *Orthopedics* 2010; 33:511.
- Garg K, Ward CL, Hurtgen BJ, Wilken JM, Stinner DJ, Wenke JC, et al. Volumetric muscle loss: persistent functional deficits beyond frank loss of tissue. *J Orthop Res* 2015; 33:40-46.
- Sicari BM, Rubin JP, Dearth CL, Wolf MT, Ambrosio F, Boninger M, et al. An acellular biologic scaffold promotes skeletal muscle formation in mice and humans with volumetric muscle loss. *Sci Transl Med* 2014; 6:234-258.
- Brown BN, Londono R, Tottey S, Zhang L, Kukla KA, Wolf MT, et al. Macrophage phenotype as a predictor of constructive remodeling following the implantation of biologically derived surgical mesh materials. *Acta Biomater* 2012; 8:978-987.
- Brown BN, Valentin JE, Stewart-Akers AM, McCabe GP, Badylak SF. Macrophage phenotype and remodeling outcomes in response to biologic scaffolds with and without a cellular component. *Biomaterials* 2009; 30:1482-1491.
- Wolf MT, Dearth CL, Sonnenberg SB, Lobo E, Badylak SF. Naturally derived and synthetic scaffolds for skeletal muscle reconstruction. *Adv Drug Deliv Rev* 2014.
- Carey LE, Dearth CL, Johnson SA, Londono R, Medberry CJ, Daly KA, et al. In vivo degradation of 14C-labeled porcine dermis biologic scaffold. *Biomaterials* 2014; 35:8297-8304.
- Gilbert TW, Stewart-Akers AM, Badylak SF. A quantitative method for evaluating the degradation of biologic scaffold materials. *Biomaterials* 2007; 28:147-150.
- Ma J, Sahoo S, Baker AR, Derwin KA. Investigating muscle regeneration with a dermis/small intestinal submucosa scaffold in a rat full-thickness abdominal wall defect model. *J Biomed Mater Res B Appl Biomater* 2014.
- Merritt EK, Hammers DW, Tierney M, Suggs LJ, Walters TJ, Farrar RP. Functional assessment of skeletal muscle regeneration utilizing homologous extracellular matrix as scaffolding. *Tissue Eng Part A* 2010; 16:1395-1405.
- Sicari BM, Agrawal V, Siu BF, Medberry CJ, Dearth CL, Turner NJ, et al. A murine model of volumetric muscle loss and a regenerative medicine approach for tissue replacement. *Tissue Eng Part A* 2012; 18:1941-1948.
- Turner NJ, Badylak JS, Weber DJ, Badylak SF. Biologic scaffold remodeling in a dog model of complex musculoskeletal injury. *J Surg Res* 2012; 176:490-502.
- Garg K, Ward CL, Rathbone CR, Corona BT. Transplantation of devitalized muscle scaffolds is insufficient for appreciable *de novo* muscle fiber regeneration after volumetric muscle loss injury. *Cell & Tissue Research* 2014; In Press.
- Keane TJ, Badylak SF. The host response to allogeneic and xenogeneic biological scaffold materials. *J Tissue Eng Regen Med* 2014.
- Ghins E, Colson-Van Schoor MM, Marechal G. The origin of muscle stem cells in rat triceps surae regenerating after mincing. *J Muscle Res Cell Motil* 1984; 5:711-722.
- Carlson BM, Gutmann E. Development of contractile properties of minced muscle regenerates in the rat. *Exp Neurol* 1972; 36:239-249.
- Snow MH. Metabolic activity during the degenerative and early regenerative stages of minced skeletal muscle. *Anat Rec* 1973; 176:185-203.
- Relaix F, Zammit PS. Satellite cells are essential for skeletal muscle regeneration: the cell on the edge returns centre stage.

- Development 2012; 139:2845-2856.
22. Kuang S, Gillespie MARudnicki MA. Niche regulation of muscle satellite cell self-renewal and differentiation. *Cell Stem Cell* 2008; 2:22-31.
 23. Collins CA, Olsen I, Zammit PS, Heslop L, Petrie A, Partridge TA, *et al.* Stem cell function, self-renewal, and behavioral heterogeneity of cells from the adult muscle satellite cell niche. *Cell* 2005; 122:289-301.
 24. Cohn RDCampbell KP. Molecular basis of muscular dystrophies. *Muscle Nerve* 2000; 23:1456-1471.
 25. Morgan JEZammit PS. Direct effects of the pathogenic mutation on satellite cell function in muscular dystrophy. *Exp Cell Res* 2010; 316:3100-3108.
 26. Conboy IMRando TA. Aging, stem cells and tissue regeneration: lessons from muscle. *Cell Cycle* 2005; 4:407-410.
 27. Chen YW, Nagaraju K, Bakay M, Mcintyre O, Rawat R, Shi R, *et al.* Early onset of inflammation and later involvement of TGFbeta in Duchenne muscular dystrophy. *Neurology* 2005; 65:826-834.
 28. Lepper C, Partridge TAFan CM. An absolute requirement for Pax7-positive satellite cells in acute injury-induced skeletal muscle regeneration. *Development* 2011; 138:3639-3646.
 29. Sambasivan R, Yao R, Kissenpfennig A, Van Wittenberghe L, Paldi A, Gayraud-Morel B, *et al.* Pax7-expressing satellite cells are indispensable for adult skeletal muscle regeneration. *Development* 2011; 138:3647-3656.
 30. Ferrari G, Cusella-De Angelis G, Coletta M, Paolucci E, Stornaiuolo A, Cossu G, *et al.* Muscle regeneration by bone marrow-derived myogenic progenitors. *Science* 1998; 279:1528-1530.
 31. Dellavalle A, Maroli G, Covarello D, Azzoni E, Innocenzi A, Perani L, *et al.* Pericytes resident in postnatal skeletal muscle differentiate into muscle fibres and generate satellite cells. *Nat Commun* 2011; 2:499.
 32. Dellavalle A, Sampaolesi M, Tonlorenzi R, Tagliafico E, Sacchetti B, Perani L, *et al.* Pericytes of human skeletal muscle are myogenic precursors distinct from satellite cells. *Nat Cell Biol* 2007; 9:255-267.
 33. Sampaolesi M, Blot S, D'antona G, Granger N, Tonlorenzi R, Innocenzi A, *et al.* Mesoangioblast stem cells ameliorate muscle function in dystrophic dogs. *Nature* 2006; 444:574-579.
 34. Mitchell KJ, Pannerec A, Cadot B, Parlakian A, Besson V, Gomes ER, *et al.* Identification and characterization of a non-satellite cell muscle resident progenitor during postnatal development. *Nat Cell Biol* 2010; 12:257-266.
 35. Sicari BM, Rubin JP, Dearth CL, Wolf MT, Ambrosio F, Boninger M, *et al.* An Acellular Biologic Scaffold Promotes Skeletal Muscle Formation in Mice and Humans with Volumetric Muscle Loss. *Sci Transl Med* 2014; 6:234ra258.
 36. Turner NJ, Yates AJ, Jr., Weber DJ, Qureshi IR, Stolz DB, Gilbert TW, *et al.* Xenogeneic extracellular matrix as an inductive scaffold for regeneration of a functioning musculotendinous junction. *Tissue Eng Part A* 2010; 16:3309-3317.
 37. Camargo FD, Green R, Capetanaki Y, Jackson KAGoodell MA. Single hematopoietic stem cells generate skeletal muscle through myeloid intermediates. *Nat Med* 2003; 9:1520-1527.
 38. Arnold L, Henry A, Poron F, Baba-Amer Y, Van Rooijen N, Plonquet A, *et al.* Inflammatory monocytes recruited after skeletal muscle injury switch into antiinflammatory macrophages to support myogenesis. *J Exp Med* 2007; 204:1057-1069.
 39. Tidball JGVillalta SA. Regulatory interactions between muscle and the immune system during muscle regeneration. *Am J Physiol Regul Integr Comp Physiol* 2010; 298:R1173-1187.
 40. Tidball JGWehling-Henricks M. Macrophages promote muscle membrane repair and muscle fibre growth and regeneration during modified muscle loading in mice in vivo. *J Physiol* 2007; 578:327-336.
 41. Tidball JGVillalta SA. Regulatory interactions between muscle and the immune system during muscle regeneration. *Am J Physiol Regul Integr Comp Physiol*; 298:R1173-1187.
 42. Spiller KL, Anfang RR, Spiller KJ, Ng J, Nakazawa KR, Daulton JW, *et al.* The role of macrophage phenotype in vascularization of tissue engineering scaffolds. *Biomaterials* 2014; 35:4477-4488.
 43. Agrawal H, Tholpady SS, Capito AE, Drake DBKatz AJ. Macrophage phenotypes correspond with remodeling outcomes of various acellular dermal matrices. *Open Access Journal of Regenerative Medicine* 2012; 1:51-59.
 44. Novak ML, Weinheimer-Haus EMKoh TJ. Macrophage activation and skeletal muscle healing following traumatic injury. *J Pathol* 2014; 232:344-355.
 45. Li YP. TNF-alpha is a mitogen in skeletal muscle. *Am J Physiol Cell Physiol* 2003; 285:C370-376.
 46. Sonnet C, Lafuste P, Arnold L, Brigitte M, Poron F, Authier FJ, *et al.* Human macrophages rescue myoblasts and myotubes from apoptosis through a set of adhesion molecular systems. *J Cell Sci* 2006; 119:2497-2507.
 47. Ochoa O, Sun D, Reyes-Reyna SM, Waite LL, Michalek JE, Mcmanus LM, *et al.* Delayed angiogenesis and VEGF production in CCR2-/- mice during impaired skeletal muscle regeneration. *Am J Physiol Regul Integr Comp Physiol* 2007; 293:R651-661.
 48. Warren GL, Hulderman T, Mishra D, Gao X, Millecchia L, O'farrell L, *et al.* Chemokine receptor CCR2 involvement in skeletal muscle regeneration. *FASEB J* 2005; 19:413-415.