AWARD NUMBER: W81XWH-12-1-0407

TITLE: The Replication Stress Response in Pancreatic Cancer

PRINCIPAL INVESTIGATOR: David S. Yu

CONTRACTING ORGANIZATION: Emory University School of Medicine Atlanta, GA 30322

REPORT DATE: December 2014

TYPE OF REPORT: Final Report

PREPARED FOR: U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release; Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DO	CUMENTATION PAGE	Form Approved
Public reporting burden for this collection of information is	estimated to average 1 hour per response, including the time for review	VIVIB INO. U/U4-U188 wing instructions, searching existing data sources, gathering and maintaining the
data needed, and completing and reviewing this collection this burden to Department of Defense, Washington Headd 4302. Respondents should be aware that notwithstanding valid OMB control number. PLEASE DO NOT RETURN	of information. Send comments regarding this burden estimate or any uarters Services, Directorate for Information Operations and Reports (any other provision of law, no person shall be subject to any penalty fr OUR FORM TO THE ABOVE ADDRESS.	r other aspect of this collection of information, including suggestions for reducing 0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202- or failing to comply with a collection of information if it does not display a currently
1. REPORT DATE	2. REPORT TYPE	3. DATES COVERED
December 2014	Final Report	30 Sep 2012 - 29 Sep 2014
4. IIILE AND SUBIIILE The Replication Stress Res	ponse in Pancreatic Cancer	5a. CONTRACT NUMBER
	-	5b. GRANT NUMBER
		W81XWH-12-1-0407
		5c. PROGRAM ELEMENT NUMBER
6. AUTHOR(S)		5d. PROJECT NUMBER
David Yu		
		5e. TASK NUMBER
		5f. WORK UNIT NUMBER
E-Mail: dsyu@emory.edu		
7. PERFORMING ORGANIZATION NAME Emory University School of 1365 Clifton Road NE	(S) AND ADDRESS(ES) Medicine	8. PERFORMING ORGANIZATION REPORT NUMBER
Atlanta, GA 30322		
9. SPONSORING / MONITORING AGENC	Y NAME(S) AND ADDRESS(ES)	10. SPONSOR/MONITOR'S ACRONYM(S)
Fort Detrick Maryland 21702-501		
Tort Detrick, Maryland 21702-5012	F a	11. SPONSOR/MONITOR'S REPORT
		NUMBER(S)
12. DISTRIBUTION / AVAILABILITY STAT	EMENT	
Approved for Public Release; Distr	ibution Unlimited	
13. SUPPLEMENTARY NOTES		
14. ABSTRACT		
Pancreatic cancer is a highly lethal ma	lignancy with an expected 5-year survival of le	ess than 5% for all patients using current therapies.
Most of these therapies rely on inducin	Ig DNA damage and replication blocks to caus	e cell death; nowever the effectiveness of these
treatments is often infined to a subset of	of patients that respond to treatment. Orderstan	patient while potentially reducing toxicity. The
Replication Stress Response (RSR) is	a signaling network that recognizes challenges	to DNA replication and mobilizes diverse activities to
maintain genome integrity. The RSR is	s critical for the prevention of pancreatic cance	r by acting as a barrier against genomic instability and
tumorigenesis. We hypothesized that r	ovel RSR genes maintain genome integrity by	participating in an ATR-mediated replication stress
response and that dysregulation of RSI	R genes and their cancer barrier function result	s in the development of pancreatic cancer. Utilizing a
custom generated siRNA library target	ing genes somatically mutated in pancreatic ca	incer from the Sanger COSMIC library, we completed
a synthetic lethal screen to identify gen	nes which when silenced mediate gemcitabine	sensitivity in human pancreatic cancer cells. We
further validated positive hits be decor	volution of individual siRNAs and characteriz	ed their activities in DNA replication and the DNA
damage response. We determined whe	ther these genes were dysregulated and differe	ntially expressed in pancreatic cancer cell lines and
patient samples using published data s	ets.	
10. SUBJECT TERMS		

Pancreatic cancer, replication stress response, dna damage response, biomarker

 16. SECURITY CLASSIFICATION OF:
 17. LIMITATION
 18. NUMBER
 19a. NAME OF RESPONSIBLE PERSON

 0F ABSTRACT
 0F PAGES
 USAMRMC

a. REPORT	b. ABSTRACT	c. THIS PAGE		19b. TELEPHONE NUMBER (include area
U	U	U	UU	code)

Table of Contents

1. Introduction	1
2. Keywords	2
3. Overall Project Summary	2
4. Key Research Accomplishments	5
5. Conclusion	5
6. Publications, Abstracts, and Presentations	6
7. Inventions, Patents and Licenses	10
8. Reportable Outcomes	10
9. Other Achievements	10
10. References	10
11. Appendices	11

Introduction

Pancreatic cancer is a highly lethal malignancy with an expected 5-year survival of less than 5% for all patients using current therapies. Most of these therapies rely on inducing DNA damage and interfering with DNA replication to cause cell death: however the effectiveness of these treatments is often limited to a subset of patients that respond to treatment. Understanding why patients respond or do not respond to specific treatments would allow the personalization of therapies that are most effective for a patient while potentially reducing toxic side effects. The Replication Stress Response (RSR) is a signaling pathway that recognizes challenges to DNA replication and mobilizes diverse activities to maintain genome integrity. The RSR is critical to prevent pancreatic cancer. In human pre-cancerous lesions, aberrant DNA replication induces activation of the RSR, which maintains genome integrity or causes cell death. Mutations in the RSR promote the survival and proliferation of genetically unstable cells ultimately resulting in cancer. However, the genetic changes that lead to pancreatic cancer can also weaken the ability of cancer cells to respond to treatment by compromising DNA repair pathways. Often the cancer cell will become reliant on backup pathways, which can be targeted to cause cell death through the principle of synthetic lethality. Two genes or pathways are synthetically lethal when inactivation of one is sublethal but inactivation of both causes cell death. We hypothesize that novel RSR genes maintain genome integrity and that deregulation of RSR genes and their cancer barrier function results in the development of pancreatic cancer. However, the genetic changes that drive pancreatic cancer progression also cause susceptibility to synthetic lethal cancer treatments, which can be exploited to personalize therapy. To test this hypothesis, we propose the following specific aims: 1) Identify RSR genes which mediate sensitivity to the first-line chemotherapeutic agent gemcitabine in pancreatic cancer cells. 2) Determine the activities of RSR proteins in DNA replication and DNA damage responses. 3) Evaluate if RSR genes can function as biomarkers for response in pancreatic cancer. Completion of these aims will provide new insights into how the RSR maintains genome integrity, elucidate novel targets for the treatment of pancreatic cancer, and identify subsets of pancreatic cancers that may benefit from gemcitabine chemotherapy.

Keywords

Pancreatic cancer, replication stress response, dna damage response, dna replication, biomarker

Overall Project Summary

We completed a loss of function genetic screen to identify genes, which when silenced cause sensitization or resistance to a low dose of gemcitabine, in human pancreatic cancer cells. We reasoned that genes involved in the RSR would likely be involved in the ATR signaling pathway. We therefore optimized a high-throughput assay using ATR or CHK1 siRNA oligonucleotides as positive controls and ATM or non-targeting (NT) oligonucleotides as negative controls with cell proliferation as a read-out (Fig. 1A-B). The primary screen was completed in MIA PaCa-2 cells, which consistently gave the highest signal to noise ratio among several tested cell types (data not shown). Briefly, cells were reverse transfected with pools of 4 siRNAs targeting a unique sequence of each gene arrayed in a one gene/one well format in 96-well plates. Forty-eight hours after transfection, cells were treated with or without 13 nM gemcitabine for 72 hours prior to assaying for cell proliferation using WST-1 reagent. Each plate contained two positive controls (ATR and CHK1) and several negative controls (NT), and plate-to-plate variability was controlled by normalizing the values on each plate to the average of the negative control values on that plate. We completed three replicas of the primary screen using a

library of 1540 siRNAs, corresponding to four unique siRNA duplexes, targeting each of 385 unique human genes (Fig. 1C) somatically mutated in pancreatic cancer from the Sanger COSMIC database¹ (Task 1-1). Positive hits included a number of genes previously linked to DNA replication and/or DNA damage responses, including FANCI, BRCA2, PKP2, and CTBP2. demonstrating that our screen can yield genes involved in the RSR.

We performed bioinformatic analyses of our positive hits including cross-referencing with published putative ATM/ATR substrates² and published DNA damage sensitivity screens and putative ATM/ATR substrates3-6 (Task 3-1). Using these criteria, we selected 20 genes for further analysis in a secondary screen by deconvoluting with individual siRNAs (Task 1-2). Ten of our genes had at least 2 siRNAs causing sensitivity phenotypes, including chromodomain helicase DNA binding protein 7 (CHD7), which we focused on for further analysis. Four of four siRNAs targeting CHD7 caused gemcitabine sensitization (Fig. 2A). Western blot analysis confirmed



Fig. 1. Gemcitabine sensitivity screen. A. Primary screen flow diagram. MIA PaCa-2 cells were transfected with siRNA (pools of 4 siRNAs per gene) in a one gene/one well format in 96-well plates. 48 hours after transfection, cells were treated with or without 13 nM gemcitabine for 72 hours prior to assaying for cell proliferation using WST-1 reagent. In each plate, non-targeting (NT) targeting, ATR, and CHK1 siRNA were used as controls. Plate-to-plate variability was controlled by normalizing the values on that plate to the average of the negative controls on that plate. B. Optimization of cell viability assay using ATR, ATM, CHK1, or NT siRNA. The viability ratio of gemcitabine treated to untreated is shown. Mean and standard deviation from three replicas is shown. C. Results of primary screen. The primary screen was completed in triplicate. The viability ratio of treated versus untreated cells is shown.

decreased levels of CHD7 following siRNA knockdown as well as specificity of the anti-CHD7 antibody used for immunohistochemistry (IHC) analysis (Fig. 2B) (Task 3-3). A similar gemcitabine sensitization after CHD7 silencing was observed using a range of gemcitabine concentrations and in BxPC-3 and HPAC pancreatic cancer cells, suggesting that the phenotype is not cell-type specific (data not shown) (Task 1-3). We also determined the gemcitabine sensitivity of CHD7 depleted cells using a colony formation assay. MIA PaCa-2 cells silenced for CHD7 demonstrated a significantly reduced percentage of surviving colonies following a 24 hour pulse of gemcitabine sensitization of CHD7 depleted cells using a colony formation assay. MIA PaCa-2 cells colonies following a 24 hour pulse of gemcitabine in a dose-dependent manner compared to a NT control (Fig. 2C), confirming the gemcitabine sensitization of CHD7 depleted cells observed with WST-1 reagent. Consistent with these findings, MIA PaCa-2 and BxPC3 pancreatic cancer cells, which express lower levels of CHD7 than HPAC, CAPAN-1, and AsPC-1 pancreatic cancer cells, demonstrated increased gemcitabine sensitivity (Fig. 2D-E), suggesting that CHD7 expression may predict response to gemcitabine in pancreatic cancer cells (Task 3-5, 3-6).



Fig 2. CHD7 knockdown causes gemcitabine sensitization. A. Four siRNAs targeting CHD7 caused gemcitabine sensitization in MIA PaCa-2 cells. Treated versus untreated percent viability was calculated and the mean and standard deviation from three replicas is shown. * indicates p < 0.05. B. Western blot analysis demonstrating efficiency of CHD7 knockdown with indicated siRNAs. C. Clonogenic assay demonstrating gemcitabine sensitization with CHD7 silencing. MIA PaCa-2 cells transfected with siRNA against CHD7, ATR, or NT were seeded for colony formation, treated with indicated concentrations of gemcitabine for 24 hours, and assayed for surviving colonies 8-12 days later. Percent survival of colonies from treated versus untreated cells is indicated. Mean and standard deviation from three replicas are shown. * indicates p < 0.05. D. Western blot analysis of cell lysate from MIA PaCa-2, HPAC, CAPAN-1, BxPC-3, and AsPC-1 cells with the indicated ant bodies. The CHD7:GAPDH ratio of representative blot from three independent experiments is shown. E. Gemcitabine sensitivity of MIA PaCa-2, HPAC, CAPAN-1, BxPC-3, and AsPC-1 cells following treatment with indicated concentrations of gemcitabine for 72 hours is shown. * indicates p < 0.05.

The gemcitabine hypersensitivity of CHD7 depleted cells suggests that CHD7 may function in the RSR. CHD7 silencing significantly increased the percentage of cells staining with γH2AX, a marker for DNA damage, following treatment with gemcitabine (Fig. 3A), suggesting that CHD7 silencing potentiates gemcitabine-induced DNA damage (Task 2). However, no significant difference in repair kinetics was observed between cells silenced with CHD7 compared with a NT siRNA (Fig. 3A) (Task 2). To determine if CHD7 is involved in cell cycle progression, we examined cells for incorporation of BrdU using flow cytometry for DNA content (Task 2-1). CHD7 silenced cells showed a decreased percentage of cells in S-phase and an increased percentage of cells in G2/M phase compared with control cells (Fig. 3B). To determine if CHD7 functions in ATR-dependent signaling in response to gemcitabine treatment, we examined cells for the phosphorylation of CHK1 Ser317. CHD7 silencing significantly reduced CHK1 Ser317 phosphorylation but not total CHK1 protein levels in response to gemcitabine treatment (Fig. 3C), suggesting that CHD7 functions in controlling ATR dependent

phosphorylation of CHK1 in response to gemcitabine treatment (Task 2-4). To determine if CHD7 functions in DNA damage responses, we examined cells for S-phase and G2/M checkpoint induction following ionizing radiation but found no evidence that CHD7 is involved in the S-phase or G2/M checkpoint (data not shown) (Task 2-2, Task 2-3). We also performed mass spectrometry of purified HA-CHD7 expressed in cells (Task 2-5, Task 2-8) to examine for potential interacting partners that may function in the RSR but found no evidence that CHD7 interacts in a complex with known RSR proteins (data not shown)



Fig 3. CHD7 is a DNA dam age response protein. A. MIA PaCa-2 cells were treated with or without gemcitabine for 20 hours, washed, released for the indicated time points, and processed for yH2AX staining by indirect immunofluorescence. The percentage (mean and standard deviation) of yH2AX positive cells from three replicas is shown. * indicates p < 0.05 **B.** MIA PaCa-2 cells were transfected with NT, CHD7, or ATR siRNA, labeled with BrdU, fixed, and processed for BrdU incorporation and DNA content by flow cytometry. Mean and standard deviation of gated cells from three replicas is shown. **C.** Western blot analysis of cell lysate from MIA PaCa-2 cells treated with or without gemcitabine for 6 hours and probed with anti-CHD7, pCHK-1 Ser317, CHK1, and GAPDH antibodies.

Key Research Accomplishments

- 1. Identified novel RSR genes that are critical for determining gemcitabine sensitivity in pancreatic cancer.
- Characterized functions of novel RSR proteins, including showing that CHD7 functions in the ATR signaling pathway
- 3. Determined that expression of CHD7 can function as a biomarker for gemcitabine sensitivity in cells

Conclusion

We identified a number of novel RSR genes that are critical for determining gemcitabine sensitivity in cells and tumors, including CHD7. We determined that CHD7 functions as a novel replication stress response protein in the ATR signaling pathway. We also found that CHD7

expression is associated with gemcitabine sensitivity in pancreatic cancer cells. We are now working to determine whether expression of CHD7 and other genes identified in our screen can be utilized as prognostic and predictive biomarkers to personalize treatment for patients with pancreatic cancer.

"Publications, Abstracts, Presentations"

Manuscripts:

- 1. Lay Press: Nothing to report
- 2. Peer-Reviewed Scientific Journals:

Hall, W.A., Petrova, A.V., Colbert, L.E., Hardy, C.W., Fisher, S.B., Saka, B., Shelton, J.W., Warren, M.D., Pantazides, B.G., Gandhi, K., Kowalski, J., Kooby, D.A., El-Rayes, B.F., Staley, C.A., Adsay, N.V., Curran, W.J., Landry, J.C., Maithel, S.K., **Yu, D.S.** 2013. Low CHD5 expression activates the DNA damage response and predicts poor outcome in patients undergoing adjuvant therapy for resected pancreatic cancer. **Oncogene**, DOI:10.1038/onc.2013.488.

Colbert, L.E., Petrova, A.V., Fisher, S.B., Pantazides, B.G., Madden, M.Z., Hardy, C.W., Warren, M.D., Pan, Y., Nagaraju, G.P., Liu, E.A., Saka, B., Hall, W.A., Shelton, J.W., Gandhi, K., Pauly, R., Kowalski, J., Kooby, D.A., El-Rayes, B.F., Staley, C.A., Adsay, N.V., Curran, W.J., Landry, J.C., Maithel, S.K., **Yu, D.S.** 2014. CHD7 expression predicts survival outcomes in patients with resected pancreatic cancer. **Cancer Research**, 74(10), 2677-87. PMCID: PMC4025946.

- 3. Invited Articles: Nothing to report
- 4. Abstracts:

Landry, J.C., Pantazides, B.G., Park, W., Shelton, J.W., Maithel, S.K., El-Rayes, B., Yu, D.S. 2012. A synthetic lethal screen identifies genetic determinants for gemcitabine sensitivity in pancreatic cancer. American Association for Cancer Research (AACR) Annual Meeting, Chicago, IL. Poster presentation.

Hardy, C.W., Pantazides, B.G., Gandhi, K., Landry, J., Shelton, J., Maithel, S., El-Rayes B., Kowalski, J., Yu, D.S. 2012. A loss of function genetic screen identifies determinants of gemcitabine sensitivity in pancreatic cancer. AACR/Pancreatic Cancer: Progress and Challenges Meeting, Lake Tahoe, NV. Poster presentation.

Pantazides, B.G., Hardy, C.W., Gandhi, K., Landry, J., Shelton, J., Maithel, S., El-Rayes B., Kowalski, J., Yu, D.S. 2012. A synthetic lethal screen identifies genes that mediate gemcitabine resistance in pancreatic cancer. Radiation Research Society Annual Meeting, San Juan, Puerto Rico. Poster presentation.

Yu, D.S., Pantazides, B.G., Hardy, C.W., Gandhi, K., Landry, J., Shelton, J., Maithel, S., El-Rayes B., Kowalski, J. 2012. A synthetic lethal screen identifies genes that mediate

gemcitabine resistance in pancreatic cancer. American Society for Therapeutic Radiology and Oncology (ASTRO) 54th Annual Meeting, Boston, MA. Oral presentation.

Warren, M.D., Hardy, C.W., Pantazides, B.G., Landry, J.C., Shelton, J.W., Maithel, S.K., El-Rayes, B., Kowalski, J, Yu, D.S. 2013. A synthetic lethal screen identifies determinants of gemcitabine sensitivity in pancreatic cancer. AACR Annual Meeting, Washington, DC. Poster presentation.

Colbert, L.C., Hall, W.A., Hardy, C.W., Fisher, S.B., Yu, D.S., Maithel, S.K., El-Rayes, B., Saka, B., Adsay, N.V., Petrova, A., Pantazides, B., Kowalski, J., Gandhi, K., Kooby, D., Staley, C., Landry, J.C. 2013. A pilot clinical trial of chromodomain-helicase-DNAbinding protein 7 (CHD7) expression as a prognostic and predictive biomarker in patients with early-stage pancreatic adenocarcinoma. AACR Annual Meeting, Washington, DC. Poster presentation.

Colbert, L.E., Fisher, S.B., Hardy, C.W., Hall, W.A., Saka, B., Shelton, J.W., Petrova, A.V., Warren, M.D., Pantazides, B.G., Gandhi, K., Kowalski, J, Kooby, D.A., El-Rayes, B.F., Staley, C.A., Adsay, N.V., Curran, W.J., Landry, J.C., Maithel, S.K., Yu, D.S. 2013. Understanding and exploiting the replication stress response in pancreatic cancer. AACR Annual Meeting, Washington DC. Oral presentation.

Colbert, L.E., Fisher, S.B., Hardy, C.W., Hall, W.A., Saka, B., Shelton, J.W., Petrova, A.V., Warren, M.D., Pantazides, B.G., Gandhi, K., Kowalski, J, Kooby, D.A., El-Rayes, B.F., Staley, C.A., Adsay, N.V., Curran, W.J., Landry, J.C., Maithel, S.K., Yu, D.S. 2013. A rationale-driven approach using a gemcitabine sensitivity screen identifies CHD7 expression as a predictor of outcome in patients with resected pancreatic cancer. Radiation Research Society Annual Meeting, New Orleans, LA. Poster presentation.

Petrova, A.V., Colbert, L.E., Fisher, S.B., Hardy, C.W., Hall, W.A., Saka, B., Shelton, J.W., Warren, M.D., Pantazides, B.G., Gandhi, K., Kowalski, J, Kooby, D.A., El-Rayes, B.F., Staley, C.A., Adsay, N.V., Curran, W.J., Landry, J.C., Maithel, S.K., Yu, D.S. 2013. A synthetic lethal screen identifies determinants of gemcitabine sensitivity in pancreatic cancer. Radiation Research Society Annual Meeting, New Orleans, LA. Poster presentation.

Hall, W.A., Petrova, A.V., Colbert, L.E., Hardy, C.W., Fisher, S.B., Saka, B., Shelton, J.W., Warren, M.D., Pantazides, B.G., Gandhi, K., Kowalski, J., Kooby, D.A., El-Rayes, B.F., Staley, C.A., Adsay, N.V., Curran, W.J., Landry, J.C., Maithel, S.K., Yu, D.S. 2013. Low CHD5 expression activates the DNA damage response and predicts poor outcome in patients undergoing adjuvant therapy for resected pancreatic cancer. Radiation Research Society Annual Meeting, New Orleans, LA. Poster presentation.

Colbert, L.E., Fisher, S.B., Hardy, C.W., Hall, W.A., Saka, B., Shelton, J.W., Petrova, A.V., Warren, M.D., Pantazides, B.G., Gandhi, K., Kowalski, J, Kooby, D.A., El-Rayes, B.F., Staley, C.A., Adsay, N.V., Curran, W.J., Landry, J.C., Maithel, S.K., Yu, D.S. 2013. A rationale-driven approach using a gemcitabine sensitivity screen identifies CHD7 expression as a predictor of outcome in patients with resected pancreatic cancer. American Society for Therapeutic Radiology and Oncology (ASTRO) 55th Annual Meeting, Atlanta, GA. Oral presentation.

Hall, W.A., Petrova, A.V., Colbert, L.E., Hardy, C.W., Fisher, S.B., Saka, B., Shelton, J.W., Warren, M.D., Pantazides, B.G., Gandhi, K., Kowalski, J., Kooby, D.A., El-Rayes, B.F., Staley, C.A., Adsay, N.V., Curran, W.J., Landry, J.C., Maithel, S.K., Yu, D.S. 2013. Low CHD5 expression activates the DNA damage response and predicts poor outcome in patients undergoing adjuvant therapy for resected pancreatic cancer. American Society for Therapeutic Radiology and Oncology (ASTRO) 55th Annual Meeting, Atlanta, GA. Poster presentation.

Colbert, L.E., Fisher, S.B., Hardy, C.W., Hall, W.A., Saka, B., Shelton, J.W., Petrova, A.V., Warren, M.D., Pantazides, B.G., Gandhi, K., Kowalski, J, Kooby, D.A., El-Rayes, B.F., Staley, C.A., Adsay, N.V., Curran, W.J., Landry, J.C., Maithel, S.K., Yu, D.S. 2013. Rationale-Driven Identification of Novel Gemcitabine Sensitivity Genes as Biomarkers for Outcome in Pancreatic Cancer Patients by Exploiting the Replication Stress Response. 3rd World Congress on Cancer Science and Therapy, San Francisco, CA. Oral Presentation.

Presentations:

Landry, J.C., Pantazides, B.G., Park, W., Shelton, J.W., Maithel, S.K., El-Rayes, B., Yu, D.S. 2012. A synthetic lethal screen identifies genetic determinants for gemcitabine sensitivity in pancreatic cancer. American Association for Cancer Research (AACR) Annual Meeting, Chicago, IL. Poster presentation.

Pantazides, B.G., Hardy, C.W., Gandhi, K., Landry, J., Shelton, J., Maithel, S., El-Rayes B., Kowalski, J., Yu, D.S. 2012. A synthetic lethal screen identifies genes that mediate gemcitabine resistance in pancreatic cancer. Winship Scientific Research and Academic Development Symposium. Atlanta, GA. Poster presentation

Hardy, C.W., Pantazides, B.G., Gandhi, K., Landry, J., Shelton, J., Maithel, S., El-Rayes B., Kowalski, J., Yu, D.S. 2012. A loss of function genetic screen identifies determinants of gemcitabine sensitivity in pancreatic cancer. AACR/Pancreatic Cancer: Progress and Challenges Meeting, Lake Tahoe, NV. Poster presentation.

Pantazides, B.G., Hardy, C.W., Gandhi, K., Landry, J., Shelton, J., Maithel, S., El-Rayes B., Kowalski, J., Yu, D.S. 2012. A synthetic lethal screen identifies genes that mediate gemcitabine resistance in pancreatic cancer. Radiation Research Society Annual Meeting, San Juan, Puerto Rico. Poster presentation.

Yu, D.S., Pantazides, B.G., Hardy, C.W., Gandhi, K., Landry, J., Shelton, J., Maithel, S., El-Rayes B., Kowalski, J. 2012. A synthetic lethal screen identifies genes that mediate gemcitabine resistance in pancreatic cancer. American Society for Therapeutic Radiology and Oncology (ASTRO) 54th Annual Meeting, Boston, MA. Oral presentation.

Warren, M.D., Hardy, C.W., Pantazides, B.G., Landry, J.C., Shelton, J.W., Maithel, S.K., El-Rayes, B., Kowalski, J, Yu, D.S. 2013. A synthetic lethal screen identifies determinants of gemcitabine sensitivity in pancreatic cancer. AACR Annual Meeting, Washington, DC. Poster presentation.

Colbert, L.C., Hall, W.A., Hardy, C.W., Fisher, S.B., Yu, D.S., Maithel, S.K., El-Rayes, B., Saka, B., Adsay, N.V., Petrova, A., Pantazides, B., Kowalski, J., Gandhi, K., Kooby, D., Staley, C., Landry, J.C. 2013. A pilot clinical trial of chromodomain-helicase-DNA-

binding protein 7 (CHD7) expression as a prognostic and predictive biomarker in patients with early-stage pancreatic adenocarcinoma. AACR Annual Meeting, Washington, DC. Poster presentation.

Colbert, L.E., Fisher, S.B., Hardy, C.W., Hall, W.A., Saka, B., Shelton, J.W., Petrova, A.V., Warren, M.D., Pantazides, B.G., Gandhi, K., Kowalski, J, Kooby, D.A., El-Rayes, B.F., Staley, C.A., Adsay, N.V., Curran, W.J., Landry, J.C., Maithel, S.K., Yu, D.S. 2013. Understanding and exploiting the replication stress response in pancreatic cancer. AACR Annual Meeting, Washington DC. Oral presentation.

Colbert, L.E., Fisher, S.B., Hardy, C.W., Hall, W.A., Saka, B., Shelton, J.W., Petrova, A.V., Warren, M.D., Pantazides, B.G., Gandhi, K., Kowalski, J, Kooby, D.A., El-Rayes, B.F., Staley, C.A., Adsay, N.V., Curran, W.J., Landry, J.C., Maithel, S.K., Yu, D.S. 2013. A rationale-driven approach using a gemcitabine sensitivity screen identifies CHD7 expression as a predictor of outcome in patients with resected pancreatic cancer. Radiation Research Society Annual Meeting, New Orleans, LA. Poster presentation.

Petrova, A.V., Colbert, L.E., Fisher, S.B., Hardy, C.W., Hall, W.A., Saka, B., Shelton, J.W., Warren, M.D., Pantazides, B.G., Gandhi, K., Kowalski, J, Kooby, D.A., El-Rayes, B.F., Staley, C.A., Adsay, N.V., Curran, W.J., Landry, J.C., Maithel, S.K., Yu, D.S. 2013. A synthetic lethal screen identifies determinants of gemcitabine sensitivity in pancreatic cancer. Radiation Research Society Annual Meeting, New Orleans, LA. Poster presentation.

Hall, W.A., Petrova, A.V., Colbert, L.E., Hardy, C.W., Fisher, S.B., Saka, B., Shelton, J.W., Warren, M.D., Pantazides, B.G., Gandhi, K., Kowalski, J., Kooby, D.A., El-Rayes, B.F., Staley, C.A., Adsay, N.V., Curran, W.J., Landry, J.C., Maithel, S.K., Yu, D.S. 2013. Low CHD5 expression activates the DNA damage response and predicts poor outcome in patients undergoing adjuvant therapy for resected pancreatic cancer. Radiation Research Society Annual Meeting, New Orleans, LA. Poster presentation.

Colbert, L.E., Fisher, S.B., Hardy, C.W., Hall, W.A., Saka, B., Shelton, J.W., Petrova, A.V., Warren, M.D., Pantazides, B.G., Gandhi, K., Kowalski, J, Kooby, D.A., El-Rayes, B.F., Staley, C.A., Adsay, N.V., Curran, W.J., Landry, J.C., Maithel, S.K., Yu, D.S. 2013. A rationale-driven approach using a gemcitabine sensitivity screen identifies CHD7 expression as a predictor of outcome in patients with resected pancreatic cancer. American Society for Therapeutic Radiology and Oncology (ASTRO) 55th Annual Meeting, Atlanta, GA. Oral presentation.

Hall, W.A., Petrova, A.V., Colbert, L.E., Hardy, C.W., Fisher, S.B., Saka, B., Shelton, J.W., Warren, M.D., Pantazides, B.G., Gandhi, K., Kowalski, J., Kooby, D.A., El-Rayes, B.F., Staley, C.A., Adsay, N.V., Curran, W.J., Landry, J.C., Maithel, S.K., Yu, D.S. 2013. Low CHD5 expression activates the DNA damage response and predicts poor outcome in patients undergoing adjuvant therapy for resected pancreatic cancer. American Society for Therapeutic Radiology and Oncology (ASTRO) 55th Annual Meeting, Atlanta, GA. Poster presentation.

Colbert, L.E., Fisher, S.B., Hardy, C.W., Hall, W.A., Saka, B., Shelton, J.W., Petrova, A.V., Warren, M.D., Pantazides, B.G., Gandhi, K., Kowalski, J, Kooby, D.A., El-Rayes, B.F., Staley, C.A., Adsay, N.V., Curran, W.J., Landry, J.C., Maithel, S.K., Yu, D.S. 2013. Rationale-Driven Identification of Novel Gemcitabine Sensitivity Genes as Biomarkers

for Outcome in Pancreatic Cancer Patients by Exploiting the Replication Stress Response. 3rd World Congress on Cancer Science and Therapy, San Francisco, CA. Oral Presentation.

Petrova, A.V., Smith, S.C., Madden, M.Z., Wang, H., Pan, Y., Warren, M.D., Hardy, C.W., Liang, D., Liu, E.A., Robinson, M.H., Rudra, S., Wang, J., Ehdaivand, S., Torres, M.A., Wang, Y., Yu, D.S. 2014. A Gemcitabine Sensitivity Screen identifies a Role for NEK9 in the Replication Stress Response. Radiation Research Society Annual Meeting, Las Vegas, NM. Poster Presentation.

"Pancreatic Cancer Research and Challenges," Seminar, Winship Cancer Institute Pancreatic Cancer Support Group, January, 2013.

"Understanding and Exploiting the Replication Stress Response in Cancer," Seminar, Division of Cancer Biology Seminary Series, Emory University School of Medicine, February, 2013.

"Biomarker Analysis of Gemcitabine Sensitivity Genes in Pancreatic Cancer," Radiation Therapy Oncology Group Semi-Annual Meeting, Philadelphia, PA, June, 2013.

"Biomarker Analysis of Gemcitabine Sensitivity Genes in Pancreatic Cancer," Radiation Therapy Oncology Group Semi-Annual Meeting, Chicago, IL, July, 2014.

"Inventions, Patents, Licenses"

Nothing to report

Reportable Outcomes

Nothing to report

Other Achievements

Nothing to report

References

1. Forbes, S. A. *et al.* The Catalogue of Somatic Mutations in Cancer (COSMIC). *Curr Protoc Hum Genet* **Chapter 10**, Unit 10 11, doi:10.1002/0471142905.hg1011s57 (2008).

2. Matsuoka, S. *et al.* ATM and ATR substrate analysis reveals extensive protein networks responsive to DNA damage. *Science* **316**, 1160-1166 (2007).

3. Cotta-Ramusino, C. *et al.* A DNA damage response screen identifies RHINO, a 9-1-1 and TopBP1 interacting protein required for ATR signaling. *Science* **332**, 1313-1317, doi:332/6035/1313 [pii]10.1126/science.1203430.

4. Paulsen, R. D. *et al.* A genome-wide siRNA screen reveals diverse cellular processes and pathways that mediate genome stability. *Mol Cell* **35**, 228-239, doi:S1097-2765(09)00459-6 [pii] 10.1016/j.molcel.2009.06.021 (2009).

5. Smogorzewska, A. *et al.* Identification of the FANCI protein, a monoubiquitinated FANCD2 paralog required for DNA repair. *Cell* **129**, 289-301, doi:S0092-8674(07)00320-0 [pii]10.1016/j.cell.2007.03.009 (2007).

6. Wang, B., Hurov, K., Hofmann, K. & Elledge, S. J. NBA1, a new player in the Brca1 A complex, is required for DNA damage resistance and checkpoint control. *Genes Dev* **23**, 729-739, doi:gad.1770309 [pii]10.1101/gad.1770309 (2009).

Appendices

CHD7 Expression Predicts Survival Outcomes in Patients with Resected Pancreatic Cancer

Lauren E. Colbert¹, Aleksandra V. Petrova¹, Sarah B. Fisher⁴, Brooke G. Pantazides¹, Matthew Z. Madden¹, Claire W. Hardy¹, Matthew D. Warren¹, Yunfeng Pan¹, Ganji P. Nagaraju², Elaine A. Liu¹, Burcu Saka³, William A. Hall¹, Joseph W. Shelton¹, Khanjan Gandhi⁵, Rini Pauly⁵, Jeanne Kowalski⁵, David A. Kooby^{4,6}, Bassel F. El-Rayes², Charles A. Staley III⁴, N. Volkan Adsay³, Walter J. Currari Jr¹, Jerome C. Landry¹, Shishir K. Maithel⁴, and David S. Yu¹

Abstract

Pancreatic ductal adenocarcinoma (PDAC) is a devastating disease with poor outcomes with current therapies. Gemcitabine is the primary adjuvant drug used clinically, but its effectiveness is limited. In this study, our objective was to use a rationale driven approach to identify novel biomarkers for outcome in patients with early stage resected PDAC treated with adjuvant gemcitabine. Using a synthetic lethal screen in human PDAC cells, we identified 93 genes, including 55 genes linked to DNA damage responses (DDR), that demonstrated gemcitabine sensitization when silenced, including CHD7, which functions in chromatin remodeling, CHD7 depletion sensitized PDAC cells to gemcitabine and delayed their growth in tumor xenografts. Moreover, CHD7 silencing impaired ATR dependent phosphorylation of CHK1 and increased DNA damage induced by gemcitabine. CHD7 was dysregulated, ranking above the 90th percentile in differential expression in a panel of PDAC clinical specimens, highlighting its potential as a biomarker. Immunohistochemical analysis of specimens from 59 patients with resected PDAC receiving adjuvant gemcitabine revealed that low CHD7 expression was associated with increased recurrence free survival (RFS) and overall survival (OS), in univariate and multivariate analyses. Notably, CHD7 expression was not associated with RFS or OS for patients not receiving gemcitabine. Thus, low CHD7 expression was correlated selectively with gemcitabine sensitivity in this patient population. These results supported our rationale driven strategy to exploit dysregulated DDR pathways in PDAC to identify genetic determinants of gemcitabine sensitivity, identifying CHD7 as a novel biomarker candidate to evaluate further for individualizing PDAC treatment. Cancer Res; 74(10); 2677 87. ©2014 AACR.

Introduction

Pancreatic adenocarcinoma has a poor prognosis with a 5 year overall survival (OS) rate around 5% (1). Patients with early stage pancreatic adenocarcinoma who undergo resection demonstrate the best prognosis, particularly when resection is followed by adjuvant chemotherapy with or without radiother apy (2, 3). Still, recurrence is common and OS remains poor even for patients who undergo complete resection and adju vant therapy. Recent developments have suggested that pan creatic adenocarcinoma is a genetically heterogeneous disease

Note: Supplementary data for this article are available at Cancer Research Online (http://cancerres.aacrjournals.org/).

Corresponding Author: David S. Yu, Department of Radiation Oncology, Emory University School of Medicine, 1365 Clifton Road NE, C3008 Atlanta, Georgia 30322. Phone: 404 778 1758; Fax: 404 778 5520; E mail: dsyu@emory.edu

doi: 10.1158/0008 5472.CAN 13 1996

©2014 American Association for Cancer Research.

(4) and, as such, patients may benefit from the identification of predictive biomarkers for responsiveness to adjuvant therapy.

Gemcitabine is the primary chemotherapeutic agent used to treat patients with pancreatic adenocarcinoma in the adjuvant setting (2, 5). The cytotoxic effects of gemcitabine are mediated in part through incorporation into DNA as a terminal nucle oside analog and in part through inhibition of ribonucleotide reductase, which depletes nucleotides required for DNA syn thesis. However, the efficacy of gemcitabine for pancreatic adenocarcinoma is limited. A better understanding of which patients are likely to respond to gemcitabine treatment would facilitate personalization of therapy and optimize the clinical benefit to toxicity ratio associated with adjuvant therapy.

The DNA damage response (DDR) pathway is critical for the maintenance of genome integrity and serves as a cancer barrier by mobilizing DNA repair, cell cycle arrest, and/or apoptosis (6, 7). In human precancerous lesions, aberrant DNA replica tion induces DDR activation, which constrains tumor devel opment. Thus, the DDR acts as a barrier against genomic instability and cancer development. Tumor cells may in turn develop mutations or epigenetic silencing of protective DDR genes, leading to the proliferation of genetically unstable cells and ultimately resulting in cancer. Indeed, a large number of DDR genes are somatically mutated in pancreatic

Authors' Affiliations: Departments of ¹Radiation Oncology, ²Medical Oncology, and ³Pathology; ⁴Division of Surgical Oncology, Department of Surgery; ⁵Biostatistics and Bioinformatics Shared Resource, Winship Cancer Institute, Emory University; and ⁶Atlanta Veterans Affairs Medical Center, Atlanta, Georgia

adenocarcinoma, including ATM, BRCA2, CDKN2A, FANCI, HELB, and RAD9 (8). These genetic changes in the DDR pathway can lead to pancreatic adenocarcinoma and can also weaken the ability of cancer cells to respond to treatment by decreasing activity in DNA repair pathways. Often, the cancer cell will become reliant on backup pathways that can be targeted to cause cell death through the principal of synthetic lethality (inactivation of one gene or pathway is sublethal but inactivation of both causes cell death). As such, determining genetic alterations and cancer treatments that are syntheti cally lethal may lead to the identification of novel druggable targets as adjuncts to gemcitabine treatment or novel biomar kers to predict response to gemcitabine therapy. Using this rationale, we sought to exploit dysregulated DDR pathways in pancreatic adenocarcinoma by identifying genetic determi nants that are synthetically lethal with gemcitabine treatment and evaluating their clinical relevance as biomarkers for out come in patients with early stage resected pancreatic adeno carcinoma treated with adjuvant gemcitabine.

Chromodomain helicase DNA binding protein 7 (CHD7), is a member of a family of chromodomain enzymes that belong to the ATP dependent chromatin remodeling protein SNF2 super family. Mutations in CHD7 lead to congenital CHARGE syn drome, named for its characteristic traits: coloboma of the eye, heart defects, atresia of the nasal choanae, retardation of growth and/or development, genital and/or urinary abnormalities, and ear abnormalities and deafness (9), and Kallman Syndrome, a genetic disorder marked by hypogonadotropic hypogonadism and anosmia (10). CHD7 is also dysregulated in 13% to 35% of cases of pancreatic adenocarcinoma, with aberrant expression, copy number variation, and somatic mutations (see Supple mentary Table S3; refs. 11 13). CHD7 helps to regulate neural crest gene expression (14), regulates ribosomal RNA biogenesis (15), and interacts with SOX2 to regulate gene expression (16). CHD7 is also a putative substrate of the ATM/ATR checkpoint kinases, suggesting that it may play a role in the DDR (17, 18). The clinical significance of CHD7 expression in pancreatic adenocarcinoma has not previously been reported.

The purpose of this analysis was to use a rationale driven approach to identify novel biomarkers for outcome in patients with early stage resected pancreatic adenocarcinoma treated with adjuvant gemcitabine (Fig. 1A). We initially completed a synthetic lethal siRNA screen to identify genetic determinants of gemcitabine sensitivity in human pancreatic cancer cells and identified the top 15% of these genes for further analysis. Genes validated by a secondary screen and/or linked to the DDR were then analyzed for dysregulation and differential expression in pancreatic adenocarcinoma by mining published data sets to determine their potential as biomarkers. Finally, we correlated CHD7 gene expression characterized by immunohistochemistry (IHC) with clinical outcome in patients with early stage resected pancreatic adenocarcinoma treated with adjuvant gemcitabine.

Materials and Methods

Cell culture, siRNA, and transfection

MIA PaCa 2 cells were grown in Dulbecco's Modified Eagle Medium (DMEM; Gibco) supplemented with 10% FBS (Gibco) and 2.5% horse serum (Gibco). HPAC cells were grown in 1:1 DMEM:Hams F12 supplemented (Gibco) with 40 ng/mL hydro cortisone, 10 ng/mL EGF, and 5% FBS. BxPC 3 and AsPC 1 cells were grown in RPMI 1640 supplemented with 10% FBS, and CAPAN 1 cells were grown in Iscove's Modified Dulbecco's Medium supplemented with 20% FBS. Cell lines were grown in a humidified incubator at 37°C with 5% carbon dioxide.

Transfections were done using the HiPerFect Transfection Reagent (Qiagen) according to the manufacturer's protocol. Primary and secondary screen siRNAs were purchased from Thermo Scientific. siRNA sequences are listed below.

NT: (ATGAACGTGAATTGCTCAATT) ATR: (CCUCCGUGAUGUUGCUUGA) ATRIP: (GGTCCACAGATTATTAGA) CHK1: (CTGAAGAAGCAGTCGCAGT) CHD7 1: (UAACGUACCUAACCUAUUA) CHD7 2: (CGACAAGGCUAGUUUGAAA) CHD7 3: (GGGAAGCUAUUAUAUCUGA) CHD7 4: (GUAGAUAACCAAGAACUAA)

TRC lentiviral short hairpin RNA (shRNA) was purchased from Thermo Scientific: shControl (RHS4080), shCHD7 1 (1RHS3979 201747986), shCHD7 2 (1RHS3979 201747990).

Gemcitabine sensitivity screen

MIA PaCa 2 cells were transfected in 96 well plates using HiPerFect Transfection Reagent (Qiagen) with 25 nmol/L siRNA from a custom siGENOME siRNA library (Thermo Scientific) of 4,024 siRNAs corresponding to 1,006 unique human nuclear enzyme genes (pools of four siRNAs targeting a unique sequence of each gene) using a one gene per well format. Twenty four hours later, plates were split 1:4, and then treated following another 24 hours with or without 13 nmol/L gemcitabine (Hospira, Inc.) for 72 hours before assaying for cell proliferation using WST 1 reagent (Roche Diagnostics). Each plate contained two positive controls (ATR and CHK1) and several negative controls (NT), and plate to plate variability was controlled by normalizing the values on each plate to the average of the negative control values on that plate. A ratio of gemcitabine treated/untreated viability was calculated and normalized to that of nontargeting siRNA. Principal compo nents analysis (PCA) was used to account for possible vari ability between the cell viability of the three replicates for each gene. These genes were then sorted by increasing average cell viability via PCA, and the top 15% of genes were categorized as possible "hits."

Secondary validation screen

MIA Pa Ca 2, BxPC3, or HPAC cells were transfected in 96 well plates with 25 nmol/L siRNA, split 1:4 24 hours later, and then treated following another 24 hours with or without gemcitabine at IC_5 , IC_{25} , or IC_{50} for 72 hours before assaying for cell proliferation using WST 1 reagent. A ratio of gem citabine treated/untreated viability was calculated and normalized to that of nontargeting siRNA. MIA PaCa 2, HPAC, CAPAN 1, BxPC3, and AsPC 1 cells were treated



Figure 1. Primary gencitabine sensitivity screen. A, flow diagram of approach for identifying novel bionnarkers for outcome in patients with early stageresected pancreatic adenocarcinoma treated with adjuvant gencitabine. B, diagram of primary screen as described in text. C, results of primary screen. The log₂ ratio of treated versus untreated cell viability relative to the nontargeting (NT) siRNA for each gene is shown. Mean from three replicas of primary screen is shown. D, proportion of top 15% of gencitabine sensitivity genes with statistically significant involvement in known pathways. The top 18 pathways identified on network analysis via MetaCore ExPlain Process Network Analysis can be consolidated into the listed categories. Fifty five percent of all identified genes were involved in DDR pathways.

with the indicated concentrations of gemcitabine for 72 hours before assaying for cell proliferation using the WST 1 reagent.

Colony formation assay

Cells were transfected with 25 nmol/L siRNA. Following a 24 hour knockdown, 500 cells were seeded into 6 well plates in triplicate. Cells were allowed to culture overnight and were then treated for 24 hours with increasing concentrations of gemcitabine. Following the gemcitabine incubation, the plates were washed with PBS and fresh media were added for 8 to 12 days before staining colonies with a 0.5% crystal violet (Ampresco) solution.

Western blot analysis

MIA PaCa 2 cells transfected with siRNA for 48 hours or MIA PaCa 2, HPAC, CAPAN 1, BxPC3, and AsPC 1 cells were har vested with NP40 buffer containing 200 mmol/L NaCl, 1% NP40, 50 mmol/L Tris HCl (pH 8.0), and supplemented with fresh protease inhibitors. Samples were loaded into a SDS PAGE gel, transferred to a polyvinylidene difluoride (PVDF) membrane, and subsequently probed with an anti CHD7 antibody (NBP1 77393; Novus Biologicals) and anti GAPDH antibody (GeneTex; GTX627408) followed by LI COR IRDye secondary antibodies. Detection was performed using the Odyssey system (LI COR Biosciences).

To analyze phosphorylation of CHK1, MIA PaCa 2 cells were transfected with 25 nmol/L siRNA for 48 hours and treated with 1 mmol/L gemcitabine for 6 hours. Cells then were harvested, washed with PBS, and lysed in cold RIPA buffer (25 mmol/L Tris HCl pH 7.6, 150 mmol/L NaCl, 1% NP40, 1% sodium deoxycholate, and 0.1% SDS) with protease and phosphatase inhibitors for 30 minutes. Lysates were clarified by centrifugation at 16 $\times g$ for 10 minutes and 150 µg of protein for each sample were used for Western blot analysis. Primary antibody pCHK1 S317 (Cell Signaling Technology; #2344) and CHK1 (Santa Cruz Biotechnology; sc 8408) were used for detection of phosphorylated and total CHK1, respectively.

In vivo tumor growth inhibition assay

Male nude mice were maintained in a pathogen free envi ronment, and all *in vivo* procedures were approved by the Emory University, Institutional Animal Care and Use Commit tee. Stable MIA PaCa 2 shCHD7 2 and MIA PaCa 2 shControl cells ($1.5 \times 10^6/0.1$ mL of 20% Matrigel gel in serum free media) were injected subcutaneously into the flank of 5 week old mice. Mice bearing established tumors (100 to 125 mm³) were randomized into treatment groups of four. Animals were treated on days 0, 7, and 14 via the tail vein with either vehicle or 100 mg/kg gemcitabine. Tumor growth inhibition was determined as described previously (19).

γH2AX DNA damage assay

MIA PaCa 2 cells were treated with or without 13 nmol/L gemcitabine for 20 hours, washed, released for the indicated time points, and processed for YH2AX staining by indirect immunofluorescence. Cells were washed with $1 \times$ PBS, fixed with 2% paraformaldehyde for 10 minutes at room tempera ture, permeabilized with 0.5% Triton X 100 (Fisher Scientific), and blocked in a 5% bovine serum albumin solution (Sigma). Cells were then immunostained with anti phospho histone H2AX (Ser139) antibody (Millipore; 05 636) and anti mouse secondary antibody with AlexaFluor 488 (Invitrogen; A21206). After incubation, cells were mounted onto slides with a mounting media containing 4', 6 diamidino 2 phenylindole (DAPI) and dried. Analysis was performed using a Zeiss Observer Z1 microscope with Axiovision Rel 4.8 software using the $63 \times$ oil objective. Foci quantitation was conducted by counting 250 healthy cells and scoring cells with 10 or more foci as positive. Experiments were done in triplicate.

Cell cycle analysis

MIA PaCa 2 cells were transfected with 25 nmol/L siRNA for 48 hours and treated with or without gemcitabine (13 nmol/L) for 24 hours. After fixing with ice cold 70% ethanol, cells were washed in PBS, and propidium iodide (PI; 25 μ g/mL; Sigma) and RNase A (10 μ g/mL; Qiagen) were added to determine DNA content. Cells were analyzed on a FACSCanto II (BD Bio sciences) and FlowJo software.

Biomarker selection

Gemcitabine sensitivity genes validated on secondary screen or known DDR genes were analyzed for evidence of dysregulation by identifying genes overexpressed in The Compendium of Potential Biomarkers of Pancreatic Cancer (20) or somatically mutated in the pancreatic Catalogue for Somatic Mutations in Cancer (COSMIC) database (11). For determination of differential expression, we extracted expression data from the two Gene Expression Omnibus (GEO) submissions based on the Affymetrix U133 Plus 2.0 platform: (i) GSE12654, a 22 pancreatic cancer cell line study (19); and (ii) GSE16515, 20 pancreatic patient tumors (21, 22). Within each study, after processing and normalization, we performed a genomewide filter to identify genes with "large" expression differences among tumors, and separately, among cell lines, using a variance approach. We define "differential expression" as genes whose expression variabil ity is "large" relative to all other genes on the array, in which "large" is defined according to whether expression variability associated with a gene was greater than the 90th percentile from all genes. We then compared this list of genes with the lists in Supplementary Tables S2 and S4. CHD7 was chosen as a potential biomarker based on evidence of both dysre gulation and differential expression.

IHC patient selection

Patients were selected for this analysis from a prospectively maintained database of patients who underwent resection for early stage pancreatic adenocarcinoma between January 2000 and October 2008; data for these patients have been included in other cohorts previously reported (23 26). These 59 patients received adjuvant chemotherapy with or without adjuvant radiation. The gemcitabine patient population was composed of 42 of these patients who received gemcitabine as a com ponent of the adjuvant chemotherapy regimen. An additional 17 patients received adjuvant chemotherapy with agents other than gemcitabine. OS was calculated from date of surgery to patient death. Recurrence free survival (RFS) was measured on the basis of surveillance imaging obtained at regular intervals after resection. Patient demographics, pathologic character istics, and treatment characteristics were originally collected from pathologic record and chart review. Permission was obtained from the Emory Institutional Review Board 00048816, and patient confidentiality was maintained accord ing to the Health Insurance and Patient Accessibility Act of 1996.

Immunohistochemical analysis

An experienced pathologist identified representative sec tions of tumor and normal tissue from formalin fixed paraffin embedded slides. The tissue was stained using an anti CHD7 mouse monoclonal antibody (NBP1 77393; Novus Biologicals) at a concentration of 1:200. Specificity of the anti CHD7 antibody was validated by Western blot analysis following siRNA silencing (Fig. 2B). An expression score was calculated using a previously defined scoring system (23, 27). Overall score was dichotomized into low (<3.1) and high (>3.1) expression groups for this analysis (Supplementary Fig. S1).



Figure 2. CHD7 knockdown causes gemcitabine sensitization. A, four siRNAs targeting CHD7 caused gemcitabine sensitization in MIA PaCa 2 cells. Treated versus untreated percent viability was calculated and the mean and SD from three replicas is shown; *, *P* < 0.05. B, Western blot analysis demonstrating efficiency of CHD7 knockdown with indicated siRNAs. C, clonogenic assay demonstrating gemcitabine sensitization with CHD7 silencing. MIA PaCa 2 cells transfected with siRNA against CHD7, ATR, or NT were seeded for colony formation, treated with indicated concentrations of gemcitabine for 24 hours, and assayed for surviving colonies 8 to 12 days later. Percent survival of colonies from threated versus untreated cells is indicated. Mean and SD from three replicas are shown; *, *P* < 0.05. D, Western blot analysis of cell lysate from MIA PaCa 2, HPAC, CAPAN 1, BxPC 3, and ASPC 1 cells with the indicated antibodies. The CHD7:GAPDH ratio of representative blot from three independent experiments is shown. E, gemcitabine sensitivity of MIA PaCa 2, HPAC, CAPAN 1, BxPC 3, and ASPC 1 cells with indicated concentrations of gemcitabine for 72 hours is shown. F, Western blot analysis demonstrating efficiency of CHD7 knockdown with indicated schematis as a shown; *, *P* < 0.05. J, western blot analysis of cell lysate from three independent experiments is shown. E, gemcitabine sensitivity of MIA PaCa 2, HPAC, CAPAN 1, BxPC 3, and ASPC 1 cells following treatment with indicated concentrations of gemcitabine for 72 hours is shown. F, Western blot analysis demonstrating efficiency of CHD7 knockdown with indicated shRNAs in MIA PaCa 2 cells. G, athymic nude mice with shCHD7 and shControl MIA PaCa 2 tumor xenografts were treated with or without gemcitabine (100 mg/kg) on days 0, 7, and 14, and tumor growth was measured every 4 days. Mean and SEM from six tumors are shown; *, *P* < 0.05.

Statistical analysis

Descriptive statistics were generated for patient character istics, tumor characteristics, and treatment characteristics. Similar statistical analyses were performed for patients receiv ing adjuvant therapy, patients receiving gemcitabine based therapy, and patients receiving non gemcitabine based ther apy. Kaplan Meier log rank survival analysis was performed to determine prognostic factors for RFS and OS. Univariate and multivariate Cox regression analyses were performed for all patients to examine the correlation of CHD7 expression level on both RFS and OS. Factors examined on univariate analysis included age, sex, ethnicity, receipt of adjuvant and neoadju vant therapy, tumor size, margin status, grade, nodal status, perineural invasion, lymphovascular invasion, receipt of radio therapy, CA19 9 levels, and type of adjuvant chemotherapy. Clinically relevant covariates significant to a level of P < 0.2 on univariate analysis for either RFS or OS were included in the multivariate model; these included tumor size, margin status, nodal status, perineural invasion, lymphovascular invasion, and tumor grade. Data were analyzed using the Statistical Package for the Social Sciences 19.0 software for Windows (IBM).

Results

Gemcitabine sensitivity screen

To identify genetic determinants of gemcitabine sensitivity, we completed a siRNA screen to identify genes that when silenced cause either sensitization or resistance to a low dose of gemcitabine in human pancreatic cancer cells. Because gem citabine induces DNA damage and replication stress, we reasoned that gemcitabine sensitivity genes would likely be involved in the DDR. We, therefore, optimized a high through put assay using ATR and CHK1 siRNA as positive controls and a nontargeting siRNA as a negative control with cell proliferation as a read out (Fig. 1B). The primary screen was completed in MIA PaCa 2 cells, which consistently gave the highest signal to noise ratio among several tested cell types (Supplementary Fig. S3A and data not shown). Briefly, cells were transfected with pools of four siRNAs targeting a unique sequence of each gene arrayed in a one gene per one well format in 96 well plates. Forty eight hours after transfection, cells were treated with or without 13 nmol/L gemcitabine (equivalent to IC_{25} under these conditions, see Fig. 2E) for 72 hours before assaying for cell proliferation using WST 1 reagent. Each plate contained two positive controls (ATR and CHK1) and several negative con trols (NT), and plate to plate variability was controlled by normalizing the values on each plate to the average of the negative control values on that plate. We completed three replicas of the primary screen using a library of 4,024 siRNAs, corresponding to four unique siRNA duplexes, targeting each of 1,006 unique human genes (Fig. 1C). The library consisted predominantly of nuclear enzymes, which we reasoned were more likely to function directly in the DDR and be targetable. Results of the primary screen were ranked by PCA score (Supplementary Table S1). The top 15% of these genes (156 genes) included 55 genes linked to the DDR (Fig. 1D; Supple mentary Fig. S2; and Supplementary Table S2) including well characterized ATR signaling pathway genes ATR, CHK1, RAD9, RAD1, and HUS1 and nucleotide metabolism genes RRM1 and RRM2, known to regulate gemcitabine sensitivity (28), dem onstrating that our screen can yield DDR genes that determine gemcitabine sensitivity.

CHD7 knockdown causes gemcitabine sensitization

Sixty eight of our hits were identified in previously published DNA damage sensitivity screens (17, 20, 29 36) and 27 are putative ATM/ATR substrates (17; Supplementary Table S3). We used these criteria to validate 47 of the 99 hits not characterized in the DDR in a secondary screen using decon voluted individual siRNAs to confirm their gemcitabine sen sitivity and eliminate false positives due to off target effects, and 38 of these genes induced gemcitabine sensitivity in at least two out of four siRNAs tested, including CHD7 (Supple mentary Table S4). Four of 4 siRNAs targeting CHD7 caused gemcitabine sensitization (Fig. 2A). Western blot analysis confirmed decreased levels of CHD7 following siRNA knock down as well as specificity of the anti CHD7 antibody used for IHC analysis (Fig. 2B). A similar gemcitabine sensitization after CHD7 silencing was observed using a range of gemcitabine concentrations and in BxPC 3 and HPAC pancreatic cancer cells, suggesting that the phenotype is not cell type specific (Supplementary Fig. S3A C). CHD7 silencing in the absence of gemcitabine treatment reduced cell viability (Supplementary Fig. S3D). We also determined the gemcitabine sensitivity of CHD7 depleted cells using a colony formation assay. MIA PaCa 2 cells silenced for CHD7 demonstrated a significantly reduced percentage of surviving colonies following a 24 hour pulse of gemcitabine in a dose dependent manner compared with a NT control (Fig. 2C), confirming the gemcitabine sensitization of CHD7 depleted cells observed with WST 1 reagent. Consistent with these findings, MIA PaCa 2 and BxPC3 pancreatic cancer cells, which express lower levels of CHD7 than HPAC, CAPAN 1, and AsPC 1 pancreatic cancer cells, demonstrated increased gemcitabine sensitivity (Fig. 2D and E), suggesting that CHD7 expression may predict response to gemcitabine in pancreatic adenocarcinoma cells. To deter mine if CHD7 silencing causes gemcitabine sensitization of pancreatic cancer tumors in vivo, we generated a xenograft model using MIA PaCa 2 cells stably expressing shCHD7 or shControl (Fig. 2F). CHD7 silencing significantly delayed tumor growth in mice treated with gemcitabine compared with a control treated with gemcitabine (Fig. 2G), suggesting that CHD7 silencing also causes gemcitabine sensitization in vivo. No significant difference in body weight was observed in mice bearing tumors with shCHD7 compared with shControl and treated with or without gemcitabine (Supplementary Fig. S4).

CHD7 is a DDR protein

The gemcitabine hypersensitivity of CHD7 depleted cells suggests that CHD7 may function in the DDR. CHD7 silencing significantly increased the percentage of cells staining with γH2AX, a marker for DNA damage, following treatment with gemcitabine (Fig. 3A), suggesting that CHD7 silencing potenti ates gemcitabine induced DNA damage. However, no signifi cant difference in repair kinetics was observed between cells silenced with CHD7 compared with a nontargeting siRNA (Fig. 3A). CHD7 silenced cells showed a decreased percentage of cells in S phase and an increased percentage of cells in G₂ M in the absence of gemcitabine treatment (Supplementary Fig. S5A); however, no significant difference in cell cycle profile was observed between CHD7 depleted compared with NT control cells following gemcitabine treatment (Supplementary Fig. S5B). There was also no significant difference in protein levels of CHD7 in response to gemcitabine treatment (Supplemen tary Fig. S6). To determine whether CHD7 functions in ATR dependent signaling in response to gemcitabine treatment, we examined cells for the phosphorylation of CHK1 Ser317. CHD7 silencing significantly reduced CHK1 Ser317 phosphorylation but not total CHK1 protein levels in response to gemcitabine treatment (Fig. 3B), suggesting that CHD7 functions in con trolling ATR dependent phosphorylation of CHK1 in response to gemcitabine treatment.

CHD7 is dysregulated and differentially expressed in pancreatic adenocarcinoma

Genes validated by our secondary screen or linked to the DDR were then analyzed for dysregulation and differential expression in pancreatic adenocarcinoma by mining of pub lished data sets to determine their potential as biomarkers. Of these, eight genes demonstrate aberrant expression or somatic mutations in pancreatic adenocarcinoma (Fig. 3C and Sup plementary Table S3) as reported in The Compendium of Potential Biomarkers (20) and the COSMIC database (11). Twelve of the genes are above the 90th percentile in differential expression among a panel of 22 pancreatic adenocarcinoma cell lines or 20 pancreatic adenocarcinoma tissue samples (Fig. 3C and Supplementary Table S3; refs. 21, 22). Four of the genes exhibit both dysregulation and differential expression, includ ing *CHD7*, which was selected for further analysis as a biomarker.

Survival analyses

Patient demographics, pathologic and treatment character istics can be seen in Table 1. CHD7 expression was low in 84.7% of patients. Median tumor size was 3.4 cm (range, 1–6 cm), and 60% of patients were node positive. In addition to CHD7 expression, significant covariates on univariate analysis includ ed tumor size, margin status, lymph node status, perineural invasion (PNI), lymphovascular invasion (LVI), and grade (P < 0.2). On Kaplan Meier analysis for patients receiving gemcitabine as a component of adjuvant therapy (n = 42), low CHD7 expression was associated with increased RFS (15 vs. 7 months; P = 0.025; Fig. 4A) and increased OS (18 vs. 10 months; P = 0.015; Fig. 4B). On multivariate analysis (Table 2), low CHD7



Figure 3. CHD7 is a DDR protein. A, MIA PaCa 2 cells were treated with or without gemcitabine for 20 hours, washed, released for the indicated time points, and processed for γ H2AX staining by indirect immunofluorescence. The percentage (mean and SD) of γ H2AX positive cells from three replicas is shown; *, *P* < 0.05. B, Western blot analysis of cell lysate from MIA PaCa 2 cells treated with or without gemcitabine for 6 hours and probed with anti CHD7, pCHK 1 Ser317, CHK1, and GAPDH antibodies. C, Venn diagram showing gemcitabine sensitivity genes dysregulated and/or above the 90th percentile in differential expression among a panel of pancreatic cancer cell lines and tissue samples.

Table 1. Patient demographics, tumor characteristics, and treatment characteristics for all patients (N = 59)

	Median (range)	N	Percent- age
Patient demographics			
Male sex		31	52.5
Ethnicity			
Asian		2	3.4
Black		12	20.3
White		42	71.2
Age (y)	60.0 (37–84)		
OS (mo)	17.3 (4.8–114.6)		
RFS (mo)	14.5 (0.6–109.8)		
Tumor characteristics			
Positive margins		20	24.5
Grade			
Well differentiated		5	6.3
Moderately		46	57.5
differentiated			
Poorly differentiated		28	35.0
Positive nodes		48	60.0
PNI		70	87.5
LVI		38	47.5
Low CHD7 expression		50	84.7
Tumor size (cm)	3.4 (1–6)		
Treatment characteristics			
Neoadjuvant therapy		2	3.4
Radiation therapy		39	66.1
Received gemcitabine		42	69.5

expression remained associated with increased RFS [HR, 0.12; 95% confidence interval (CI), 0.04 0.42; P = 0.001 and increased OS (HR, 0.09; 95% CI, 0.03 0.29; P < 0.0001). In the subset of patients receiving adjuvant therapy with agents other than gemcitabine [most commonly 5 fluorouracil (5 FU)], CHD7 was not associated with RFS (P = 0.1, data not shown) or OS (P = 0.4, data not shown). On Kaplan Meier analysis for all patients (n = 59), low CHD7 expression via IHC scoring was associated with increased RFS (15 months vs. 7 months; P =0.015; Fig. 4C) and increased OS (19.5 months vs. 9 months; P =0.001; Fig. 4D). These results remained significant on multivar iate analysis (Table 3). To ensure stability of the multivariate model given the small number of events, the three least signif icant factors on univariate analysis were removed from the model (grade, PNI, and LVI) and the significance of CHD7 expression remained unchanged.

Discussion

In this study, we demonstrate a rationale driven approach for identifying novel biomarkers for outcome in patients with early stage resected pancreatic adenocarcinoma treated with adju vant gemcitabine. Using a synthetic lethal screen to identify genetic determinants of gemcitabine sensitivity in human pancreatic cancer cells, we identified 93 genes that, when



Figure 4. Kaplan Meier log rank survival analysis for CHD7 expression in patients receiving adjuvant therapy (n = 59) and in patients receiving gemcitabine therapy (n = 42). A, effect of CHD7 expression on RFS in patients receiving gemcitabine therapy (n = 42). B, effect of CHD7 expression on OS in patients receiving gemcitabine therapy (n = 59). D, effect of CHD7 expression on OS in patients receiving adjuvant therapy (n = 59). D, effect of CHD7 expression on OS in patients receiving adjuvant therapy (n = 59). D, effect of CHD7 expression on OS in patients receiving adjuvant therapy (n = 59). D, effect of CHD7 expression on OS in patients receiving adjuvant therapy (n = 59). D, effect of CHD7 expression on OS in patients receiving adjuvant therapy (n = 59).

silenced, demonstrate gemcitabine sensitization, including CHD7. CHD7 deficiency caused gemcitabine sensitization in pancreatic adenocarcinoma cells and delayed pancreatic tumor xenograft growth in mice treated with gemcitabine. We further found that CHD7 knockdown impaired ATR dependent phos phorylation of CHK1 and increased DNA damage induced by gemcitabine, revealing a novel function for CHD7 as a DDR protein, which maintains genome integrity in response to gemcitabine. We examined CHD7 as a potential biomarker based on its dysregulation and differential expression in a panel of pancreatic adenocarcinoma cell lines and tissues. Finally, we found that low CHD7 expression is associated with improved RFS and OS in patients with early stage resected pancreatic adenocarcinoma treated with adjuvant gemcitabine. These findings support our rationale driven approach in exploiting dysregulated DDR pathways in pancreatic adenocarcinoma to identify genetic determinants of gemcitabine sensitivity that can be translated to novel biomarkers or drug targets.

A third of the genes identified in our primary gemcitabine sensitivity screen are linked to the DDR, including ATR sig naling pathway genes ATR, CHK1, RAD9, RAD1, HUS1, and CDK9 (37) and nucleotide metabolism genes RRM1 and RRM2. CHD7 was previously identified as a putative ATM/ATR sub strate (17). Our finding that CHD7 silencing in human pan creatic cancer cells potentiates gemcitabine induced DNA damage and impairs CHK1 Ser317 phosphorylation in response to gemcitabine treatment suggests that CHD7 also functions in the ATR signaling pathway and helps to explain at least in part why CHD7 knockdown causes gemcitabine sen sitization in cells and in vivo. Still, the cell cycle effects of CHD7 expression require further understanding through future stud ies, which remain ongoing. For example, CHK1 inhibition has been shown to potentiate gemcitabine induced cytotoxicity by inducing premature mitosis (38). A number of genes identified in our screen, including RRM1, RRM2, and CHK1 have previ ously been shown to determine gemcitabine sensitivity in

		RFS		OS			
Outcome	HR	95% CI	Р	HR	95% CI	Р	
Tumor size	1.708	1.223-2.387	0.002	1.685	1.171-2.425	0.005	
Positive margins	0.252	0.083-0.767	0.015	0.187	0.059-0.590	0.004	
Higher grade	1.126	0.566-2.242	0.735	1.064	0.555-2.040	0.851	
Positive nodes	1.177	0.5402-0.564	0.682	0.965	0.420-2.216	0.933	
PNI	0.574	0.210-1.567	0.279	4.133	1.044–16.358	0.043	
LVI	1.004	0.462-2.180	0.992	1.429	0.619-3.298	0.402	
Low CHD7 expression	0.122	0.035-0.420	0.001	0.086	0.025-0.292	<0.000	

NOTE: Bold denotes statistical significance.

^aMultivariate analysis includes all clinically relevant covariates with *P* < 0.2 on univariate analysis.

human pancreatic cancer cells (39), and low RRM2 expression has been shown to be associated with improved outcome in patients with pancreatic adenocarcinoma (24) and specifically those treated with adjuvant gemcitabine (28), providing val idation for our screen in identifying gemcitabine sensitivity genes that may function as potential biomarkers. Several of the gemcitabine sensitivity genes, including PLK1 and AURKB, are involved in mitotic progression that is in part targeted by nanoparticle albumin bound (nab) paclitaxel (Abraxane; Cel gene), which potentiates gemcitabine sensitivity and improves survival in patients with metastatic pancreatic adenocarcino ma treated with gemcitabine (40, 41). It is thus possible that the gemcitabine sensitivity genes reported in this study may also be novel druggable targets to be used in combination with gemcitabine. Indeed, PARP2, a target of PARP inhibitors that sensitizes pancreatic cancer cells to gemcitabine (42, 43), was also identified in our screen.

In our clinical data, low CHD7 expression was associated with increased OS and RFS in all patients receiving adjuvant therapy, although this was likely driven by the inclusion of patients receiving gemcitabine. The association of low CHD7 expression with increased survival was magnified in patients receiving gemcitabine as a component of their adjuvant ther apy despite smaller patient numbers, indicating that low CHD7 expression may indeed be associated with gemcitabine sensi tivity in these patients. In contrast, CHD7 expression in patients not receiving gemcitabine was not statistically signif icant. This analysis is underpowered with limitation of small sample size and selection bias, but our findings provide valuable hypothesis generating data suggesting that CHD7 may have predictive value in these patients.

Given the evidence that patients with low CHD7 expression demonstrate improved outcomes, it is possible that adjuvant therapy regimens could be tailored to individualize patient treatment based on CHD7 expression. This should be exam ined in future prospective trials and in larger secondary analyses of completed prospective studies. Although adjuvant chemotherapy for patients with pancreatic adenocarcinoma is advantageous, the ideal drug regimen remains unclear. The benefit of adjuvant gemcitabine compared with adjuvant 5 FU in patients with early stage resected pancreatic adenocarci noma has not been demonstrated in any large trials. Both the ESPAC 3 trial, which randomized patients with resected pan creatic adenocarcinoma to adjuvant gemcitabine versus 5 FU, and the RTOG 97 04 trial, which randomized patients with resected pancreatic adenocarcinoma to adjuvant pre and postchemoradiotherapy gemcitabine versus 5 FU, reported no significant difference in disease free survival (DFS) or OS

Outcome		RFS		OS			
	HR	95% CI	Р	HR	95% CI	Р	
Tumor size	1.622	1.228-2.144	0.001	1.527	1.130-2.063	0.006	
Positive margins	0.849	0.408-1.768	0.662	0.724	0.347-1.511	0.389	
Higher grade	1.051	0.579-1.907	0.870	1.057	0.592-1.887	0.851	
Positive nodes	1.638	0.777-3.452	0.194	1.729	0.818-3.652	0.151	
PNI	0.410	0.161-1.044	0.062	2.028	0.591-6.960	0.261	
LVI	1.064	0.529-2.139	0.862	1.082	0.537-2.178	0.826	
Low CHD7 expression	0.271	0.107-0.687	0.006	0.203	0.085-0.486	<0.000	

NOTE: Bold denotes statistical significance.

^aMultivariate analysis includes all clinically relevant covariates with P < 0.2 on univariate analysis.

between the two arms (3, 44). Our finding that low CHD7 expression is associated with improved outcome in patients with early stage pancreatic adenocarcinoma treated with adju vant gemcitabine suggest that, once validated, CHD7 expres sion could potentially be used as a predictive biomarker to individualize adjuvant therapy for these patients. In addition, the optimal radiation dose and fractionation for patients with resected pancreatic adenocarcinoma remains unknown, and molecular biomarkers to guide adjuvant therapy decisions are essential (45). The potential utility of CHD7 expression as a prognostic and potentially predictive biomarker still remains a hypothesis generating observation and requires validation in a prospective clinical trial, in which regimen dosing and duration are more homogenous.

Interest in genetic sequencing data, such as with The Cancer Genome Atlas and other similar projects, continues to increase (46, 47), leading to rapidly increasing knowledge of genes expressed and mutated in specific cancer types including pancreatic adenocarcinoma. As this knowledge becomes available, it is crucial that an approach be devel oped to help identify those genes that may serve as clinically relevant prognostic or predictive biomarkers or potential drug targets for novel therapeutics. The successful identification and validation of *CHD7* as a novel gemcitabine sensitivity gene that is associated with outcome in patients with early stage pancreatic adenocarcinoma treated with adjuvant gemcitabine is evidence that our approach may be successful in identifying other clinically relevant biomarkers or drug targets.

It is worth noting that recent advances in chemotherapy have increased the use of FOLFIRINOX therapy in the meta static setting, impacting the potential utility of this study. Still, National Comprehensive Cancer Network guidelines in the metastatic setting equivalently recommend FOLFIRINOX or two gemcitabine based regimens (gemcitabine with the addi tion of erlotinib or nab paclitaxel), both with category one evidence (24). In addition, gemcitabine or fluoropyrimidine therapies are still recommended in the adjuvant setting, which is where this study's clinical focus remains. Future studies should evaluate the predictive role of CHD7 in a larger, randomized prospective trial to validate potential gemcitabine sensitivity genes in a similar fashion to other identified pre

- Heinemann V, Haas M, Boeck S. Systemic treatment of advanced pancreatic cancer. Cancer Treat Rev 2012;38:843 53.
- Neoptolemos JP, Stocken DD, Bassi C, Ghaneh P, Cunningham D, Goldstein D, et al. Adjuvant chemotherapy with fluorouracil plus folinic acid vs gemcitabine following pancreatic cancer resection: a random ized controlled trial. JAMA 2010;304:1073 81.
- Karhu R, Mahlamaki E, Kallioniemi A. Pancreatic adenocarcinoma genetic portrait from chromosomes to microarrays. Genes Chromo somes Cancer 2006;45:721 30.
- Oettle H, Post S, Neuhaus P, Gellert K, Langrehr J, Ridwelski K, et al. Adjuvant chemotherapy with gemcitabine vs observation in patients undergoing curative intent resection of pancreatic cancer: a random ized controlled trial. JAMA 2007;297:267 77.

dictive biomarkers (48). The current study suggests that CHD7 may be a useful biomarker for determining which patients will derive greater benefit from gemcitabine therapy, providing clinicians a way to better select patients for specific adjuvant therapy regimens in the future.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Disclaimer

The content of this article is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.

Authors' Contributions

Conception and design: L.E. Colbert, S.B. Fisher, C.W. Hardy, G.P. Nagaraju, W.A. Hall, J.C. Landry, S.K. Maithel, D.S. Yu

Development of methodology: L.E. Colbert, S.B. Fisher, M.Z. Madden, C.W. Hardy, G.P. Nagaraju, W.A. Hall, J.C. Landry, S.K. Maithel, D.S. Yu

Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.): L.E. Colbert, A.V. Petrova, S.B. Fisher, B.G. Pantazides, M.Z. Madden, C.W. Hardy, M.D. Warren, Y. Pan, G.P. Nagaraju, E.A. Liu, B. Saka, W.A. Hall, C.A. Staley III, W.J. Curran Jr, J.C. Landry

Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis): L.E. Colbert, A.V. Petrova, S.B. Fisher, C.W. Hardy, G.P. Nagaraju, W.A. Hall, J.W. Shelton, K. Gandhi, R. Pauly, J. Kowalski, N.V. Adsay, S.K. Maithel, D.S. Yu

Writing, review, and/or revision of the manuscript: L.E. Colbert, A.V. Petrova, S.B. Fisher, C.W. Hardy, W.A. Hall, J.W. Shelton, J. Kowalski, D.A. Kooby, B.F. El-Rayes, C.A. Staley III, W.J. Curran Jr, J.C. Landry, S.K. Maithel, D.S. Yu Administrative, technical, or material support (i.e., reporting or orga-

nizing data, constructing databases): L.E. Colbert, C.W. Hardy, Y. Pan, G.P. Nagaraju, W.A. Hall, R. Pauly, D.S. Yu

Study supervision: W.A. Hall, W.J. Curran Jr, J.C. Landry, S.K. Maithel, D.S. Yu

Acknowledgments

The authors thank members of the Yu laboratory and Chris Crane for helpful comments during article preparation.

Grant Support

This work was supported by the National Center for Advancing Translational Sciences of the NIH (ULI TR000454 to L.E. Colbert and S.B. Fisher, and TLITR000456 to L.E. Colbert); Pancreatic Cancer Action Network/American Association for Cancer Research (16982; D.S. Yu); Department of Defense/Peer Reviewed Cancer Research Program (CA110535; D.S. Yu); and Georgia Cancer Coalition (11072; D.S. Yu).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received July 15, 2013; revised February 11, 2014; accepted February 13, 2014; published OnlineFirst March 13, 2014.

- Bartkova J, Horejsi Z, Koed K, Kramer A, Tort F, Zieger K, et al. DNA damage response as a candidate anti cancer barrier in early human tumorigenesis. Nature 2005;434:864 70.
- Gorgoulis VG, Vassiliou LV, Karakaidos P, Zacharatos P, Kotsinas A, Liloglou T, et al. Activation of the DNA damage checkpoint and genomic instability in human precancerous lesions. Nature 2005;434:907 13.
- Forbes S, Clements J, Dawson E, Bamford S, Webb T, Dogan A, et al. Cosmic 2005. Br J Cancer 2006;94:318 22.
- Vissers LE, van Ravenswaaij CM, Admiraal R, Hurst JA, de Vries BB, Janssen IM, et al. Mutations in a new member of the chromodomain gene family cause CHARGE syndrome. Nat Genet 2004;36:955 7.
- Kim HG, Kurth I, Lan F, Meliciani I, Wenzel W, Eom SH, et al. Mutations in CHD7, encoding a chromatin remodeling protein, cause idiopathic hypogonadotropic hypogonadism and Kallmann syndrome. Am J Hum Genet 2008;83:511 9.

References

Siegel R, Naishadham D, Jemal A. Cancer statistics, 2012. CA Cancer J Clin 2012;62:10 29.

- Bamford S, Dawson E, Forbes S, Clements J, Pettett R, Dogan A, et al. The COSMIC (Catalogue of Somatic Mutations in Cancer) database and website. Br J Cancer 2004;91:355 8.
- 12. Gao J, Aksoy BA, Dogrusoz U, Dresdner G, Gross B, Sumer SO, et al. Integrative analysis of complex cancer genomics and clinical profiles using the cBioPortal. Sci Signal 2013;6:pl1.
- Cerami E, Gao J, Dogrusoz U, Gross BE, Sumer SO, Aksoy BA, et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. Cancer Discov 2012;2: 401 4.
- Bajpai R, Chen DA, Rada Iglesias A, Zhang J, Xiong Y, Helms J, et al. CHD7 cooperates with PBAF to control multipotent neural crest formation. Nature 2010;463:958 62.
- Zentner GE, Hurd EA, Schnetz MP, Handoko L, Wang C, Wang Z, et al. CHD7 functions in the nucleolus as a positive regulator of ribosomal RNA biogenesis. Hum Mol Genet 2010;19:3491 501.
- Engelen E, Akinci U, Bryne JC, Hou J, Gontan C, Moen M, et al. Sox2 cooperates with Chd7 to regulate genes that are mutated in human syndromes. Nat Genet 2011;43:607 11.
- Matsuoka S, Ballif BA, Smogorzewska A, McDonald ER III, Hurov KE, Luo J, et al. ATM and ATR substrate analysis reveals extensive protein networks responsive to DNA damage. Science 2007;316:1160 6.
- Batsukh T, Schulz Y, Wolf S, Rabe TI, Oellerich T, Urlaub H, et al. Identification and characterization of FAM124B as a novel component of a CHD7 and CHD8 containing complex. PLoS ONE 2012;7:e52640.
- Nagaraju GP, Zhu S, Wen J, Farris AB, Adsay VN, Diaz R, et al. Novel synthetic curcumin analogues EF31 and UBS109 are potent DNA hypomethylating agents in pancreatic cancer. Cancer Lett 2013;341: 195 203.
- Harsha HC, Kandasamy K, Ranganathan P, Rani S, Ramabadran S, Gollapudi S, et al. A compendium of potential biomarkers of pancreatic cancer. PLoS Med 2009;6:e1000046.
- Maupin KA, Sinha A, Eugster E, Miller J, Ross J, Paulino V, et al. Glycogene expression alterations associated with pancreatic cancer epithelial mesenchymal transition in complementary model systems. PLoS ONE 2010;5:e13002.
- Pei H, Li L, Fridley BL, Jenkins GD, Kalari KR, Lingle W, et al. FKBP51 affects cancer cell response to chemotherapy by negatively regulating Akt. Cancer Cell 2009;16:259 66.
- 23. Maithel SK, Coban I, Kneuertz PJ, Kooby DA, El Rayes BF, Kauh JS, et al. Differential expression of ERCC1 in pancreas adenocarcinoma: high tumor expression is associated with earlier recurrence and short ened survival after resection. Ann Surg Oncol 2011;18:2699 705.
- 24. Fisher SB, Patel SH, Bagci P, Kooby DA, El Rayes BF, Staley CA III, et al. An analysis of human equilibrative nucleoside transporter 1, ribonucleoside reductase subunit M1, ribonucleoside reductase sub unit M2, and excision repair cross complementing gene 1 expression in patients with resected pancreas adenocarcinoma: implications for adjuvant treatment. Cancer 2013;119:445 53.
- 25. Colbert LE, Fisher SB, Hardy CW, Hall WA, Saka B, Shelton JW, et al. Pronecrotic mixed lineage kinase domain like protein expression is a prognostic biomarker in patients with early stage resected pancreatic adenocarcinoma. Cancer 2013;119:3148 55.
- 26. Hall WA, Petrova AV, Colbert LE, Hardy CW, Fisher SB, Saka B, et al. Low CHD5 expression activates the DNA damage response and predicts poor outcome in patients undergoing adjuvant therapy for resected pancreatic cancer. Oncogene 2013 Nov 25. [Epub ahead of print].
- 27. Basturk O, Singh R, Kaygusuz E, Balci S, Dursun N, Culhaci N, et al. GLUT 1 expression in pancreatic neoplasia: implications in pathogen esis, diagnosis, and prognosis. Pancreas 2011;40:187 92.
- 28. Fujita H, Ohuchida K, Mizumoto K, Itaba S, Ito T, Nakata K, et al. Gene expression levels as predictive markers of outcome in pancreatic cancer after gemcitabine based adjuvant chemotherapy. Neoplasia 2010;12:807 17.
- 29. Cotta Ramusino C, McDonald ER III, Hurov K, Sowa ME, Harper JW, Elledge SJ. A DNA damage response screen identifies RHINO, a 9 1 1 and TopBP1 interacting protein required for ATR signaling. Science 2011;332:1313 7.

- 30. Paulsen RD, Soni DV, Wollman R, Hahn AT, Yee MC, Guan A, et al. A genome wide siRNA screen reveals diverse cellular processes and pathways that mediate genome stability. Mol Cell 2009;35:228 39.
- Agata N, Ahmad R, Kawano T, Raina D, Kharbanda S, Kufe D. MUC1 oncoprotein blocks death receptor mediated apoptosis by inhibiting recruitment of caspase 8. Cancer Res 2008;68:6136 44.
- 32. Akada M, Crnogorac Jurcevic T, Lattimore S, Mahon P, Lopes R, Sunamura M, et al. Intrinsic chemoresistance to gemcitabine is asso ciated with decreased expression of BNIP3 in pancreatic cancer. Clin Cancer Res 2005;11:3094 101.
- 33. Azorsa DO, Gonzales IM, Basu GD, Choudhary A, Arora S, Bisanz KM, et al. Synthetic lethal RNAi screening identifies sensitizing targets for gemcitabine therapy in pancreatic cancer. J Transl Med 2009;7:43.
- Dong N, Wang M, Li H, Cui Y, Guo Q. Gemcitabine in combination with vinorelbine in elderly patients with anthracycline and taxane pre treated metastatic breast cancer. Cancer Chemother Pharmacol 2012;69:1315 22.
- 35. O'Connell BC, Adamson B, Lydeard JR, Sowa ME, Ciccia A, Brede meyer AL, et al. A genome wide camptothecin sensitivity screen identifies a mammalian MMS22L NFKBIL2 complex required for geno mic stability. Mol Cell 2010;40:645 57.
- Whitehurst AW, Bodemann BO, Cardenas J, Ferguson D, Girard L, Peyton M, et al. Synthetic lethal screen identification of chemosensi tizer loci in cancer cells. Nature 2007;446:815 9.
- Yu DS, Zhao R, Hsu EL, Cayer J, Ye F, Guo Y, et al. Cyclin dependent kinase 9 cyclin K functions in the replication stress response. EMBO Rep 2010;11:876 82.
- Morgan MA, Parsels LA, Parsels JD, Lawrence TS, Maybaum J. The relationship of premature mitosis to cytotoxicity in response to check point abrogation and antimetabolite treatment. Cell Cycle 2006;5: 1983 8.
- 39. Zhou J, Chen Z, Malysa A, Li X, Oliveira P, Zhang Y, et al. A kinome screen identifies checkpoint kinase 1 (CHK1) as a sensitizer for RRM1 dependent gemcitabine efficacy. PLoS ONE 2013;8:e58091.
- 40. Von Hoff D, Ervin TJ, Arena FP, Chiorean EG, Infante JR, Moore MJ, et al., editors. Randomized phase III study of weekly nab paclitaxel plus gemcitabine versus gemcitabine alone in patients with metastatic adenocarcinoma of the pancreas (MPACT). J Clin Oncol 30: 2012 (suppl 34; abstr LBA148).
- Frese KK, Neesse A, Cook N, Bapiro TE, Lolkema MP, Jodrell DI, et al. nab Paclitaxel potentiates gemcitabine activity by reducing cytidine deaminase levels in a mouse model of pancreatic cancer. Cancer Discov 2012;2:260 9.
- 42. Jacob DA, Bahra M, Langrehr JM, Boas Knoop S, Stefaniak R, Davis J, et al. Combination therapy of poly (ADP ribose) polymerase inhibitor 3 aminobenzamide and gemcitabine shows strong antitumor activity in pancreatic cancer cells. J Gastroenterol Hepatol 2007;22:738 48.
- 43. Porcelli L, Quatrale AE, Mantuano P, Leo MG, Silvestris N, Rolland JF, et al. Optimize radiochemotherapy in pancreatic cancer: PARP inhi bitors a new therapeutic opportunity. Mol Oncol 2013;7:308 22.
- 44. Regine WF, Winter KA, Abrams RA, Safran H, Hoffman JP, Konski A, et al. Fluorouracil vs gemcitabine chemotherapy before and after fluorouracil based chemoradiation following resection of pancreatic adenocarcinoma: a randomized controlled trial. JAMA 2008;299: 1019 26.
- 45. Hall WA, Colbert LE, Liu Y, Gillespie T, Lipscomb J, Hardy C, et al. The influence of adjuvant radiotherapy dose on overall survival in patients with resected pancreatic adenocarcinoma. Cancer 2013;119:2350 7.
- 46. Biankin AV, Waddell N, Kassahn KS, Gingras MC, Muthuswamy LB, Johns AL, et al. Pancreatic cancer genomes reveal aberrations in axon guidance pathway genes. Nature. 2012;491:399 405.
- 47. Iacobuzio Donahue CA, Velculescu VE, Wolfgang CL, Hruban RH. Genetic basis of pancreas cancer development and progression: insights from whole exome and whole genome sequencing. Clin Can cer Res 2012;18:4257 65.
- Farrell JJ, Elsaleh H, Garcia M, Lai R, Ammar A, Regine WF, et al. Human equilibrative nucleoside transporter 1 levels predict response to gemcitabine in patients with pancreatic cancer. Gastroenterology 2009;136:187 95.

www.nature.com/onc

ORIGINAL ARTICLE

Low CHD5 expression activates the DNA damage response and predicts poor outcome in patients undergoing adjuvant therapy for resected pancreatic cancer

WA Hall¹, AV Petrova¹, LE Colbert¹, CW Hardy¹, SB Fisher², B Saka³, JW Shelton¹, MD Warren¹, BG Pantazides¹, K Gandhi⁴, J Kowalski⁴, DA Kooby^{2,5}, BF El-Rayes⁶, CA Staley 3rd², N Volkan Adsay³, WJ Curran¹, JC Landry¹, SK Maithel² and DS Yu¹

The DNA damage response (DDR) promotes genome integrity and serves as a cancer barrier in precancerous lesions but paradoxically may promote cancer survival. Genes that activate the DDR when dysregulated could function as useful biomarkers for outcome in cancer patients. Using a siRNA screen in human pancreatic cancer cells, we identified the *CHD5* tumor suppressor as a gene, which, when silenced, activates the DDR. We evaluated the relationship of CHD5 expression with DDR activation in human pancreatic cancer cells and the association of CHD5 expression in 80 patients with resected pancreatic adenocarcinoma (PAC) by immunohistochemical analysis with clinical outcome. CHD5 depletion and low CHD5 expression in human pancreatic cancer cells lead to increased H2AX-Ser139 and CHK2-Thr68 phosphorylation and accumulation into nuclear foci. On Kaplan–Meier log-rank survival analysis, patients with low CHD5 expression had a median recurrence-free survival (RFS) of 5.3 vs 15.4 months for patients with high CHD5 expression (P = 0.03). In 59 patients receiving adjuvant chemotherapy, low CHD5 expression was associated with decreased RFS (4.5 vs 16.3 months; P = 0.001) and overall survival (OS) (7.2 vs 21.6 months; P = 0.003). On multivariate Cox regression analysis, low CHD5 expression remained associated with worse OS (HR: 3.187 (95% CI: 1.49–6.81); P = 0.003) in patients undergoing adjuvant chemotherapy. Thus, low CHD5 expression activates the DDR and predicts for worse OS in patients with resected PAC receiving adjuvant chemotherapy. Our findings support a model in which dysregulated expression of tumor suppressor genes that induce DDR activation can be utilized as biomarkers for poor outcome.

Oncogene (2014) 33, 5450-5456; doi:10.1038/onc.2013.488; published online 25 November 2013

Keywords: CHD5; DNA damage response; biomarker; pancreatic cancer; tumor suppressor

INTRODUCTION

Pancreatic adenocarcinoma (PAC) is a devastating malignancy, and 5-year overall survival (OS) rates for all patients remain less than 10%.¹ Despite large, prospective randomized trials, the optimal management of PAC following surgical resection is still unknown. Several trials have resulted in equivocal conclusions regarding the optimal use of various combinations of adjuvant chemotherapy and radiation therapy or chemotherapy alone.^{2–4} Currently the data supporting prognostic biomarkers in PAC continue to evolve, which make optimal treatment decisions difficult. Given the increasingly apparent genetic heterogeneity of PAC, it seems conceivable that genetic profiles exist to provide prognostic information, which could help to predict response to adjuvant therapy.^{5–8}

The DNA damage response (DDR) is critical for the maintenance of genome integrity and serves as a cancer barrier by mobilizing DNA repair, cell cycle arrest or apoptosis.^{9,10} In human precancerous lesions, replication stress resulting from activated oncogenes and inactivated tumor suppressor proteins induces activation of the DDR, which constrains tumor development. This creates a selection pressure for mutations or epigenetic silencing of DDR genes leading to the proliferation of genetically unstable cells and ultimately resulting in cancer. Thus, the DDR acts as a barrier against genomic instability and cancer development. Indeed, a large number of DDR genes are somatically mutated in PAC, including *ATM*, *BRCA2*, *CDKN2A*, *FANCI*, *HELB* and *RAD9*.¹¹ Although the DDR promotes genome integrity and helps to prevent progression of precancerous lesions, it may paradoxically promote the survival of PAC that outgrow the selection pressure of DDR activation. These cancers could potentially be more resistant to the replication stress and DNA damage induced by chemotherapy and radiation therapy that is used in the adjuvant setting for early-stage PAC. The clinical significance of dysregulated expression of genes that induce DDR activation in PAC and other cancers is currently not known.

Chromodomain helicase DNA-binding protein 5 (CHD5) is a member of a family of chromodomain enzymes that belong to the ATP-dependent chromatin remodeling protein SNF2 superfamily and characterized as a tumor suppressor. CHD5 has been shown to be mutated, deleted or silenced in a number of malignancies including neuroblastoma, gastric cancer, leukemia/lymphoma, melanoma, breast, prostate, ovarian and lung cancers, squamous cell of the larynx and gliomas.^{12–24} The most thorough characterization of CHD5 is in neural tissues where it controls proliferation, apoptosis and senescence via the p19(Arf)/p53

E mail: dsyu@emory.edu

Received 6 May 2013; revised 17 August 2013; accepted 4 October 2013; published online 25 November 2013

¹Department of Radiation Oncology, Winship Cancer Institute, Emory University, Atlanta, GA, USA; ²Division of Surgical Oncology, Department of Surgery, Winship Cancer Institute, Emory University, Atlanta, GA, USA; ³Department of Pathology, Winship Cancer Institute, Emory University, Atlanta, GA, USA; ⁴Biostatistics and Bioinformatics Shared Resource, Winship Cancer Institute, Atlanta, GA, USA; ⁵Atlanta Veterans Affairs Medical Center, Atlanta, GA, USA and ⁶Department of Medical Oncology, Winship Cancer Institute, Emory University, Atlanta, GA, USA. Correspondence: Dr DS Yu, Department of Radiation Oncology, Emory University School of Medicine, 1365 Clifton Rd NE, C3008, Atlanta, GA 30322, USA.

Low CHD5 expression activates DDR predicts outcome WA Hall et al

Nucleus

5451

pathway.¹² CHD5 binds to histone H3 for its tumor suppressor function.²⁵ CHD5 is also a putative substrate of the ATM/ATR checkpoint kinases, suggesting that it may have a role in the DDR.²⁶ Furthermore, low CHD5 expression has been shown to correlate with poor patient outcomes in a number of other malignancies, independently of other known prognostic factors.^{21,24,27–29} The role of CHD5 expression in PAC has not previously been explored.

RESULTS

CHD5 silencing activates the DDR

a

siRNA Gene

F

CHD5-1

To identify genes, which, when dysregulated, induce DDR activation, we completed a loss-of-function genetic screen in

YH2AX

Nucleus

MIA PaCa-2 human pancreatic cancer cells for genes, which when silenced increase DDR signaling. Our library included 200 siRNAs targeting 50 nuclear enzyme genes. MIA PaCa-2 cells were transfected with siRNA and analyzed 72 h later by indirect immunofluorescence microscopy for phosphorylation of H2AX Ser139 (γ H2AX), an ATM/ATR substrate and early marker of DDR activation. Two siRNAs targeting a different region of CHD5 significantly induced γ H2AX foci formation above a non-targeting control (Figures 1a and b). Similarly two siRNAs targeting CHD5 significantly induced formation of phospho-CHK2 Thr68 foci (Figures 1c and d), an ATM/ATR substrate and second marker of DDR activation, compared with a non-targeting control. Western blot analysis confirmed knockdown of CHD5 expression, specificity

Merge



C

siRNA Gene

CHD5-1

ŧ

Merge

pCHK2 Thr68

5452

of the anti-CHD5 antibody, as well as increased phosphorylation of CHK2 Thr68 and γ H2AX (Figure 1e). Consistent with these findings, MIA PaCa-2 and BxPC3 pancreatic cancer cells, which express lower levels of CHD5 than PANC1 and HPAC pancreatic cancer cells, demonstrated increased phosphorylation of CHK2 Thr68 and γ H2AX (Figure 1f). Collectively, these findings suggest that CHD5 silencing and low CHD5 expression in cells activates the DDR.

To determine whether CHD5 expression might function as an ideal biomarker, we mined published genome-wide data sets to determine if its expression is differentially expressed among pancreatic cancer cell lines and PAC patient samples. CHD5 was found to be above the 90th percentile of genes for differential expression among a panel of 22 PAC cell lines and 20 PAC patient samples.^{30,31}

Clinical results

Patient characteristics can be seen in Table 1. Median patient age was 63.1 years (37–84), and median follow-up for survivors was 53 months (range 6–114). The median OS for all patients was 15.4 months (range 2.8–114.6), and median RFS was 9.3 months (range 0.6–119.8). At last known follow-up, 14 (17.5%) patients had no evidence of disease, and 62 (77.5%) of patients were deceased. A total of 59 (73.8%) patients received adjuvant therapy, with most of these patients receiving gemcitabine-based regimens (n = 41/59; 69.5%) and radiation therapy (39/59; 66.1%). Forty-two patients (52.5%) were male, and 59 (73.8%) were white. Tumor size ranged from 1 cm to 6 cm with a median of 3.3 cm; 20 patients (24.5%) had positive margins and 48 (60%) had positive lymph nodes.

Survival analyses: all patients

Eighty patients were analyzed in this group. Kaplan–Meier analyses showed an association between low CHD5 expression and decreased OS in all patients (8.5 months vs 18.1 months; P = 0.12; Figure 2a) and recurrence-free survival (RFS) (5.3 months vs 15.4 months; P = 0.03; Figure 2a). The complete results of the univariate and multivariate Cox regression analyses for all patients can be seen in Table 2. Low CHD5 expression was associated with decreased OS on multivariate analysis (HR: 2.3 (95% CI: 1.2–4.5); P = 0.01). The only other significant prognostic factor on multivariate analysis was lymphovascular invasion (HR: 1.4 (95% CI: 0.6–3.4); P = 0.04).

Subset analyses: patients receiving adjuvant therapy

In the subset of patients receiving adjuvant therapy, 59 patients were analyzed. Kaplan–Meier analyses showed an association between low CHD5 expression and decreased OS in patients undergoing adjuvant therapy (7.2 vs 21.6 months; P = 0.003; Figure 2b) and RFS (4.5 vs 16.3 months; P = 0.001; Figure 2b). This relationship persisted on multivariate analysis, examining the effect of CHD5 expression on OS (HR: 3.19 (95% CI: 1.5–6.81); P = 0.003). The full multivariate results for patients receiving adjuvant therapy can be seen in Table 3.

As the initial exploratory intent of these genetic markers was to provide prognostic data for patients undergoing adjuvant gemcitabine-based therapy, a second subset analysis was performed in the patients receiving only gemcitabine-based therapy, and the previously defined relationship remained significant on multivariate analysis, with low CHD5 expression associated with decreased RFS (HR: 3.08 (95% CI: 1.09–8.73); P = 0.035; Supplementary Table S1) and decreased OS (HR: 3.041 (95% CI: 1.13–8.16); P = 0.027; Supplementary Table S1). Finally an exploratory multivariate analysis was performed comparing the prognostic significance of CHD5 with other known prognostic genetic markers, including excision repair cross-complementing gene 1 (ERCC1) and ribonucleotide reductase subunit 2 (RRM2) that have been previously published and shown to be significant Table 1. Patient demographics, tumor characteristics and treatmentcharacteristics for all patients (N80)

		Ν	Percen
Patient Demographics	The second second		
Age (years)	63.1 (37 84) ^a		
Male sex		42	52.5
Ethnicity			
Asian		2	2.5
Black		15	18.8
White		59	73.8
Overall survival (months)	15.4 (2.8 114.6) ^a		
Recurrence free survival	9.3 (0.6 119.8) ^a		
(months)			
Pathological characteristics			
Positive margins		20	24.5
Grade			
1		5	6.3
2		46	57.5
3		28	35.0
Positive nodes		48	60.0
Perineural invasion		70	87.5
Lymphovascular invasion		38	47.5
High CHD5 expression		63	78.75
Tumor size (cm)	3.3 (1 6) ^a		
Treatment characteristics			
Neoadjuvant therapy		2	2.5
Radiation therapy		39	48.8
Received gemcitabine		41	51.3

using this patient cohort.⁸ The results of this multivariate analysis can be seen in Table 4; of note, CHD5 remained significant independent of other known prognostic genetic markers in this population, ERCC1 and RRM2.^{7,8}

DISCUSSION

Our results demonstrate that silencing of the CHD5 tumor suppressor and low CHD5 expression in human pancreatic cancer cells lead to activation of the DDR and that low CHD5 expression is associated with worse RF5 and OS in patients with PAC treated by resection and adjuvant chemotherapy. These findings support a model in which dysregulated expression of tumor suppressor genes that induce DDR activation can function as biomarkers for poor outcome.

The role and function of CHD5 have been explored in several malignancies, and CHD5 has been well characterized as a tumor suppressor protein.^{12,21,23,29} Multiple studies have demonstrated that silencing or downregulation of CHD5 is associated with the development of various malignancies including lung,²² gastric,¹⁵ neuroblastoma,23 breast,21 and laryngeal squamous cell carcinoma.³² We have presented unique data that demonstrate the prognostic ability of CHD5 for patients receiving adjuvant chemotherapy following resection for PAC. In addition to the clinically significant association of CHD5 expression, we have presented a mechanistic explanation as to the etiology of these clinical findings. Our preclinical data demonstrate that low CHD5 expression in human pancreatic cancer cells leads to activation of the DDR with increased yH2AX and phosho-CHK2 Thr68 foci accumulation and expression. We hypothesize that the increased activation of the DDR in cells with low CHD5 expression could contribute to the worse clinical outcomes seen in this

Low CHD5 expression activates DDR predicts outcome WA Hall et al



Figure 2. Kaplan Meier log-rank survival analysis examining CHD5 expression associated with OS and RFS in (a) all patients and (b) patients undergoing adjuvant therapy only.

Outcome	2	RFS			OS			
	Univariate		Univariate Multivariate ^a		Univariate		Multivariate ^a	
	HR (95% CI)	P value	HR (95% CI)	P value	HR (95% CI)	P value	HR (95% CI)	P value
Tumor size	1.3 (1.0 1.6)	0.02	1.3 (1.0 1.3)	0.02	1.3 (1.0 1.6)	0.02	1.2 (1.0 1.6)	0.1
Positive margins	1.8 (1.1 3.0)	0.02	1.4 (0.8 2.5)	0.2	1.7 (1.1 2.8)	0.03	1.3 (0.7 2.2)	0.4
Positive nodes	1.8 (1.0 3.0)	0.04	1.5 (0.8 2.9)	0.2	2.0 (1.1 3.4)	0.01	1.5 (0.8 2.9)	0.2
PNI	1.0 (0.5 2.1)	0.9	0.9 (0.4 2.0	0.8	3.9 (1.2 12.4)	0.02	3.8 (1.2 12.7)	0.03
LVI	1.5 (.9 2.5)	0.1	1.4 (0.8 2.5)	0.3	2.0 (1.2 3.3)	0.01	1.4 (0.6 3.4)	0.04
Higher grade	1.3 (0.9 2.0)	0.2	1.3 (0.8 2.0)	0.3	1.3 (0.8 1.9)	0.3	1.7 (0.9 3.0)	0.08
Low CHD5 expression ^b	2.0 (1.0 3.3)	0.03	2.5 (1.3 4.6)	0.006	1.6 (0.8 2.9)	0.1	2.3 (1.2 4.5)	0.01

Abbreviations: LVI, lymphovascular invasion; OS, overall Survival; PNI, perineural invasion; RFS, recurrence free survival. "Multivariate analysis includes all clinically relevant covariates and covariates with *P* value < 0.1 on univariate analysis. ^bCHD5 Score is decreasing from high expression to low expression.

patient cohort. The dysregulation of CHD5 expression and activation of the DDR could promote the survival of PAC that outgrow the selection pressure of DDR activation through compensatory DNA repair or pro-survival pathways. Specifically, these cancers may be more resistant to the replication stress and DNA damage induced by chemotherapy and radiation therapy that is used in the adjuvant setting for early-stage PAC, and thus contribute to the worse clinical outcomes seen in these patients.

CHD5 expression levels have been shown to be prognostic in various other disease sites. In a series by Wong *et al.*, down-regulation of CHD5 expression was shown to be an independent adverse prognostic factor in patients with ovarian cancer. Similar

5454

Outcome		RFS		OS		
	HR	95% Cl	P value	HR	95% Cl	P value
Tumor size	1.411	1.095 1.817	0.008	1.276	0.967 1.684	0.085
Positive margins	0.999	0.488 2.043	0.998	0.996	0.471 2.109	0.993
Higher grade	1.072	0.593 1.940	0.817	1.157	0.645 2.075	0.625
Positive nodes	1.964	0.910 4.236	0.085	1.770	0.841 3.726	0.133
PNI	0.488	0.191 1.252	0.136	1.790	0.541 5.919	0.340
LVI	1.078	0.543 2.138	0.830	1.346	0.673 2.691	0.401
Low CHD5 expression ^b	3.013	1.383 6.563	0.006	3.187	1.492 6.808	0.003

Abbreviations: LVI, lymphovascular invasion; OS, overall Survival; PNI, perineural invasion; RFS, recurrence free survival. ^aMultivariate analysis includes all clinically relevant covariates and covariates with P value < 0.1 on univariate analysis. ^bCHD5 Score is decreasing from high expression to low expression.

Table 4. Multivariate^a Cox regression analyses for CHD5, ERCC1 and RRM2 in adjuvant patients (N 31)

Outcome		RFS		-	OS	
	HR	95% CI	P value	HR	95% CI	P value
Tumor size	1.253	0.930 1.687	0.138	1.214	0.889 1.658	0.223
Positive margins	1.624	0.730 3.614	0.235	1.456	0.635 3.337	0.374
Higher grade	1.654	0.858 3.187	0.133	1.363	0.750 2.479	0.310
Positive Nodes	1.573	0.750 3.297	0.231	1.360	0.642 2.880	0.422
PNI	0.575	0.219 1.508	0.260	1.944	0.587 6.431	0.276
LVI	1.256	0.639 2.471	0.508	1.377	0.694 2.733	0.361
Low CHD5 expression ^b	3.777	1,736 8,217	0.001°	2.672	1,234 5.784	0.013
High RRM2 expression	3.173	1.118 9.000	0.030	1.319	0.431 4.036	0.627
High ERCC1 expression	9.058	3.36724.369	< 0.0001	7.173	2.673 19.247	< 0.0001

Abbreviations: ERCC, excision repair cross complementing gene 1; LVI, lymphovascular invasion; OS, overall Survival; PNI, perineural invasion; RFS, recurrence free survival; RRM2, ribonucleotide reductase subunit 2. ^aMultivariate analysis includes all clinically relevant covariates with *P* value <0.2 on univariate analysis. ^bCHD5 score is decreasing from high expression to low expression. ^cBold font denotes statistical significance.

to the current study, the decreased expression of CHD5 was prognostic independent of other clinically significant variables such as age, tumor type, grade and clinical stage.²⁸ In a second series by Du et al., low CHD5 expression was prognostic for decreased RFS and OS in patients with primary gallbladder carcinoma when compared with patients with high CHD5 expression. Again, CHD5 expression was a predictor of poor outcomes independent of other known prognostic factors.³³ In a third series by Garcia et al., the expression of CHD5 was again shown to be a marker of outcome in patients with neuroblastoma independent of other relevant biological or clinical parameters. In the Garcia series, higher CHD5 expression was associated with an improved OS and event-free survival. It was also shown that after induction chemotherapy, reactivation of CHD5 expression was correlated with prolonged OS and improved clinical response. It appears consistent across a variety of malignancies that decreased CHD5 expression is associated with worse clinical outcomes.

The current series presents a cohort of 80 patients in whom CHD5 was associated with a statistically significant RFS difference but not statistically significant OS difference on KM analysis when analyzed in the cohort as a whole. This lack of significance for OS may be a consequence of the relatively small number of patients with low CHD5 expression and limitation of insufficient statistical power. However, when the analysis was repeated excluding the patients who did not receive adjuvant therapy, CHD5 became a highly significant prognostic factor for both OS and RFS, indicating that there may be a stronger correlation in this group. This finding is consistent with the hypothesis that CHD5 expression levels are correlated with cells' ability to respond to or repair DNA damage in patients who receive adjuvant therapy. CHD5 function has been explored primarily in the central nervous system (CNS). The tumor suppressor activity of CHD5 has been shown to be mediated through the p19arf/p53 pathway.¹² CHD5 was also found to be a putative ATM/ATR substrate.²⁶ Our finding that CHD5 silencing in cells activates the DDR provides support for CHD5 as a genome maintenance protein that responds to replication stress. The rationale is that, in the absence of exogenous DNA damage, the DDR will be activated as a result of spontaneous DNA damage resulting from collapsed replication forks.

Given the poor prognosis of PAC even following R0 surgical resection, genetic markers that could provide additional prognostic information for these patients are critical. Such genetic markers may help identify subsets of PAC patients at particularly high risk for poor outcomes. Improving our ability to identify such patients will help to identify those patients who may benefit the most from aggressive therapy. With further validation, given that patients with low CHD5 expression have poor outcomes, it is possible that adjuvant therapy regimens could be tailored to individualize patient treatment based on CHD5 expression. Patients with low CHD5 expression for instance may benefit from participation in clinical trials of novel therapeutic regimens, as they are less likely to benefit from standard adjuvant treatment.

The current study is the first to demonstrate the prognostic significance of CHD5 expression in patients with resected PAC. Further study of CHD5 expression in patients with resected PAC, including validation in a prospective randomized trial, is needed to better characterize the prognostic significance of this gene. In addition, CHD5 warrants further analysis of its function in the DDR given its clinical relevance as both a tumor suppressor and prognostic factor in a number of malignancies.

MATERIALS AND METHODS

Cell culture and transfections

Human MIA PaCa 2, PANC1, HPAC and BxPC3 pancreatic cancer cells were obtained from the American Type Culture Collection (ATCC, Manassas, VA, USA) and grown under standard cell culture conditions. siRNA duplexes were obtained from Dharmacon (Pittsburgh, PA, USA) and transfected with HiPerfect (Qiagen, Venlo, Limburg, Netherlands), according to the manufacturer's instructions.

Individual siRNA sequences are as follows:

non targeting (ATGAACGTGAATTGCTCAATT);

CHD5 1 (GGAAAGACCUGCCCUACGA); D 009878 01;

CHD5 2 (GAAAGAACGACAUGGAUGA); D 009878 04 renamed to CHD5 2.

Immunofluorescence

Seventy two hours after transfection, cells were washed with 1xPBS, fixed with 2% paraformaldehyde for 10 min at room temperature, permeabilized with 0.5% Triton X 100 (Fisher Scientific, Waltham, MA, USA) and blocked in a 5% bovine serum albumin solution (Sigma, St Louis, MO, USA). Cells were then immunostained with an anti phospho histone H2AX (Ser139) antibody (Millipore, Billerica, MA, USA, 05 636) and anti mouse secondary antibody with AlexaFluor 488 (Invitrogen, Carlsbad, CA, USA, A21202) or an anti phospho CHK2 (Thr68) antibody (Cell Signaling, Danvers, MA, USA, 2661) and anti rabbit secondary antibody with AlexaFluor 488 (Invitrogen, A21206). After incubation, slides were mounted onto slides with a mounting media containing 4', 6 diamidino 2 phenylindole (DAPI) and dried. Analysis was performed using a Zeiss Observer Z1 (Carl Zeiss Microscopy, Jena, Germany) microscope with Axiovision Rel 4.8 software (Zeiss, Jena, Germany) using the ×63 oil objective. Foci quantitation was conducted by counting 100 healthy cells and scoring cells with 10 or more foci as positive. Experiments were done in triplicate. Images were processed using the Adobe Photoshop software (San Jose, CA, USA).

Western blot analysis

Seventy two hours after transfection, cells were collected in PBS and then lysed for 30 min on ice with NP40 buffer containing 200 mM NaCl, 1% NP40, 50 mM Tris HCl (pH 8.0), supplemented with fresh protease inhibitors. Samples were centrifuged at 13 000 r.p.m. for 10 min at 4 °C, afterward clear lysate was extracted. A Bradford assay was used to determine protein concentration, where in samples were loaded into a SDS PAGE gel, transferred to a PVDF membrane and subsequently probed with antibodies against CHD5 (Novus, 23320002), phospho CHK2 (Thr68) (Cell Signaling, 2661), phospho histone H2AX (Ser139) antibody (Millipore, 05 636), H2AX (Bethyl, Montgomery, TX, USA, A300 082A), CHK2 (Santa Cruz, Dallas, TX, USA, sc 17748) and GAPDH (GeneTex, Irvine, CA, USA, GTX627408) followed by LI COR IRDye secondary antibodies. Detection and densitometric analysis was performed using the Odyssey system. Images were processed using the Adobe Photoshop software.

Differential expression analysis

For determination of differential expression, we extracted gene expression data from the two Gene Expression Omnibus (GEO) submissions based on the Affymetrix U133 Plus 2.0 platform: (1) GSE12654: a 22 pancreatic cancer cell line study and (2) GSE16515: 20 pancreatic patient tumors.³⁰³¹ Within



each study, after processing and normalization, we performed a genome wide filter to identify genes with 'large' expression differences among tumors, and, separately, among cell lines, using a variance approach. We define 'differential expression' as genes whose expression variability is 'large' relative to all other genes on the array, where 'large' is defined according to whether expression variability associated with a gene was greater than the 90th percentile from all the genes. It was in this analysis that CHDS emerged as highly statistically significant for further exploration in PAC.

Patient selection

A total of 80 patient specimens with adequate tissue available for analysis were obtained from a prospectively maintained surgical tissue bank of primary tumor samples. Each of patients included in the study had undergone pancreaticoduodenectomy or distal pancreatectomy for PDAC between the dates of January 2000 and October 2008. Analysis and genetic screens have been previously published using the same data set, and much of the methods for data acquisition have been previously reported.^{7,8} RFS was measured based on surveillance imaging of the chest, abdomen and pelvis, obtained at regular intervals after resection. OS was determined based on the clinical follow up data available in each patient record and was measured from the day of surgery. Relevant prognostic factors for PAC were obtained from both clinical records and individual pathology reports. Institutional Review Board approval was obtained before any data review or analysis, and patient confidentiality was maintained according to the Health Insurance and Patient Accessibility Act.

Immunohistochemical analysis

The immunohistochemical (IHC) analysis was limited to primary tumor tissue and adjacent normal tissue. Slides that had been formalin fixed and paraffin embedded were reviewed by a staff pathologist to confirm the presence and quality of tumor. The tissue was stained using an anti CHD5 antibody (Novus, 23320002). Specificity of the anti CHD5 antibody was validated by western blot analysis following siRNA silencing (Figure 1e). The percentage of cell staining and intensity of staining were factored into a previously published scoring system.8 A singe pathologist (BS) blinded to each of the patient outcomes reviewed each case. Fine granular nuclear staining was regarded as positive, and the percentage of cells that showed labeling was scored as 0 < 1%, 1 1 10%, 2 11 50% and 3 51 100%. In addition to the total percentage of staining that was given, the intensity of the staining was scored as weak 1, moderate 2 or strong 3. For the cases that demonstrated heterogeneous staining, the dominant intensity pattern was used. Representative IHC staining for each of the staining groups can be found in Figure 3. The gene expression scoring system used has been previously published and gives more weight to the percentage of cells staining, as this is thought to be more biologically significant.⁸ In general, for this scoring system, a score of ≤1 was classified as low expression, whereas a score of >1 was classified as high expression.

Statistical analysis

Univariate and multivariate Cox regression analyses were performed for all patients to examine the impact of CHD5 expression levels on both OS and RFS. Kaplan Meier log rank survival analysis was performed for RFS and OS based on CHD5 expression levels. Factors that were significant to a level of P < 0.2 on univariate analysis were included in the multivariate analysis; these included tumor size, margin status, nodal status, perineural invasion,



Figure 3. Representative immunohistochemical analysis for CHD5 expression in primary tumor tissue. The intensity of nuclear expression of CHD5 was graded as (a) weak - 1, (b) moderate - 2 or (c) strong - 3. Normal pancreatic tissue is also pictured with CHD5 staining characteristics showing islets and acini, demonstrating usually cytoplasmic positivity and staining of islets appearing more than acini. The duct epithelium show nuclear staining.



5456

lymphovascular invasion, tumor grade and CHD5 expression levels. Subset analyses were performed for patients receiving only adjuvant therapy of any kind and patients receiving only adjuvant gemcitabine based chemotherapy. All analyses were conducted using the Statistical Package for the Social Sciences 20.0 for Windows (SPSS Inc., Chicago, IL, USA).

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

We thank members of the Yu lab for helpful discussion. This work is supported by the National Center for Advancing Translational Sciences of the National Institutes of Health under Award Numbers ULI TR000454 to LEC and SBF and TLITR000456 to LEC, PanCAN/AACR 16982 to DSY, DOD/PRCRP CA110535 to DSY and Georgia Cancer Coalition 11072 to DSY.

DISCLAIMER

The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

REFERENCES

- 1 Jemal A, Siegel R, Xu J, Ward E. Cancer statistics, 2010. Cancer J Clin 2010; 60: 277 300.
- 2 Neoptolemos JP, Stocken DD, Bassi C, Ghaneh P, Cunningham D, Goldstein D et al. Adjuvant chemotherapy with fluorouracil plus folinic acid vs gemcitabine following pancreatic cancer resection: a randomized controlled trial. JAMA 2010; 304: 1073 1081.
- 3 Oettle H, Post S, Neuhaus P, Gellert K, Langrehr J, Ridwelski K et al. Adjuvant chemotherapy with gemcitabine vs observation in patients undergoing curative intent resection of pancreatic cancer: a randomized controlled trial. JAMA 2007; 297: 267 277.
- 4 Regine WF, Winter KA, Abrams RA, Safran H, Hoffman JP, Konski A et al. Fluor ouracil vs gemcitabine chemotherapy before and after fluorouracil based che moradiation following resection of pancreatic adenocarcinoma: a randomized controlled trial. JAMA 2008; 299: 1019–1026.
- 5 Iacobuzio Donahue CA, Fu B, Yachida S, Luo M, Abe H, Henderson CM et al. DPC4 gene status of the primary carcinoma correlates with patterns of failure in patients with pancreatic cancer. J Clin Oncol 2009; 27: 1806-1813.
- 6 Colbert LE, Fisher SB, Hardy CW, Hall WA, Saka B, Shelton JW et al. Pronecrotic mixed lineage kinase domain like protein expression is a prognostic biomarker in patients with early stage resected pancreatic adenocarcinoma. *Cancer* 2013; **119**: 3148–3155.
- 7 Fisher SB, Patel SH, Bagci P, Kooby DA, El Rayes BF, Staley 3rd CA et al. An analysis of human equilibrative nucleoside transporter 1, ribonucleoside reductase subunit M1, ribonucleoside reductase subunit M2, and excision repair cross complementing gene 1 expression in patients with resected pancreas adeno carcinoma: Implications for adjuvant treatment. *Cancer* 2012; 14: 514 522.
- 8 Maithel SK, Coban I, Kneuertz PJ, Kooby DA, El Rayes BF, Kauh JS et al. Differential expression of ERCC1 in pancreas adenocarcinoma: high tumor expression is associated with earlier recurrence and shortened survival after resection. AnnSurg Oncol 2011; 18: 2699 2705.
- 9 Bartkova J, Horejsi Z, Koed K, Kramer A, Tort F, Zieger K et al. DNA damage response as a candidate anti cancer barrier in early human tumorigenesis. *Nature* 2005; **434**: 864 870.
- 10 Gorgoulis VG, Vassiliou LV, Karakaidos P, Zacharatos P, Kotsinas A, Liloglou T et al. Activation of the DNA damage checkpoint and genomic instability in human precancerous lesions. *Nature* 2005; 434: 907 913.
- 11 Forbes S, Clements J, Dawson E, Bamford S, Webb T, Dogan A et al. Cosmic 2005. Br J Cancer 2006; 94: 318 322.

- 12 Bagchi A, Papazoglu C, Wu Y, Capurso D, Brodt M, Francis D et al. CHD5 is a tumor suppressor at human 1p36. Cell 2007; 128: 459 475.
- 13 Robbins CM, Tembe WA, Baker A, Sinari S, Moses TY, Beckstrom Stemberg S et al. Copy number and targeted mutational analysis reveals novel somatic events in metastatic prostate tumors. Genome Res 2011; 21: 47 55.
- 14 Law ME, Templeton KL, Kitange G, Smith J, Misra A, Feuerstein BG et al. Molecular cytogenetic analysis of chromosomes 1 and 19 in glioma cell lines. Cancer Genet Cytogenet 2005; 160: 1 14.
- 15 Wang X, Lau KK, So LK, Lam YW. CHD5 is down regulated through promoter hypermethylation in gastric cancer. J Biomed Sci 2009; 16: 95.
- 16 Gorringe KL, Choong DY, Williams LH, Ramakrishna M, Sridhar A, Qiu W et al. Mutation and methylation analysis of the chromodomain helicase DNA binding 5 gene in ovarian cancer. *Neoplasia* 2008; 10: 1253 1258.
- 17 Mulero Navarro S, Esteller M. Chromatin remodeling factor CHD5 is silenced by promoter CpG island hypermethylation in human cancer. *Epigenetics* 2008; 3: 210 215.
- 18 Ng D, Yang XR, Tucker MA, Goldstein AM. Mutation screening of CHD5 in mela noma prone families linked to 1p36 revealed no deleterious coding or splice site changes. BMC Res Note 2008; 1: 86.
- 19 Okawa ER, Gotoh T, Manne J, Igarashi J, Fujita T, Silverman KA et al. Expression and sequence analysis of candidates for the 1p36.31 tumor suppressor gene deleted in neuroblastomas. Oncogene 2008; 27: 803 810.
- 20 Mokarram P, Kumar K, Brim H, Naghibalhossaini F, Saberi firoozi M, Nouraie M et al. Distinct high profile methylated genes in colorectal cancer. PloS ONE 2009; 4: e7012.
- 21 Wu X, Zhu Z, Li W, Fu X, Su D, Fu L et al. Chromodomain helicase DNA binding protein 5 plays a tumor suppressor role in human breast cancer. Breast Cancer Res 2012; 14: R73.
- 22 Zhao R, Yan Q, Lv J, Huang H, Zheng W, Zhang B et al. CHD5, a tumor suppressor that is epigenetically silenced in lung cancer. Lung Cancer 2012; 76: 324 331.
- 23 Fujita T, Igarashi J, Okawa ER, Gotoh T, Manne J, Kolla V et al. CHD5, a tumor suppressor gene deleted from 1p36.31 in neuroblastomas. J Natl Cancer Inst 2008; 100: 940 949.
- 24 Wang L, He S, Tu Y, Ji P, Zong J, Zhang J et al. Downregulation of chromatin remodeling factor CHD5 is associated with a poor prognosis in human glioma. J Clin Neurosci 2013; 20: 958 963.
- 25 Paul S, Kuo A, Schalch T, Vogel H, Joshua Tor L, McCombie WR et al. Chd5 requires PHD mediated histone 3 binding for tumor suppression. *Cell Reports* 2013; 3: 992 102.
- 26 Matsuoka S, Ballif BA, Smogorzewska A, McDonald 3rd ER, Hurov KE, Luo J et al. ATM and ATR substrate analysis reveals extensive protein networks responsive to DNA damage. Science 2007; 316: 1160–1166.
- 27 Garcia I, Mayol G, Rodriguez E, Sunol M, Gershon TR, Rios J et al. Expression of the neuron specific protein CHD5 is an independent marker of outcome in neuro blastoma. *Mol Cancer* 2010; 9: 277.
- 28 Wong RR, Chan LK, Tsang TP, Lee CW, Cheung TH, Yim SF et al. CHD5 Down regulation Associated with Poor Prognosis in Epithelial Ovarian Cancer. Gynec Obstet Invest 2011; 72: 203 207.
- 29 Koyama H, Zhuang T, Light JE, Kolla V, Higashi M, McGrady PW et al. Mechanisms of CHD5 Inactivation in neuroblastomas. Clin Cancer Res 2012; 18: 1588 1597.
- 30 Maupin KA, Sinha A, Eugster E, Miller J, Ross J, Paulino V et al. Glycogene expression alterations associated with pancreatic cancer epithelial mesenchymal transition in complementary model systems. PloS One 2010; 5: e13002.
- 31 Pei H, Li L, Fridley BL, Jenkins GD, Kalari KR, Lingle W et al. FKBP51 affects cancer cell response to chemotherapy by negatively regulating Akt. Cancer Cell 2009; 16: 259 266.
- 32 Wang J, Chen H, Fu S, Xu ZM, Sun KL, Fu WN. The involvement of CHD5 hypermethylation in laryngeal squamous cell carcinoma. Oral Oncol 2011; 47: 601 608.
- 33 Du X, Wu T, Lu J, Zang L, Song N, Yang T et al. Decreased expression of chromodomain helicase DNA binding protein 5 is an unfavorable prognostic marker in patients with primary gallbladder carcinoma. *Clin Transl Oncol* 2012; 15: 198 204.

Supplementary Information accompanies this paper on the Oncogene website (http://www.nature.com/onc)