

Ecohab: Hyperspectral Optical Properties of Red-Tide Blooms

Helmut Maske

Center for Hydro-Optics and Remote Sensing

San Diego State University

6505 Alvarado Rd., Suite 206

San Diego, CA 92120

phone: (619) 594-2272 fax: (619) 594-8670 email: hmaske@oceanos.cicese.mx

James L. Mueller

and

Charles C. Trees

Center for Hydro-Optics and Remote Sensing

San Diego State University

6505 Alvarado Rd., Suite 206

San Diego, CA 92120

phone: (619) 594-2230 fax: (619) 594-8670 email: jim@chors.sdsu.edu, chuck@chors.sdsu.edu

Award #: N000149810778

http://www.onr.navy.mil/sci_tech/ocean/onrpgahj.htm

LONG-TERM GOALS

The goal of our project is the bio-optical characterization of dinoflagellate and other red-tide blooms, to facilitate the optical detection and monitoring of these blooms from above the sea surface.

OBJECTIVES

We want to explain the origin of the red surface color of dense dinoflagellate blooms, and the difference to other dense phytoplankton blooms that do not show this characteristic color, e.g. diatoms (brown tides) or chlorophytes. Our working hypotheses assumes that the red color of dinoflagellate blooms originate from *in vivo* fluorescence of chl *a*, but we are also considering the hypothesis that the back scattering properties of dinoflagellate cells or other bloom associated particles might change the backscattering properties and hence the upwelling radiance. The interpretation of the surface color will be based on a comparison of measured upwelling radiance and numerical simulation (Hydrolight v.4.1). The input to the simulation will be based on measured depth profiles of optically relevant variables.

APPROACH

Given the strong light absorption of water in the red, the depth of origin of the surface color is close to the surface (Maske, et al., 1998). We are measuring the fine scale (0.2 m) near surface distribution of red-tide forming organisms, and want to compare them to the apparent and inherent optical properties near the surface. To obtain near surface profiles we build an electrically triggered sampler that is taking 12 samples (4 x 1.8 liter and 8 x 300 ml) simultaneously between 0 and 2.2 m ($dz = 0.2m$). The sampler is floated away from the ship into a high density, undisturbed red-tide patch before triggering. The water samples are being processed for pigment concentration (HPLC), particulate absorption spectra

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 2001		2. REPORT TYPE		3. DATES COVERED 00-00-2001 to 00-00-2001	
4. TITLE AND SUBTITLE Ecohab: Hyperspectral Optical Properties of Red-Tide Blooms				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Center for Hydro-Optics and Remote Sensing,,San Diego State University,6505 Alvarado Rd., Suite 206,,San Diego,,CA, 92120				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT The goal of our project is the bio-optical characterization of dinoflagellate and other red-tide blooms, to facilitate the optical detection and monitoring of these blooms from above the sea surface.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 6	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

and particle counts. The vertical profiles of spectral downwelling irradiance and upwelling radiance will also be modeled with high vertical resolution, using the Hydrolight (v 4.1) radiative transfer model with measured profiles of inherent optical properties. Some unknown parameters that have to be estimated and adjusted to fit the measured radiometric data are the quantum efficiency of the fluorescence of Chl a, the volume scattering function of empty dinoflagellate thecae, and other particles (dinoflagellates, bacteria, virus).

From comparisons of modeled and measured irradiance and radiance profiles and reflectance, we expect to gain information on the dominant source of the red-tide surface color. The origin of red upwelling radiance perceived by the eye should be obvious from the spectral shape of the upwelling radiance. On the other hand, fluorescence emission would give us a well-defined spectrum with a maximum at 685nm or longer, and the enhanced elastic backscatter produced by the particles would give a wider peak.

WORK COMPLETED

A high *Lingulodinium polyedrum* bloom was measured off La Jolla/San Diego Jolla, 26 + 27. Sept. 2001 on two days. We concentrated on near surface water sampling and radiometry evaluating 3 profiles.

The 0-220 cm Linear Array Vertical Surface Sampler (LAVSS) with up to 12 sampler positions and electrical remote release was improved and used in our recent red tide sampling off San Diego. The remote release mechanism allows us to sample undisturbed surface layers away from the research platform. The sampler is more reliable now and some 350ml samplers were substituted by 1.8 liter samplers. WETLabs upgraded our HISTAR *in situ* spectrophotometer, which measures hyperspectral absorption and beam transmission. The instrument was used for near surface profiling but its performance has not yet been tested in the laboratory after the reconfiguration. The Hydrolight-2 was used to measure hyperspectral radiance and irradiance both above and below water. These data will be used to measure *in situ* and remote sensing reflectance for comparison with Hydrolight modeling results. The Hydrolight (v 4.1) model was installed and is being used to simulate measured conditions in the Monterey Bay (2000) and La Jolla (2001) blooms, and in hypothetical sensitivity studies. Some modeling results from Monterey Bay were presented at Ocean Optics XV (Maske et al. 2000).

RESULTS

The red-tide bloom observed off San Diego showed the expected very high chlorophyll a concentrations ($\sim 100 \text{ mg Chl a m}^{-3}$). About 60 percent of the homogeneously pigmented *Lingulodinium* cells had shed their thecae, the suspended thecae were still suspended in the water samples, increasing the particle concentration 60 percent. We found surprisingly high bacteria concentrations ($\sim 10^{11} \text{ cell m}^{-3}$). A preliminary sample also indicated very high virus concentration. We don't know yet if the numbers are sufficiently high to significantly affect the optical properties. In this year's samples we observed a well mixed layer of extremely high pigments and particle concentration in the top 2.2 meters, with none of the fine structure suggested by the data from Monterey Bay in August 2000. The difference could have been due to the different species composition, the lower pigment concentration or the hydrographic conditions in Monterey Bay. We expect to complete analysis of these data soon, and will include them both in our final report and in a manuscript to be submitted for publication.

PARAMETER	METHOD	COMMENTS
Phytoplankton cell concentration	Transmission microscopy	
Phytoplankton cell size	Transmission light and epifluorescence microscopy	
Empty Dinoflagellate thecae concentration	Transmission light microscopy	
Bacteria cell concentration	DAPI	
Virus cell concentration	Sybr-Green	
Absorption coefficient of particles, a_p -depigmented	Filter pad, spectrophotometry	50% duplicate samples
a -total	Filter pad	50% duplicate samples
c -total	HiStar in situ	
HPLC pigments	HiStar in situ	
Fluorometric Chl a concentration	Wright et al., 1991	50% duplicate samples
E_s (7 wavelength, continuous)	Trees et al., 2000	
	Satlantic	410, 440, 489, 511, 553, 668, 684 nm
$Lu(z, \lambda)$ and $Ed(z, \lambda)$	Hydrorad	$d\lambda = 0.5\text{nm}$,
$Lu\text{-air}(\lambda)$	HydroRad	

IMPACT/APPLICATIONS

Our characterization of the near-surface hyperspectral inherent and apparent optical properties associated with red-tide blooms will provide a basis for assessing the feasibility of using hyperspectral remote sensing methods to identify the occurrence and distribution of specific red-tide organisms.

TRANSITIONS



None during this grant period.

RELATED PROJECTS

None.



Figure 1. Vertical array surface sampler (VASS) pictured under water. 12 individual sample bottles (0-2.2m, $\Delta z=0.2m$) are simultaneously triggered electrically.



*Figure 2. *Lingulodinium polyedrum* cell without theca (about 60% of total cells) from the July bloom off San Diego, comparing light transmission and autofluorescence images (images 240 x 200 μm). Note the spherical shape without strong cell wall, this cell shape might represent a type of cycle.*



Figure 3. Empty thecae of *Lingulodinium polyedrum* from the July bloom off San Diego (images 150 x 140 μ m). The number of empty thecae found corresponds to about 60% of live cells, thus is about the number of cells found without thecae.




Figure 4. One example of a near surface profile of optically relevant biological variables, July 26, 2001 *Lingulodinium polyedrum* bloom off San Diego.