

Development of an Adaptive Vertical Coordinate Capability for Community Ocean Models

PI name: Y. Tony Song

Jet Propulsion Laboratory, 4800 Oak Grove Drive, Pasadena, CA 91109
Phone: (818) 393-4876 fax: (818) 393-6720 email: song@pacific.jpl.nasa.gov

CO-PI name: Yi Chao

Jet Propulsion Laboratory, 4800 Oak Grove Drive, Pasadena, CA 91109
Phone: (818) 354-8168 fax: (818) 393-6720 email: yc@pacific.jpl.nasa.gov

Award Number: N0001401F0162

<http://oceans-www.jpl.nasa.gov>

LONG-TERM GOALS

The goal of this project is to develop state-of-the-art modeling technologies for accurate representation of the ocean system as it evolves in time and space. The proposed adaptive vertical coordinate system is one of the innovative technologies and will be applied to the ONR-initiated Expert System for use in a variety of ocean-related areas, including coupled physical-biogeochemical studies, climate simulations using combined atmosphere and sea-ice models, and coastal ocean predictions.

OBJECTIVES

The main objective of the proposed study is to develop an adaptive vertical coordinate capability for numerical ocean models. The coordinate system will be based on the best attributes of current known vertical coordinates, hence, should have the ability to enhance vertical resolution in the surface mixed layer for proper representation of thermo-dynamical and biogeochemical processes, to resolve the bottom boundary layer for coastal ocean processes, and to retain water mass characteristics for long-term simulations. The second objective of the study is to implement the proposed vertical system to community users ocean models. This useful tool will allow diverse ocean modelers to choose the optimal vertical model structure for a hierarchy of scales from coastal to global, and to easily coordinate and share modeling resources.

APPROACH

Our technical approach is based on several, recently developed modeling techniques: a smooth transition scheme, the general pressure gradient formulation, and the finite volume method.

The idea of smoothly transitioning among different coordinate structures is based on the successful implementation of the s-coordinate formulation in the S-coordinate Rutgers University Model (SCRUM, Song and Haidvogel 1994) and its expanded version, the Regional Ocean Modeling System (ROMS, Shchepetkin and McWilliams 2001). The s-coordinate system is a generalized sigma-coordinate to permit uniformly high resolution near the surface (like the z-coordinate) and preserve the bottom-following characteristic of the sigma-coordinate. The result is a smooth transition region in the vertical column, centered at the thermohaline depth h_c . The s-coordinate has the simple, infinitely

Report Documentation Page

Form Approved
OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE 30 SEP 2001		2. REPORT TYPE		3. DATES COVERED 00-00-2001 to 00-00-2001	
4. TITLE AND SUBTITLE Development of an Adaptive Vertical Coordinate Capability for Community Ocean Models				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Jet Propulsion Laboratory,,4800 Oak Grove Drive,,Pasadena,,CA, 91109				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT The goal of this project is to develop state-of-the-art modeling technologies for accurate representation of the ocean system as it evolves in time and space. The proposed adaptive vertical coordinate system is one of the innovative technologies and will be applied to the ONR-initiated Expert System for use in a variety of ocean-related areas, including coupled physical-biogeochemical studies, climate simulations using combined atmosphere and sea-ice models, and coastal ocean predictions.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 5	19a. NAME OF RESPONSIBLE PERSON
a REPORT unclassified	b ABSTRACT unclassified	c THIS PAGE unclassified			

continuous, functional form: $z = \zeta(1+s) + h_c s + (h-h_c)C(s)$, where $\zeta(x,y,t)$ and $h(x,y)$ are surface elevation and bathymetry, respectively, and $C(s)$ is a set of s -curves, depending on two controlling parameters for surface layer and for bottom layer. By choosing the parameter appropriately, the highest resolution is achieved near the surface layer and/or bottom layer, independent of varying bottom topography.

Another key technique for developing the adaptive vertical scheme is Jacobian formulation method of calculating the pressure gradient force in a general vertical coordinate system (Song 1998). This scheme has proven to be more accurate than the traditional approach (Haney 1991) and greatly reduces hydrostatic inconsistency. Applications of this scheme are successful in ROMS for simulations of the California Current System (Song et al. 2001) and in a z -level model for BBL dynamics (Song and Chao 2000).

The finite volume method, which has been used for both incompressible and compressible flow, has two major advantages. First, it has good conservation (of mass, etc.) properties. Conservation of physical properties in ocean models is critical for long-term simulations, such as climate studies and for equilibrium circulation of coastal currents. Second, the finite volume method allows complicated computational domains to be discretized in a simpler way than either the isoparametric finite element formulation or generalized curvilinear coordinates. This method does not depend on the mesh regularity, but is suited to approximate mixed derivatives and degenerates into the finite difference method when the mesh is regular. This is another important issue in ocean modeling as the complex geometry, including islands, coastline, and headlands.

WORK COMPLETED

We have completed our first task in solving the technical problems by systematically testing the performance and evaluating the efficiency of the adaptive coordinate system. We have performed intercomparisons of different vertical structures. Three test problems, a coastal canyon, a dense water plume, and a global ocean circulation, have been carried out and compared against. Mesh doubling calculations (Roache 1990) for ascertaining grid convergence has been used as the general practice for evaluating the numerical accuracy in engineering code development. We used this technique to test our numerical schemes. For example, there is no exact solution for the canyon test problem of Haidvogel and Beckmann (1999). To have benchmark solutions for the comparisons, we run the problem with a mesh doubling and with a mesh halving resolution. Then the solutions are compared to see where they converge to the mesh doubling solution. Although this method seems costly, it is so far the most reliable method to debug the code and to generate a reliable model for the community.

RESULTS

To test our proposed method of improving the simulation of the down slope transport of dense water, we applied the EBBL formulation of Song and Chao (2000) to a z -coordinate ocean model. The coupling between the interior z -level model and the EBBL model is achieved by exchanging entrainment/detrainment and pressure gradients at the bottom layer surface, which allows temporal and spatial variations. The nice feature of this simple test problem is that it allows the model to adjust itself by transporting cool water at lower layers to the deep ocean in exchange for warmer water from upper layers, rather than by a specified inflow, which forces the system. The dynamical processes associated with coastal density fronts and dense water plumes have been investigated (e.g., Whelless and Klinck

1995). Our model solutions (Figure 1) show dense coastal water at the bottom flowing down slope, being self-advected to the right and forming a plume, which are consistent with those early results.




Figure 1: Model results for a plume test using EBBL scheme.

IMPACT/APPLICATIONS

As ocean models have become multi-disciplinary research tools, a variety of applications from coastal to global scale and from physical to biogeochemical problems require numerical model to be flexible and highly optimized (Haidvogel and Beckmann 1999). There is a community-wide need to coordinate the development, testing, maintenance, and sharing of ocean models. The restrictions among the model classes should be reduced, if possible, to allow easy communication and coordination. Our proposed adaptive vertical coordinate system provides the capability for ocean modelers to share a common modeling platform and will benefit the scientific and operational ocean modeling community at large.

TRANSITIONS

The developed adaptive vertical coordinate scheme and tested results will be made available on-line (via the Internet) to the community for further applications. We will be responsible for questions and for providing help to implement the technique into other community models.

RELATED PROJECTS

The developed adaptive vertical coordinate capability can be readily applied to several national programs for modeling studies, including the ONR Eastern Boundary Current (CBE) program, the NOAA/NSF Global Ocean Ecosystem Dynamics (GLOBEC) project, the NSF Coastal Ocean Process (CoOP) program, and the NOAA Coastal Watch program. Specifically, the proposed work will contribute to NOPP funded project, "Modeling the Central California Coastal Upwelling System: Physics, Ecosystems and Resource Management", for which Song and Chao are partners. The improved capability in the ROMS will directly benefit the NOPP effort to model the U. S. west coast circulation and variability.

REFERENCES

- Haidvogel, D. B., and A. Beckmann, 1999: *Numerical Ocean Circulation Modeling*. Imperial College Press, pp 318.
- Haney, R. L., 1991: On the pressure gradient force over steep topography in sigma coordinate ocean models. *J. Phys. Oceanogr.*, **21**, 610-619.
- Roache, P. J., 1990: Need for control of numerical accuracy, *J. Spacecraft*, **27**(2), 98-102.
- Shchepetkin, A. F., and J. C. McWilliams, 2001: Regional ocean modeling system: Development of a split, free-surface, topography-following coordinate ocean model, (manuscript).
- Song, Y. T., 1998: A general pressure gradient formulation for ocean models. Part I: Scheme design and diagnostic analysis, *Mon. Wea. Rev.*, **126**, 3212-3230.
- Song, Y. T., and Y. Chao, 2000: An embedded bottom boundary layer formulation for z-coordinate ocean models, *J. Atmos. Oceanic. Tech.*, **17**, 546-560.
- Song, Y. T., Y. Chao, P. Marchesiello, and J. McWilliams, 2001: The role of topography in coastal upwelling and cross-shore exchange. *J. Phys. Oceanogr.* (Submitted and Revised).
- Song, Y. T. and D. B. Haidvogel, 1994: A semi-implicit ocean circulation model using a generalized topography-following coordinate. *J. Comput. Phys.*, 115, 228-244.
- Whelless, G. H., and J. M. Klinck, 1995: The evolution of density-driven circulation over sloping bottom topography, *J. Phys. Oceanogr.*, **25**, 888-901.

PUBLICATIONS

- Song, T. Y., and T. Tang, 2001: Eddy-resolving simulations of the Asian Marginal Seas and Kuroshio using ROMS, *Proc. 11th PAMS/JECSS*, 101-104, April 2001, Cheju, Korea.