Lagrangian Floats for Deep Convection

Eric A. D'Asaro APL/UW 1013 NE 40th Str, Seattle, WA 98105

phone: (206) 685-2982 fax: (206) 543-6785 email: dasaro@apl.washington.edu

Award Number: N000149410025

http://poseidon.apl.washington.edu/~dasaro/HOME/

LONG-TERM GOALS

I aim to understand the process of deep convection in the ocean.

OBJECTIVES

Near surface water is mixed to great depth at a few high latitude locations, thereby forming the deep and bottom masses of the ocean. This proposal has supported the development and deployment of neutrally buoyant floats in the Labrador Sea in the winters of 1997 and 1998 and the analysis of the resulting data. These floats provide detailed information on the processes and rates of deep convection. The objective during FY01 was to finish analysis of this data.

APPROACH



Lagrangian Floats (see figure) accurately follow water motions through a combination of a density which matches that of seawater and a high drag. The density is matched to that of the ambient water by actively changing the float's volume and will stay matched, despite changes in pressure and temperature, though a combination of active control and a hull compressibility which is close to that of seawater. High drag is achieved through a large circular cloth drogue attached to the float. The horizontal motion of the float is determined by acoustic tracking (RAFOS) and its vertical motion is determined from pressure. Data is relayed at the end of the 2-month mission via satellite (ARGOS). These data are supplemented by meteorological and oceanographic data from other investigators involved in the Labrador Sea Deep Convection experiment.

maintaining the data needed, and c including suggestions for reducing	llection of information is estimated to completing and reviewing the collect this burden, to Washington Headqu, uld be aware that notwithstanding an OMB control number	tion of information Send comment arters Services, Directorate for Inf	s regarding this burden estimate of formation Operations and Reports	or any other aspect of the property of the pro	nis collection of information, Highway, Suite 1204, Arlington	
1. REPORT DATE 30 SEP 2001		2. REPORT TYPE		3. DATES COVE 00-00-2001	ERED 1 to 00-00-2001	
4. TITLE AND SUBTITLE				5a. CONTRACT NUMBER		
Lagrangian Floats	for Deep Convection	n		5b. GRANT NUMBER		
				5c. PROGRAM E	ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER		
				5e. TASK NUMBER		
				5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) APL/UW 1013 NE 40th Str,,Seattle,,WA, 98105				8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)		
				11. SPONSOR/M NUMBER(S)	ONITOR'S REPORT	
12. DISTRIBUTION/AVAII Approved for publ	LABILITY STATEMENT ic release; distributi	ion unlimited				
13. SUPPLEMENTARY NO	OTES					
bottom masses of the buoyant floats in the These floats provideduring FY01 was to	r is mixed to great do he ocean. This propo he Labrador Sea in t le detailed informati o finish analysis of t	osal has supported the winters of 1997 ion on the processe	the development a and 1998 and the	and deployme analysis of the	ent of neutrally he resulting data.	
15. SUBJECT TERMS						
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON	
a REPORT unclassified	ь abstract unclassified	c THIS PAGE unclassified	Same as Report (SAR)	4		

Report Documentation Page


Form Approved OMB No. 0704-0188

WORK COMPLETED

Our data from both 1997 (13 floats) and 1998 (7 floats) has been completely processed and calibrated and most of the basic data analysis completed. Two papers have been accepted for publication in a special Labrador Sea Convection issue of the *Journal of Physical Oceanography*, one describing the floats and data and a second, with Ramsey Harcourt of NPS, comparing these with the predictions of an LES model. Much of this work was done by graduate student Elizabeth Steffen as part of her Ph.D. work.

RESULTS

The figure shows the strong correlation between vertical velocity as measured by the floats and the surface heat flux, from calibrated NCEP surface heat flux. This shows that mixing in the Labrador sea is really convection: it is driven by heat flux and is primarily vertical, not slant.



IMPACT/APPLICATIONS

These data have produced direct confirmation of the existing theoretical ideas about how the "rapid mixing" phase of deep convection works. This has direct implications for the parameterization of convective mixing in numerical models.

TRANSITIONS

None

RELATED PROJECTS

Relatives of these floats are being used in the ONR supported CBLAST study of hurricanes and in studies of circulation, upwelling and mixing off the Oregon Coast.

PUBLICATIONS

Harcourt et al. (2001), Fully Lagrangian floats in Labrador Sea deep convection: Comparison of Numerical and experimental results, J. Phys. Oceanogr., in press.

Steffen, E. and E. D'Asaro (2001), Deep convection in the Labrador Sea as observed by Lagrangian Floats, J. Phys. Oceanogr., in press