



EDGEWOOD CHEMICAL BIOLOGICAL CENTER

U.S. ARMY RESEARCH, DEVELOPMENT AND ENGINEERING COMMAND
Aberdeen Proving Ground, MD 21010-5424

ECBC-TR-1333

EXTRACTION AND ANALYSIS OF V-TYPE AGENTS (VX, RVX, CVX, AND VM) FROM VARIOUS FOOD MATRICES BY ULTRA-PERFORMANCE LIQUID CHROMATOGRAPHY-TIME-OF-FLIGHT MASS SPECTROMETRY

Sue Y. Bae
Mark D. Winemiller

RESEARCH AND TECHNOLOGY DIRECTORATE

December 2015

Approved for public release; distribution is unlimited.



Disclaimer

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorizing documents.

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 h per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. **PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.**

1. REPORT DATE (DD-MM-YYYY) XX-12-2015		2. REPORT TYPE Final	3. DATES COVERED (From - To) Oct 2014 – Jun 2015		
4. TITLE AND SUBTITLE Extraction and Analysis of V-Type Agents (VX, RVX, CVX, and VM) from Various Food Matrices by Ultra-Performance Liquid Chromatography–Time-of-Flight Mass Spectrometry			5a. CONTRACT NUMBER		
			5b. GRANT NUMBER		
			5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S) Bae, Sue Y.; and Winemiller, Mark D.			5d. PROJECT NUMBER		
			5e. TASK NUMBER		
			5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Director, ECBC, ATTN: RDCB-DRC-C, APG, MD 21010-5424			8. PERFORMING ORGANIZATION REPORT NUMBER ECBC-TR-1333		
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSOR/MONITOR'S ACRONYM(S)		
			11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT: Liquid chromatography–electrospray ionization mass spectrometry with positive-ion modes of operation was used to analyze several V-type chemical warfare agents (VX, RVX, CVX, and VM) in various food matrices. The development of a solid-phase extraction method using normal-phase silica gel columns for the extraction of V-type agents in food is described. In support of this objective, we examined select food samples using individual agents; mixtures of agents have not been studied. Various agent quantities, ranging from 1.7 to 3.1 mg, were spiked into food samples. The Agent Chemistry Branch at the U.S. Army Edgewood Chemical Biological Center has developed three analytical techniques for use, depending on the matrix. These matrices include orange juice, apple juice, whole milk, 2% reduced fat milk, Egg Beaters egg whites, tomato sauce, and several meats, including hamburger meat (80% lean and 20% fat), hot dogs, chicken nuggets, and turkey deli meat (99% fat free). The total percent recoveries (and percent relative standard deviations) for VX, RVX, CVX, and VM in various food samples are reported.					
15. SUBJECT TERMS V-type agents Chemical warfare agent (CWA) Food Ultra-performance liquid chromatography–time-of-flight mass spectrometry (UPLC–TOF-MS)					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			19b. TELEPHONE NUMBER (include area code)
U	U	U	UU	36	Renu B. Rastogi (410) 436-7545

Blank

PREFACE

This work was started in October 2014 and completed in June 2015.

The use of either trade or manufacturers' names in this report does not constitute an official endorsement of any commercial products. This report may not be cited for purposes of advertisement.

This report has been approved for public release.

Blank

CONTENTS

1.	INTRODUCTION	1
2.	EXPERIMENTAL METHODS.....	2
2.1	Reagents and Chemicals	2
2.2	Instrumentation	2
2.3	Sample Preparation and Extraction Procedures	3
2.3.1	Group 1: Apple Juice, Orange Juice, Whole Milk, and 2% Reduced Fat Milk	3
2.3.2	Group 2: Egg Whites and Tomato Sauce.....	3
2.3.3	Group 3: Hot Dogs, Chicken Nuggets, Turkey Deli Meat, and Ground Beef.....	3
2.3.4	Extraction Procedures	4
3.	RESULTS AND DISCUSSION	4
3.1	LC Separation and Analytical Figures of Merit.....	4
3.2	Extraction of VX from Foodstuffs	5
3.3	Extraction of RVX from Foodstuffs	9
3.4	Extraction of CVX from Foodstuffs	12
3.5	Extraction of VM from Foodstuffs	16
4.	CONCLUSION.....	19
	LITERATURE CITED	21
	ACRONYMS AND ABBREVIATIONS	25

FIGURES

1.	Structure of nerve agents VX, RVX, CVX, and VM.....	2
2.	A RediSep Rf normal-phase silica column.....	4
3.	A representative TIC and EIC for VX extracted from apple juice	6
4.	A representative mass spectrum for VX extracted from apple juice	7
5.	External calibration curve for VX in CH ₃ CN	8
6.	A representative TIC and EIC for RVX extracted from apple juice.....	10
7.	A representative mass spectrum for RVX extracted from apple juice.....	11
8.	External calibration curve for RVX in CH ₃ CN	12
9.	A representative TIC and EIC for CVX extracted from apple juice.....	13
10.	A representative mass spectrum for CVX extracted from apple juice.....	14
11.	External calibration curve for CVX in CH ₃ CN	15
12.	A representative TIC and EIC for VM extracted from apple juice.....	17
13.	A representative mass spectrum for VM extracted from apple juice.....	18
14.	External calibration curve for VM in CH ₃ CN	19

TABLES

1.	Analytical Figures of Merit Obtained for V-Type Agents.....	5
2.	Percent Recoveries of Extracted VX from Various Food Matrices.....	8
3.	Percent Recoveries of Extracted RVX from Various Food Matrices.....	9
4.	Percent Recoveries of Extracted CVX from Various Food Matrices.....	15
5.	Percent Recoveries of Extracted VM from Various Food Matrices.....	16

EXTRACTION AND ANALYSIS OF V-TYPE AGENTS (VX, RVX, CVX, AND VM) FROM VARIOUS FOOD MATRICES BY ULTRA-PERFORMANCE LIQUID CHROMATOGRAPHY–TIME-OF-FLIGHT MASS SPECTROMETRY

1. INTRODUCTION

The continued threat from traditional chemical warfare agents (CWAs) such as VX (Figure 1) is evident on an almost daily basis, as current events in Syria have demonstrated. Issues ranging from food and environmental safety to compliance with treaties makes the need for low-level detection of VX of great importance. The mere existence of these molecules in either the environment or the food supply could indicate a compliance breach, even if the CWA levels were not high enough to cause any real personal harm.

Although the detection of VX metabolites and adducts from food and biological sample matrices has been reported,^{1–9} limited literature exists regarding direct detection of the actual CWAs in food.^{10,11} The pesticide literature often includes sample-preparation techniques that are commercially available and affordable, such as solid-phase extraction cartridges^{12–20} or QuEChERS systems (Quick, Easy, Cheap, Effective, Rugged, and Safe);^{21–29} however, the CWA literature seems to focus more on new techniques and specialized equipment that may not be as readily accessible to every laboratory.^{9,30–35}

This document reports the efforts of the Agent Chemistry Team from the Research and Technology Directorate of the U.S. Army Edgewood Chemical Biological Center (ECBC; Aberdeen Proving Ground, MD) in developing new extraction and analytical detection methodologies using liquid chromatography–mass spectrometry (LC–MS). The objective of this task was to provide development and laboratory support for extraction of V-type agents (Figure 1) from various food samples. This includes detection and quantitative and qualitative analysis of complex matrices such as foods with high salt and fat contents. In support of this objective, we examined 10 food samples using individual agents. Mixtures of agents have not been studied. Apple juice, orange juice, whole milk, 2% reduced fat milk, Egg Beaters processed egg whites, tomato sauce, precooked turkey deli meat (99% fat free), chicken nuggets, hot dogs, and 80/20 hamburger meat (80% lean and 20% fat) represented food types that are commonly associated with school lunch programs. The choice of food types arose from collaborations and conversations with U.S. Department of Agriculture personnel. Foods were tested using commercially available normal-phase separation columns.

The use of ultra-performance liquid chromatography with time-of-flight mass spectroscopy (UPLC–TOF-MS), or any comparable high-resolution LC–MS, has become more common. From an affordability standpoint, it is currently within reach for most laboratories. For this work, extracted agent was analyzed using UPLC–TOF-MS, and percent recovery was calculated from an external calibration curve.




Figure 1. Structure of nerve agents VX, RVX, CVX, and VM.

2. EXPERIMENTAL METHODS

2.1 Reagents and Chemicals

The following nerve agents were provided by ECBC:

- VX: *O*-ethyl *S*-[2-(diisopropylamino)ethyl] methylphosphonothioate;
- CVX: *O*-butyl *S*-[2-(diethylamino)ethyl] methylphosphonothioate;
- RVX: *S*-[2-(diethylamino)ethyl] *O*-isobutyl methylphosphonothioate; and
- VM: *O*-ethyl *S*-(2-diethylaminoethyl) methylphosphonothioate.

For all nerve agents, purity was >99%. All reagents and solvents were LC-MS grade. Acetonitrile, water, and triethylamine (TEA) were purchased from Sigma-Aldrich (St. Louis, MO). Apple juice, orange juice, whole milk, 2% reduced fat milk, Egg Beaters egg whites, tomato sauce, and hot dog food samples were purchased from a local grocery store (Food Lion; Edgewood, MD).

2.2 Instrumentation

All samples were characterized using an Acquity UPLC Synapt G2-S system (Waters Corp.; Milford, MA) equipped with an electrospray ionization (ESI) interface. The sampling cone voltage was 20 V. The source and desolvation temperatures were 120 and 500 °C,

respectively, and the nitrogen desolvation gas flow rate was 800 L/h. The LC–ESI–TOF–multiple reaction monitoring and LC–ESI–TOF–MS data were acquired in positive-ion scan mode over a mass range of 50–1200 Da. The leucine–enkephalin solution (1 ng/μL) was used as reference mass with a flow rate of 10 μL/min. The LC separations for all extracted samples were performed on a Waters Acquity UPLC BEH amide column (50 × 2.1 mm, 1.7 μm). The mobile phase consisted of 10 mM ammonium acetate and 0.04% NH₄OH in 90/10 (v/v %) H₂O/CH₃CN (mobile phase A) and 100% acetonitrile (mobile phase B) with a 10 μL injection volume. Separation was achieved using an isocratic condition of 20/80 (v/v %) A/B with a flow rate of 0.5 mL/min. A thermostatted column-manager compartment was used to maintain the column temperature at 35 °C and the test samples at 5 °C.

2.3 Sample Preparation and Extraction Procedures

2.3.1 Group 1: Apple Juice, Orange Juice, Whole Milk, and 2% Reduced Fat Milk

Juicy Juice apple juice (Nestlé USA; Glendale, CA), Minute Maid orange juice (Coca Cola Company; Atlanta, GA), and Food Lion brand whole milk and 2% reduced fat milk were purchased from the Edgewood Food Lion supermarket. The pH of each sample was measured and recorded before the sample was extracted. Each 2 mL sample was spiked with approximately 1.8–2.0 mg of V-type agent. The sample was diluted with 20 mL of CH₃CN. The food sample was then passed through a RediSep Rf column (Teledyne Isco; Lincoln, NE), and the eluents were collected.

2.3.2 Group 2: Egg Whites and Tomato Sauce

Egg Beaters Original egg whites (ConAgra Foods; Omaha, NE) and Ragú tomato sauce (Old World Style Ragú flavored with meat; R&B Foods; Mount Prospect, IL) were purchased from the Edgewood Food Lion supermarket. The pH of each sample was measured and recorded before the sample was extracted. Approximately 5 g of each food sample was spiked with the desired V-type agent and diluted with 10 mL of CH₃CN. The mixture was centrifuged for 15 min at 6000 rpm, and the supernatant was decanted. A second portion of 10 mL of CH₃CN was added, and the mixture was vortexed or sonicated for 1 min and again centrifuged for 15 min at 6000 rpm. The supernatant was removed, and the first and second portions were combined and passed through a silica gel column. The eluent was collected for analysis.

2.3.3 Group 3: Hot Dogs, Chicken Nuggets, Turkey Deli Meat, and Ground Beef

Orioles hot dogs (Esskay; Baltimore, MD), Smart Option chicken nuggets (Food Lion private brand), Buddig Original Deli Thin turkey deli meat (Carl Buddig & Co.; Homewood, IL), and 80/20 ground beef chuck (Food Lion sourced) were purchased from the Edgewood Food Lion supermarket. Approximately 5 g of each food sample was spiked with V-type agent and diluted with 10 mL of CH₃CN. The whole sample was homogenized using a Polytron homogenizer (Kinematica; Luzern, Switzerland) at 20,000 rpm for 1–2 min. The mixture was then centrifuged for 15 min at 6000 rpm, and the supernatant was removed. A second portion of 10 mL of CH₃CN was added, and the sample was vortexed or sonicated for

1 min and centrifuged for 15 min at 6000 rpm, and the supernatant was removed. The first and second portions of supernatant were combined and passed through a silica gel column, and the eluent was collected for analysis.

2.3.4 Extraction Procedures

A packed RediSep Rf normal-phase silica gel column (shown in Figure 2) was used in this study to separate the V-type agents from the food samples. First, the RediSep Rf column was eluted with 25 mL of 0.1% TEA/CH₃CN using in-house air to pass the solution through the column. The 0.1% TEA/CH₃CN solution was collected for later use. Second, the supernatant was passed through the column, and the sample was collected from the column. Third, 1 mL of 0.1% TEA/CH₃CN solution was added to the column and pushed slightly into the silica gel until 1 mL had just cleared the top of the silica gel. This step was repeated four times. Finally, the remaining 0.1% TEA/CH₃CN solution was added to the column and passed through the bed. The final solution was diluted with mobile phase and analyzed using LC–MS.




Figure 2. A RediSep Rf normal-phase silica column.

3. RESULTS AND DISCUSSION

3.1 LC Separation and Analytical Figures of Merit

For LC–MS analysis, the MS system was operated to obtain both total ion chromatograms (TICs) at m/z 50–1200 and extracted ion chromatograms (EICs) at m/z 268.1505 for VX, RVX, and CVX. EICs at m/z 240.1189 were obtained for VM. The TIC mode was used to ensure identification of any hydrolysis products. No hydrolysis products of V agent were found from these extracted samples. The EIC mode was used to determine the limits of detection (LODs), limits of quantitation (LOQs), and the linear dynamic ranges (LDRs) of the V-type agents. The calibration curve was plotted over concentration ranges of 0.057–960, 0.065–925, 0.126–0.700, and 0.5–0.890 ng/mL for VX, RVX, CVX, and VM, respectively, using 1 μ L injections at each concentration level. The LODs for the nerve agents were calculated using 2 μ L injections at concentrations as low as 0.1 ng/mL with a signal-to-noise ratio of 3:1. The LOQs for the analyte were calculated with a signal-to-noise ratio of 10:1. The linear regression equations were calculated by a least-squares analysis for the LDRs, LODs, and LOQ. The linear regression equations and the correlation coefficients are tabulated in Table 1.

Table 1. Analytical Figures of Merit Obtained for V-Type Agents^a

Nerve Agent	LDR (ng/mL)	LOD (fg on column)	LOQ (fg on column)	Correlation Coefficient
VX	0.057–960	115	570	0.9984
RVX	0.065–925	65	130	0.9959
CVX	0.126–700	126	253	0.9956
VM	0.5–0.890	248	495	0.9953

^aNerve agent standard in CH₃CN.

3.2 Extraction of VX from Foodstuffs

In Figure 3b, a representative TIC is shown for each agent in various food matrices. For each agent, the EIC was obtained from the TIC of the extracted agent and is shown in Figure 3c. The mass spectrum, shown in Figure 4, exhibits mass ions at m/z 268.1501 due to $[M + H]^+$ and at m/z 128.1439 due to the loss of *O*-ethyl *S*-hydrogen methylphosphonothioate, $[M - C_3H_9O_2PS]^+$ for VX. The percent recovery calculations were based on an external calibration curve of VX (Figure 5). The results showed a >90% recovery of VX from various food matrices (Table 2).




Figure 3. A representative TIC and EIC for VX extracted from apple juice: (a) sample blank, (b) TIC, and (c) EIC.

L




Figure 4. A representative mass spectrum for VX extracted from apple juice.




Figure 5. External calibration curve for VX in CH₃CN.

Table 2. Percent Recoveries of Extracted VX from Various Food Matrices

Food Matrix	% Recovered ± % RSD
Apple juice	94.2 ± 1.17
Orange juice	96.7 ± 2.18
Reduced fat (2%) milk	93.8 ± 2.88
Whole milk	97.4 ± 2.78
Egg Beaters egg whites	96.8 ± 1.07
Tomato sauce	97.1 ± 2.40
Turkey deli meat	93.8 ± 2.27
Hot dogs	92.4 ± 1.86
Chicken nuggets	97.8 ± 2.02
Ground beef (80/20)	96.1 ± 1.91

3.3 Extraction of RVX from Foodstuffs

In Figure 6b, a representative TIC is shown for each agent in various food matrices. For each agent, the EIC was obtained from the TIC of the extracted agent and is shown in Figure 6c. The mass spectrum, shown in Figure 7, exhibits mass ions at m/z 268.1505 due to $[M + H]^+$ and m/z 100.1127 due to the loss of *O*-isobutyl *S*-hydrogen methylphosphonothioate, $[M - C_5H_{13}O_2PS]^+$ for RVX. The percent recovery calculations were based on an external calibration curve of RVX (Figure 8). The results showed a >90% recovery of RVX from various food matrices (Table 3).

Table 3. Percent Recoveries of Extracted RVX from Various Food Matrices

Food Matrix	% Recovered \pm % RSD
Apple juice	93.8 \pm 2.62
Orange juice	96.4 \pm 1.95
Reduced fat (2%) milk	96.4 \pm 1.18
Whole milk	95.4 \pm 3.73
Egg Beaters egg whites	93.9 \pm 1.76
Tomato sauce	94.1 \pm 2.73
Turkey deli meat	96.6 \pm 2.31
Hot dogs	93.4 \pm 3.51
Chicken nuggets	94.7 \pm 2.95
Ground beef (80/20)	96.0 \pm 2.18




Figure 6. A representative TIC and EIC for RVX extracted from apple juice: (a) sample blank, (b) TIC, and (c) EIC.




Figure 7. A representative mass spectrum for RVX extracted from apple juice.




Figure 8. External calibration curve for RVX in CH₃CN.

3.4 Extraction of CVX from Foodstuffs

In Figure 9b, a representative TIC is shown for each agent in various food matrices. The EIC for each agent was extracted from the TIC and is shown in Figure 9c. The mass spectrum, shown in Figure 10, exhibits mass ions at m/z 268.1501 due to $[M + H]^+$ and m/z 100.1124 due to the loss of *O*-isobutyl *S*-hydrogen methylphosphonothioate, $[M - C_5H_{13}O_2PS]^+$ for CVX. The percent recovery calculations were based on an external calibration curve of CVX (Figure 11). The results showed a >90% recovery of CVX from various food matrices (Table 4).




Figure 9. A representative TIC and EIC for CVX extracted from apple juice: (a) matrix blank, (b) TIC, and (c) EIC.




Figure 10. A representative mass spectrum for CVX extracted from apple juice.




Figure 11. External calibration curve for CVX in CH₃CN.

Table 4. Percent Recoveries of Extracted CVX from Various Food Matrices

Food Matrix	% Recovered ± % RSD
Apple juice	97.8 ± 1.76
Orange juice	97.6 ± 1.58
Reduced fat (2%) milk	98.6 ± 0.58
Whole milk	99.8 ± 1.61
Egg Beaters egg whites	96.9 ± 2.02
Tomato sauce	97.2 ± 2.52
Turkey deli meat	94.9 ± 2.55
Hot dogs	93.5 ± 1.81
Chicken nuggets	96.8 ± 1.57
Ground beef (80/20)	94.5 ± 1.90

3.5 Extraction of VM from Foodstuffs

In Figure 12b, a representative TIC is shown for each agent in various food matrices. For each agent, the EIC was obtained from the TIC of the extracted agent and is shown in Figure 12c. The mass spectrum, shown in Figure 13, exhibits mass ions at m/z 240.1187 due to $[M + H]^+$ and m/z 100.1124 due to the loss of *O*-ethyl *S*-hydrogen methylphosphonothioate, $[M - C_3H_9O_2PS]^+$ for VM. The percent recovery calculations were based on an external calibration curve of VM (Figure 14). The results showed a >90% recovery of VM from various food matrices (Table 5).

Table 5. Percent Recoveries of Extracted VM from Various Food Matrices

Food Matrix	% Recovered \pm % RSD
Apple juice	93.7 \pm 4.09
Orange juice	96.1 \pm 2.30
Reduced fat (2%) milk	98.1 \pm 3.04
Whole milk	99.4 \pm 1.31
Egg Beaters egg whites	94.4 \pm 2.82
Tomato sauce	97.5 \pm 2.27
Turkey deli meat	94.5 \pm 4.48
Hot dogs	91.4 \pm 1.93
Chicken nuggets	95.8 \pm 3.80
Ground beef (80/20)	92.2 \pm 2.07




Figure 12. A representative TIC and EIC for VM extracted from apple juice: (a) matrix blank, (b) TIC, and (c) EIC.




Figure 13. A representative mass spectrum for VM extracted from apple juice.




Figure 14. External calibration curve for VM in CH₃CN.

4. CONCLUSION

In this study, we set out to examine whether a commercial solution could be found for the extraction of nerve agents from various food matrices. Ideally, the method would be simple and easily available anywhere in the United States. Recoveries were >90% for the extraction techniques developed for the V-type agents in all food matrices. This report details the extraction and analysis techniques used to study V-type CWAs. The extraction method was easy to use, and it was easy to observe parts-per-million levels of VX, RVX, CVX, and VM in single food matrices.

Blank

LITERATURE CITED

1. Røen, B.T.; Sellevåg, S.R.; Dybendal, K.E.; Lundanes, E. Trace Determination of Primary Nerve Agent Degradation Products in Aqueous Soil Extracts by On-Line Solid Phase Extraction-Liquid Chromatography-Mass Spectrometry Using ZrO₂ for Enrichment. *J. Chromatogr. A* **2014**, *1329*, 90–97.
2. Bao, Y.; Liu, Q.; Chen, J.; Lin, Y.; Wu, B.D.; Xie, J.W. Quantification of Nerve Agent Adducts with Albumin in Rat Plasma Using Liquid Chromatography-Isotope Dilution Tandem Mass Spectrometry. *J. Chromatogr. A* **2012**, *1229*, 164–171.
3. Koller, M.; Becker, C.; Thiermann, H.; Worek, F. GC-MS and LC-MS Analysis of Nerve Agents in Body Fluids: Intra-Laboratory Verification Test Using Spiked Plasma and Urine Samples. *J. Chromatogr. B Analyt. Technol. Biomed. Life Sci.* **2010**, *878*, 1226–1233.
4. Mawhinney, D.B.; Hamelin, E.I.; Fraser, R.; Silva, S.S.; Pavlopoulos, A.J.; Kobelski, R.J. The Determination of Organophosphonate Nerve Agent Metabolites in Human Urine by Hydrophilic Interaction Liquid Chromatography Tandem Mass Spectrometry. *J. Chromatogr. B Analyt. Technol. Biomed. Life Sci.* **2007**, *852*, 235–243.
5. Ciner, F.L.; McCord, C.E.; Plunkett, R.W.; Martin, M.F.; Croley, T.R. Isotope Dilution LC/MS/MS for the Detection of Nerve Agent Exposure in Urine. *J. Chromatogr. B Analyt. Technol. Biomed. Life Sci.* **2007**, *846*, 42–50.
6. Noort, D.; Fidler, A.; van der Schans, M.J.; Hulst, A.G. Verification of Exposure to Organophosphates: Generic Mass Spectrometric Method for Detection of Human Butyrylcholinesterase Adducts. *Anal. Chem.* **2006**, *78*, 6640–6644.
7. Tsuge, K.; Seto, Y. Detection of Human Butyrylcholinesterase-Nerve Gas Adducts by Liquid Chromatography-Mass Spectrometric Analysis after In Gel Chymotryptic Digestion. *J. Chromatogr. B Analyt. Technol. Biomed. Life Sci.* **2006**, *838*, 21–30.
8. Crow, B.S.; Pantazides, B.G.; Quinones-Gonzalez, J.; Garton, J.W.; Carter, M.D.; Perez, J.W.; Watson, C.M.; Tomcik, D.J.; Crenshaw, M.D.; Brewer, B.N.; Riches, J.R.; Stubbs, S.J.; Read, R.W.; Evans, R.A.; Thomas, J.D.; Blake, T.A.; Johnson, R.C. Simultaneous Measurement of Tabun, Sarin, Soman, Cyclosarin, VR, VX, and VM Adducts to Tyrosine in Blood Products by Isotope Dilution UHPLC–MS/MS. *Anal. Chem.* **2014**, *86*, 10397–10405.
9. Knaack, J.S.; Zhou, Y.T.; Abney, C.W.; Prezioso, S.M.; Magnuson, M.; Evans, R.; Jakubowski, E.M.; Hardy, K.; Johnson, R.C. High-Throughput Immunomagnetic Scavenging Technique for Quantitative Analysis of Live VX Nerve Agent in Water, Hamburger, and Soil Matrixes. *Anal. Chem.* **2012**, *84*, 10052–10057.
10. Feng, C.L.; Zhou, Q.X.; Hu, Q.Y. Analysis of 7 Chemical Warfare Agents in Contaminated Grain by Gas Chromatography–Flame Photometric Detection. *Chinese J. Anal. Chem.* **2000**, *28*, 1245–1247.
11. Klemm, M. Device for Extracting Electrically Charged Molecules. German Patent DE10149875, 3 July 2003.
12. Chen, J.; Duan, C.F.; Guan, Y.F. Sorptive Extraction Techniques in Sample Preparation for Organophosphorus Pesticides in Complex Matrices. *J. Chromatogr. B* **2010**, *878*, 1216–1225.

13. Han, Q.; Wang, Z.H.; Xia, J.F.; Zhang, X.Q.; Wang, H.W.; Ding, M.Y. Application of Graphene for the SPE Clean-Up of Organophosphorus Pesticides Residues from Apple Juices. *J. Sep. Sci.* **2014**, *37*, 99–105.
14. Liu, M.; Hashi, Y.; Song, Y.Y.; Lin, J.M. Simultaneous Determination of Carbamate and Organophosphorus Pesticides in Fruits and Vegetables by Liquid Chromatography–Mass Spectrometry. *J. Chromatogr. A* **2005**, *1097*, 183–187.
15. Picó, Y.; Fernández, M.; Ruiz, M.J.; Font, G. Current Trends in Solid-Phase-Based Extraction Techniques for the Determination of Pesticides in Food and Environment. *J. Biochem. Biophys. Methods* **2007**, *70*, 117–131.
16. Picó, Y.; Moltó, J.C.; Mañes, J.; Font, G. Solid Phase Techniques in the Extraction of Pesticides and Related Compounds from Foods and Soils. *J. Microcolumn Sep.* **1994**, *6*, 331–359.
17. Schenck, F.J.; Donoghue, D.J. Determination of Organochlorine and Organophosphorus Pesticide Residues in Eggs Using a Solid Phase Extraction Cleanup. *J. Agric. Food Chem.* **2000**, *48*, 6412–6415.
18. Schenck, F.J.; Lehotay, S.J. Does Further Clean-Up Reduce the Matrix Enhancement Effect in Gas Chromatographic Analysis of Pesticide Residues in Food? *J. Chromatogr. A* **2000**, *868*, 51–61.
19. Schenck, F.J.; Podhorniak, L.V.; Hobbs, J.; Casanova, J.; Donoghue, D. Liquid Chromatographic Determination of N-Methyl Carbamate Pesticide Residues at Low Parts-per-Billion Levels in Eggs. *J. AOAC Int.* **2006**, *89*, 196–200.
20. Vidal, J.L.M.; Plaza-Bolanos, P.; Romero-Gonzalez, R.; Frenich, A.G. Determination of Pesticide Transformation Products: A Review of Extraction and Detection Methods. *J. Chromatogr. A* **2009**, *1216*, 6767–6788.
21. Bruzzoniti, M.C.; Checchini, L.; De Carlo, R.M.; Orlandini, S.; Rivoira, L.; Del Bubba, M. QuEChERS Sample Preparation for the Determination of Pesticides and Other Organic Residues in Environmental Matrices: A Critical Review. *Anal. Bioanal. Chem.* **2014**, *406*, 4089–4116.
22. Carneiro, R.P.; Oliveira, F.A.S.; Madureira, F.D.; Silva, G.; de Souza, W.R.; Lopes, R.P. Development and Method Validation for Determination of 128 Pesticides in Bananas by Modified QuEChERS and UHPLC–MS/MS Analysis. *Food Control* **2013**, *33*, 413–423.
23. Sinha, S.N.; Vasudev, K.; Rao, M.V.V. Quantification of Organophosphate Insecticides and Herbicides in Vegetable Samples Using the “Quick Easy Cheap Effective Rugged and Safe” (QuEChERS) Method and a High-Performance Liquid Chromatography–Electrospray Ionisation–Mass Spectrometry (LC–MS/MS) Technique. *Food Chem.* **2012**, *132*, 1574–1584.
24. Seiber, J.N.; Kleinschmidt, L.A. Contributions of Pesticide Residue Chemistry to Improving Food and Environmental Safety: Past and Present Accomplishments and Future Challenges. *J. Agric. Food Chem.* **2011**, *59*, 7536–7543.
25. Lehotay, S.J. QuEChERS Sample Preparation Approach for Mass Spectrometric Analysis of Pesticide Residues in Foods. *Methods Mol. Biol.* **2011**, *747*, 65–91.
26. Fernandes, V.C.; Domingues, V.F.; Mateus, N.; Delerue-Matos, C. Determination of Pesticides in Fruit and Fruit Juices by Chromatographic Methods. An Overview *J. Chromatogr. Sci.* **2011**, *49*, 715–730.

27. Chung, S.W.; Chan, B.T. Validation and Use of a Fast Sample Preparation Method and Liquid Chromatography-Tandem Mass Spectrometry in Analysis of Ultra-Trace Levels of 98 Organophosphorus Pesticide and Carbamate Residues in a Total Diet Study Involving Diversified Food Types *J. Chromatogr. A* **2010**, *1217*, 4815–4824.
28. Schenck, F.; Wong, J.; Lu, C.S.; Li, J.; Holcomb, J.R.; Mitchell, L.M. Multiresidue Analysis of 102 Organophosphorus Pesticides in Produce at Parts-per-Billion Levels Using a Modified QuEChERS Method and Gas Chromatography with Pulsed Flame Photometric Detection. *J. AOAC Int.* **2009**, *92*, 561–573.
29. Nguyen, T.D.; Yu, J.E.; Lee, D.M.; Lee, G.H. A Multiresidue Method for the Determination of 107 Pesticides in Cabbage and Radish Using QuEChERS Sample Preparation Method and Gas Chromatography Mass Spectrometry *Food Chem.* **2008**, *110*, 207–213.
30. Kolakowski, B.M.; D'Agostino, P.A.; Chenier, C.; Mester, Z. Analysis of Chemical Warfare Agents in Food Products by Atmospheric Pressure Ionization-High Field Asymmetric Waveform Ion Mobility Spectrometry-Mass Spectrometry. *Anal. Chem.* **2007**, *79*, 8257–8265.
31. Kanamori-Kataoka, M.; Seto, Y. Laboratory Identification of the Nerve Gas Hydrolysis Products Alkyl Methylphosphonic Acids and Methylphosphonic Acid, by Gas Chromatography–Mass Spectrometry after *tert*-Butyldimethylsilylation. *J. Health Sci.* **2008**, *54*, 513–523.
32. Karpas, Z. Applications of Ion Mobility Spectrometry (IMS) in the Field of Foodomics. *Food Res. Int.* **2013**, *54*, 1146–1151.
33. Simonian, A.L.; Flounders, A.W.; Wild, J.R. FET-Based Biosensors for the Direct Detection of Organophosphate Neurotoxins. *Electroanalysis* **2004**, *16*, 1896–1906.
34. Vautz, W.; Zimmermann, D.; Hartmann, M.; Baumbach, J.I.; Nolte, J.; Jung, J. Ion Mobility Spectrometry for Food Quality and Safety. *Food Addit. Contam.* **2006**, *23*, 1064–1073.
35. Lumor, S.E.; Diez-Gonzalez, F.; Labuza, T.P. Detection of Warfare Agents in Liquid Foods Using the Brine Shrimp Lethality Assay. *J. Food Sci.* **2011**, *76*, T16–T19.

Blank

ACRONYMS AND ABBREVIATIONS

80/20	80% lean and 20% fat
CVX	<i>O</i> -butyl <i>S</i> -[2-(diethylamino)ethyl] methylphosphonothioate
CWA	chemical warfare agent
ECBC	U.S. Army Edgewood Chemical Biological Center
EIC	extracted ion chromatogram
ESI	electrospray ionization
LC	liquid chromatography
LDR	linear dynamic range
LOD	limit of detection
LOQ	limit of quantitation
MS	mass spectroscopy
QuEChERS	Quick, Easy, Cheap, Effective, Rugged, and Safe
R VX	<i>S</i> -[2-(diethylamino)ethyl] <i>O</i> -isobutyl methylphosphonothioate
TEA	triethylamine
TIC	total ion chromatogram
TOF	time of flight
UPLC	ultra-performance liquid chromatography
VM	<i>O</i> -ethyl <i>S</i> -(2-diethylaminoethyl) methylphosphonothioate
VX	<i>O</i> -ethyl <i>S</i> -[2-(diisopropylamino)ethyl] methylphosphonothioate

DISTRIBUTION LIST

The following organizations were provided with one Adobe portable document format (pdf) version of this report:

U.S. Army Edgewood Chemical
Biological Center (ECBC)
RDCB-DRC-C
ATTN: Bae, S.
Winemiller, M.
Berg, F.

Defense Threat Reduction Agency
DTRA-RD-CBD-T
ATTN: Ward, T.
J9-CBS
ATTN: Moore, E.

ECBC Technical Library
RDCB-DRB-BL
ATTN: Foppiano, S.
Stein, J.

Defense Technical Information Center
ATTN: DTIC OA

G-3 History Office
U.S. Army RDECOM
ATTN: Smart, J.

Department of Homeland Security
DHS-ORD-CSAC
ATTN: Famini, G.

Office of the Chief Counsel
AMSRD-CC
ATTN: Upchurch, V.

ECBC Rock Island
RDCB-DES
ATTN: Lee, K.

