# **Evolution of Tropical Cyclone Characteristics**

Patrick A. Harr Department of Meteorology Naval Postgraduate School Monterey, CA 93943-5114 Telephone : (831) 656-3787 FAX: (831) 656-3061 email: paharr@nps.navy.mil

Award No: N0001401WR20175

### LONG-TERM GOALS

The long-term goals are to understand how variabilities in the large-scale atmospheric environment and the internal tropical cyclone structure influence tropical cyclone track and intensity characteristics and define how these influences differ between developing, mature, and decaying tropical cyclones. During the initial stages of tropical cyclone development, structure and track characteristics can exhibit large variabilities that pose difficult forecast situations. Because decaying tropical cyclones often transition to fast-moving and rapidly-developing extratropical cyclones that may contain gale- or storm-force winds, there is a need to improve understanding and prediction of the extratropical transition phase of a decaying tropical cyclone. Therefore, a tropical cyclone throughout its life cycle has the potential for impacting many fleet components.

### **OBJECTIVES**

Recent research has concentrated on three primary objectives. Extended periods of increased and reduced tropical cyclone activity occur several times during a typical tropical cyclone season (Harr and Elsberry 1991;1995a,b). If reliable forecasts of extended periods of increased or reduced tropical cyclone activity could be made, maritime operations could be coordinated appropriately. Therefore, increased understanding of the primary factors responsible for initiation, maintenance, and decay of extended periods with tropical cyclone activity or inactivity is a primary objective of the current research.

When a tropical cyclone moves into the midlatitudes, extratropical transition (ET) often results in a fast-moving, rapidly-developing midlatitude cyclone that may contain gale-, storm-, or even typhoon-force winds capable of causing significant damage to coastal and maritime interests. Furthermore, these extreme conditions may occur during summer months when such conditions do not normally occur in association with extratropical cyclones. Based on previous studies of the structural changes of the tropical cyclone during ET (Harr and Elsberry 2000, Harr et al. 2000, Klein et al. 2000), the primary objective of the current research is to define how different types of interactions between the decaying tropical cyclone and midlatitude circulation influence the development of the extratropical cyclone.

Because of the increased need for tropical cyclone forecasts at intervals beyond 72 h, it is possible that a tropical cyclone may form and move a long distance during any given forecast sequence. Because numerical forecast guidance tends to be less accurate during the early stages of the tropical cyclone

<b>Report Documentation Page</b>					Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302 Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number							
1. REPORT DATE 30 SEP 2001		2. REPORT TYPE		3. DATES COVE 00-00-2001	RED L <b>to 00-00-2001</b>		
4. TITLE AND SUBTITLE		5a. CONTRACT	NUMBER				
Evolution of Tropi		5b. GRANT NUMBER					
			5c. PROGRAM ELEMENT NUMBER				
6. AUTHOR(S)			5d. PROJECT NUMBER				
		5e. TASK NUMBER					
					5f. WORK UNIT NUMBER		
	ZATION NAME(S) AND AE teorology,,Naval Pos	8. PERFORMING ORGANIZATION REPORT NUMBER					
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)			
		11. SPONSOR/MONITOR'S REPORT NUMBER(S)					
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited							
13. SUPPLEMENTARY NO	DTES						
the internal tropica define how these in initial stages of tro variabilities that pe fast-moving and ra there is a need to in	ls are to understand al cyclone structure ifluences differ betw pical cyclone develo ose difficult forecast pidly-developing ex mprove understandi herefore, a tropical tents.	influence tropical c veen developing, ma pment, structure ar situations. Because tratropical cyclones ing and prediction o	yclone track and ture, and decayin nd track character e decaying tropica s that may contain of the extratropica	intensity cha ag tropical cy ristics can ex al cyclones off a gale- or stor al transition p	racteristics and clones. During the hibit large ten transition to rm-force winds, phase of a decaying		
15. SUBJECT TERMS							
16. SECURITY CLASSIFIC	17. LIMITATION OF	18. NUMBER	19a. NAME OF				
a REPORT unclassified	ь abstract unclassified	c THIS PAGE unclassified	ABSTRACT Same as Report (SAR)	OF PAGES <b>7</b>	RESPONSIBLE PERSON		

Standard Form 298 (Rev. 8-98) Prescribed by ANSI Std Z39-18 life cycle, a new objective is to identify the capability of operational numerical models to identify circulations that will eventually develop into tropical cyclones.

# APPROACH

Tropical cyclone activity/inactivity may be related to mechanisms that act over a variety of space and time scales. It is hypothesized that the mechanisms responsible for clustering of tropical cyclone activity can be put into a framework of interactions between several modes of circulation variability. Low-level circulation and outgoing longwave radiation (OLR) characteristics are partitioned into various modes, then the interactions between modes are examined in relation to tropical cyclone activity/inactivity.

To investigate the structural evolution of a tropical cyclone to an extratropical cyclone, the approach has been to define extratropical transition to occur in two stages, transformation and re-intensification (Klein et al. 2000a). The impact of various factors such as the evolving internal tropical cyclone features (e.g., convective activity, inner-core structure), midlatitude circulation into which the tropical cyclone is moving, and relative roles of upper- and lower-level processes on the transition process. It is hypothesized that the development of the extratropical cyclone during re-intensification depends on the phasing of the poleward-moving tropical cyclone and a critical region in the midlatitude circulation into which the tropical cyclone is moving is examined by numerical experimentation in which only the phasing between the tropical cyclone and midlatitude circulations is altered. The movement of the tropical cyclone into the midlatitudes is then delayed or accelerated to examine the relative roles of tropical cyclone.

A catalog of forecast and analyzed circulations will be used to assess the ability of each operational numerical forecast model to predict tropical cyclone formation and initial intensification. The catalog will identify successful forecasts, forecast circulations that never verified, and analyzed circulations that were never forecast.

## WORK COMPLETED

A wavelet analysis was used to define the dominant modes of low-level circulation and OLR variability over the western North Pacific in a time-frequency reference. Although the wavelet analysis was applied over the entire data record, the zoom capability of the wavelet transform allows isolation of spectral power during specific time periods and over specific frequency ranges. Of interest in this study is the distribution of spectral power during the June-October period of typhoon activity over the western North Pacific. Partitions have been identified to represent an intraseasonal mode (30-60 days), a western North Pacific monsoon trough mode (10-25 days), and a synoptic mode (2-7 days). A singular value decomposition (SVD) analysis was used to describe the primary patterns of large-scale variability associated with the covariance between circulation features and OLR identified with predominant peaks in spectral power.

The sensitivity of the two-stage process of ET to the interaction between the tropical cyclone and the midlatitude circulation into which the tropical cyclone is moving was examined with the Coupled Ocean Atmosphere Mesoscale Prediction System (COAMPS) developed at the Naval Research Laboratory, Monterey. The sensitivity was examined by modifying the initial conditions of several COAMPS simulations. The original wind, temperature, height, and moisture fields associated with the tropical cyclone were removed and re-located to a specified location. The tropical cyclone-related values were placed into a new location via an iterative statistical interpolation procedure. The new location was chosen based on the desired effect, which may be to either delay or accelerate the motion of the tropical cyclone into the midlatitudes and thus alter the phasing with the midlatitude circulation (Klein et al. 2001).

To examine the capability of operational numerical models to identify circulations that may become tropical cyclones, an algorithm has been developed to detect and track circulations throughout the forecast sequence. A matching technique was used to associate predicted circulations with subsequent analyzed circulations.

#### RESULTS

Based on the SVD analysis, dominant coupled modes of variability between low-level circulation and OLR have been identified for the three time modes defined by the wavelet analysis. Over intraseasonal scales, a global-scale circulation/OLR coupled mode has been associated with the slow eastward propagation of the Madden Julian Oscillation (MJO). Analysis of the SVD coefficients defines considerable variability in the MJO structural characteristics that projects onto circulation modes that act on longer time scales (Chang et al. 2001) and shorter time scales. Over the shorter time scales, the variable MJO-mode amplitude is associated with large variability in the modulation of synoptic-scale features by the intraseasonal events. Therefore, the key to defining potential initiation of intraseasonal periods of tropical cyclone activity or inactivity is identification of the variability in the principal features of the MJO that influence the synoptic scale directly. Over the monsoon trough time scale, the primary coupled circulation/OLR modes define large-scale circulations confined to the western North Pacific. The structure of this mode is related to the northwestward propagation of Rossby waves from enhanced convection over the equatorial western Pacific. Coupled low-level circulation/OLR modes over the synoptic scales define a combination of equatorial waves. Over the central Pacific, a mixed Rossby-gravity (MRG) wave signal is identified. This signal becomes modified over the tropical western North Pacific to be representative of northwestward-moving synoptic-scale waves that have been identified in previous studies (e.g., Chang et al. 1996). Because of the variability associated with each coupled mode, and variability associated with combinations of large-scale and mesoscale factors that influence tropical cyclone activity, the relationship between the various modes and tropical cyclone activity is very complex. However, the unique partitioning of the circulation/OLR modes enables assessment of the amount of variability in tropical cyclone activity that may be explained by each coupled mode.

One aspect of the two-stage process associated with ET is that the transformation stage is rather independent of the tropical cyclone and large-scale circulation characteristics. However, the reintensification stage is highly dependent on the phasing between the decaying tropical cyclone and the midlatitude circulation into which it is moving. The contributions of the midlatitude circulation and by the tropical cyclone remnants to the outcome of the re-intensification stage of ET are depicted in a matrix (Fig. 1). The midlatitude contribution is based on the degree to which it is favorable for extratropical cyclogenesis even without the tropical cyclone (Klein et al. 2001). The characterization of the contribution from the decaying tropical cyclone to the re-intensification stage of ET is based on its interaction with the midlatitude circulation. An optimal interaction results in a dynamic coupling of the transformed tropical cyclone with the mid- and upper-tropospheric components of the midlatitude circulation plus the low-level baroclinity such that a deep re-intensification (minimum sea-level pressure less than 980 mb) results. For cases in which the interactions between the transformed tropical cyclone contribution is minor. If the transformed tropical cyclone contribution is described as none. All contributions are defined from the sensitivity analysis based on modification of the initial conditions in a COAMPS simulation such that the decaying tropical cyclone characteristics are changed to alter the interaction with the midlatitude circulation. The results can be conveyed (Fig. 1) as a shift in the tropical cyclone contribution within the same category of midlatitude contribution.

Re-intensification Stage of ET		Midlatitude Contributions				
		Unfavorable	Neutral	Favorable		
utions	Significant	Little or	Deep and/or Rapid Re- intensification			
<b>Contributions</b>	Minor		Moderate			
TC	None	Decayers		Re- intensification		

Fig. 1: Characterization of the midlatitude circulation and tropical cyclone contributions to the re-intensification stage of ET depicted as a matrix, where the combinations of these contributions may produce three outcomes: (i) deep and/or rapid re-intensification (upper right); (ii) little to moderate re-intensification (within the shaded cells); or (iii) decay of the tropical cyclone without completing ET (lower left).

The algorithm that allows forecast and analyzed circulations to be cataloged, which has just been completed, will be applied to several operational models to develop a knowledge data base that defines the capability of each model to accurately predict tropical cyclone formation at all forecast ranges.

## **IMPACT/APPLICATIONS**

Identification of the interactions between various modes of tropical circulation variability will lead to a statistical forecast scheme of extended periods of tropical cyclone activity/inactivity. If reliable forecasts of extended periods of inactivity (i.e., at least 20 days with no tropical cyclones) could be made, maritime operations could be coordinated to take advantage of the period of reduced threat from tropical cyclones.

Knowledge of the degree of the midlatitude and tropical cyclone contributions during ET will provide increased forecaster awareness of the potential re-intensification of the extratropical cyclone that results from the completed ET process.

Knowledge of the potential for each operational numerical forecast model for prediction of tropical cyclone formation will lead to a consensus technique for identifying the probability of tropical cyclone formation.

### TRANSITIONS

The two-stage model of ET has been incorporated into official training material at the Joint Typhoon Warning Center.

### SUMMARY

Results from research of ET conducted under ONR sponsorship at the Naval Postgraduate School has triggered a major sense of awareness of the need for increased understanding of this process (Jones et al. 2001). The primary structural characteristics associated with ET have been identified by Harr and Elsberry (2000), Harr et al. (2000), and Klein et al. (2000,2001). Harr and Elsberry (2001) identified how these characteristics are associated with key weather hazards such as extreme amounts of precipitation. During the next year, results from the examination of the interaction between modes of tropical circulation variability will provide a sound forecast system for assessment of tropical cyclone activity that may be expected over ranges between 10-25 days. Finally, the assessment of the capability of each operation numerical model to predict tropical cyclone formation will provide operational forecasters with information to efficiently assess model guidance.

### REFERENCES

Chang, C.-P., J. M. Chen, P. A. Harr, and L. E. Carr, 1996: Northwestward-propagating wave patterns over the tropical western North Pacific during summer. *Mon. Wea. Rev.*, **124**, 2245-2266.

Chang, C.–P., P. A. Harr, and J. Ju, 2001: Possible roles of Atlantic circulations on the weakening Indian monsoon rainfall-ENSO relationship. *J. Climate*, **14**, 2376-2380.

Harr, P. A., and R. L. Elsberry, 1991: Tropical cyclone track characteristics as a function of large-scale circulation anomalies. *Mon. Wea. Rev.*, **119**, 1448-1468.

Harr, P. A., and R. L. Elsberry, 1995a: Large-scale circulation variability over the tropical western North Pacific. Part I: Spatial patterns and tropical cyclone characteristics. *Mon. Wea. Rev.*, **123**, 1225-1246.

Harr, P. A., and R. L. Elsberry, 1995b: Large-scale circulation variability over the tropical western North Pacific. Part II: Persistence and transition characteristics. *Mon. Wea. Rev.*, **123**, 1247-1268.

Harr, P. A., and R. L. Elsberry, 2000a: Extratropical transition of tropical cyclones over the western North Pacific. Part I: Evolution of structural characteristics during the transition process. *Mon. Wea. Rev.*, **128**, 2613-2633.

Harr, P. A., R. L. Elsberry, and T. F. Hogan, 2000: Extratropical transition of tropical cyclones over the western North Pacific. Part II: The impact of midlatitude circulation characteristics. *Mon. Wea. Rev.*, **128**, 2634-2653.

Harr, P. A., and R. L. Elsberry, 2001: An overview of heavy precipitation associated with the extratropical transition of tropical cyclones. *Preprints, Symposium on Precipitation Extremes: Prediction, Impacts and Responses.* Amer. Meteor. Soc., Boston, MA 02108, 288-292.

Jones, S. C., P. A. Harr, J. Abraham, L. F. Bosart, P. J Bowyer, B. N. Hanstrum, M. R. Sinclair, R. K. Smith, and C. Thorncroft, 2001: The extratropical transition of tropical cyclones: Forecast challenges, current understanding and future directions. *Wea. and Forecasting*, In revision.

Klein, P. M., P. A. Harr, and R. L. Elsberry, 2000a: Extratropical transition of western North Pacific tropical cyclones: An overview and conceptual model of the transformation stage. *Wea. Forecasting*, **15**, 373-395.

Klein, P. M., P. A. Harr, and R. L. Elsberry, 2001: Extratropical transition of western North Pacific tropical cyclones: Midlatitude and tropical cyclone contributions to re-intensification. *Mon. Wea. Rev.* In revision.

## PUBLICATIONS

Chang, C.–P., P. A. Harr, and J. Ju, 2001: Possible roles of Atlantic circulations on the weakening Indian monsoon rainfall-ENSO relationship. *J. Climate*, **14**, 2376-2380.

Harr, P. A., and R. L. Elsberry, 2001: An overview of heavy precipitation associated with the extratropical transition of tropical cyclones. *Preprints, Symposium on Precipitation Extremes: Prediction, Impacts and Responses.* Amer. Meteor. Soc., Boston, MA 02108, 288-292.

Jones, S. C., P. A. Harr, J. Abraham, L. F. Bosart, P. J Bowyer, B. N. Hanstrum, M. R. Sinclair, R. K. Smith, and C. Thorncroft, 2001: The extratropical transition of tropical cyclones: Forecast challenges, current understanding and future directions. *Wea. and Forecasting*, In revision

Klein, P. M., P. A. Harr, and R. L. Elsberry, 2001: Extratropical transition of western North Pacific tropical cyclones: Midlatitude and tropical cyclone contributions to re-intensification. *Mon. Wea. Rev.* In revision.