

Investigation of Processes and Factors Regulating the Generation, Maintenance and Breakdown of Bioluminescent Thin Layers

Dr. Edith Widder
Harbor Branch Oceanographic Institution
phone (561) 465-2400 x315 fax (561) 468-0757 email widder@hboi.edu

Grant Number: N00014-00-1-0176
<http://www.hboi.edu/marine/biolum1.html>

LONG-TERM GOALS

My research objectives involve determining how low light phenomena, both bioluminescence and solar radiation below 200 meters, influence the distribution and behavior of marine organisms.

OBJECTIVES

To investigate the phenomenon of bioluminescent thin layers discovered in Wilkinson Basin, Gulf of Maine (Widder et al., 1999). These thin layers (0.5 m) are composed of aggregations of the bioluminescent copepod *Metridia lucens* at density discontinuities in the water column. Three testable hypotheses were formulated to account for these aggregations: 1.) these copepods may enhance the repellent capacity of their bioluminescence by forming aggregations 2.) aggregation into thin layers may be an avoidance response to current shears or turbulent regions 3.) the copepods may have been orienting to some chemical cue or food resource such as marine snow that has been found at density discontinuities in the water column (MacIntyre et al., 1995).

APPROACH

Using a computer algorithm, which we have recently developed for 3D reconstruction and statistical analysis of spatial point patterns of identified bioluminescent displays, we measured the median nearest neighbor distances of the bioluminescent copepods in these layers (Widder and Johnsen, 2000). Since this analysis indicated that the distance between individuals was nearly 10 times greater than the maximum perceptive distance of a copepod (Haury and Yamazaki 1995) we believe the first hypothesis, requiring as it does behavioral interaction between individuals, is the least probable of the three. To test the other two hypotheses and to further examine the factors that regulate the generation, maintenance and breakdown of these bioluminescent thin layers we conducted a field study in Wilkinson Basin in collaboration with Dr. Joe Katz of Johns Hopkins University. Thin layers were located using the High-Intake-Defined-Excitation-Bathypotometer (Widder et al., 1993). Immediately following HIDEX-BP profiles the *Johnson-Sea-Link* submersible was launched with Dr. Katz's submersible holographic camera (Katz et al., 1999) mounted on the upper work platform (Figure 1). Thin layers were located using real-time sensor feedback from intensified video recordings of stimulated bioluminescence (Widder et al., 1989). Identification of organisms responsible for the bioluminescence was based on the spatial and temporal properties of the recorded displays. These displays were compared to our existing database of identified displays from this

Report Documentation Page

Form Approved
OMB No. 0704-0188


Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE 30 SEP 2001		2. REPORT TYPE		3. DATES COVERED 00-00-2001 to 00-00-2001	
4. TITLE AND SUBTITLE Investigation of Processes and Factors Regulating the Generation, Maintenance and Breakdown of Bioluminescent Thin Layers				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Harbor Branch Oceanographic Institution,,Fort Pierce,FL				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT My research objectives involve determining how low light phenomena, both bioluminescence and solar radiation below 200 meters, influence the distribution and behavior of marine organisms.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a REPORT unclassified	b ABSTRACT unclassified	c THIS PAGE unclassified			

region (Widder et al., 1992; Widder, 1997). Both single and double exposure holograms (the latter for velocity measurements) were collected during profiles through the layers and during transects in the layers. In order to analyze the content of the holograms, each of which contain roughly 35 GB of unprocessed image data, a computerized scanning procedure capable of automatically selecting only in focus particles was required. Developing this procedure has been the emphasis in the second year of this investigation.

WORK COMPLETED

Field work was carried out in the first year of this study. In order to mount the 1500 lb. holographic camera on the *Johnson-Sea-Link* submersible all other sampling gear including the robotic arm and the entire lower work platform had to be removed and a support frame had to be designed and fabricated that could support the weight of the camera in air. The electronics and control systems were mounted in front of the JSL and the optical system was mounted completely above it in order to minimize the effect of the submersible on the flow and plankton in the same volume (Fig. 1). During the field investigation in Wilkinson Basin, Gulf of Maine (42°20'N; 69°47'W) which took place June 29 – July 6, 2000 holograms were collected by locating a thin layer by its bioluminescence and then trimming the sub out just below the layer after first moving out from under any water disturbed by the submersible. A minimum amount of positive trim was then applied allowing the sub to drift up very slowly through the layer while a series of holograms were collected in rapid succession. The cylindrical sample volume for each hologram was 6.3 cm diameter and 28 cm long.




1. Johnson-Sea-Link submersible with holocam mounted on upper work platform, port side. Sample volume is between upper most portion of blue fins.

After the deployment, and during the second year the Johns Hopkins group focused on developing automatic processing procedures for detecting particles in the recorded hologram reconstructions. The system, just recently completed, has been optimized to detect and locate the numerous small (<50 μm) particles (e.g. dinoflagellates), which tend to have a very low signal to noise ratio, without reducing the detection of the larger particles such as marine snow and fecal pellets. A paper to be submitted to Applied Optics, detailing the procedure is in preparation.


RESULTS

During the field investigation a total of 47 HIDEX-BP casts were collected. Bioluminescent thin layers, similar to those recorded in the 1992 investigation, were present. Oblique bioluminescence video transects proved very effective in locating thin layers from the submersible. Over 600 single and double exposure holograms were recorded in the upper 100 m of the water column. These are the clearest and sharpest holograms ever recorded with this submersible system (Figure 2).



2. (a) *Metridia lucens* oriented toward (b) marine snow; c) anterior view of a neighboring small copepod. The scale in (a) is 1 mm and the scale in (b,c) is 100 μm .

During the second year, computer programs designed to identify and locate particles within the holograms and to generate 3D vector arrays of instantaneous velocity distributions were perfected (Figure 3).



3. *Calanus finmarchicus* in a double exposure hologram. *The velocities represented have the mean component subtracted i.e. the relative flow of the sub moving upward (0.1m/s) and forward. The copepod is not moving relative to its surroundings, which have an RMS velocity over the sample volume of approximately 3.4 mm/s. The inhomogeneities in the background, are due to refractive index fluctuations within the sample volume.*

Because of low signal to noise ratios inherent to in-line holographic recording and because of non-uniform exposures, it was necessary to devise new optical and digital methods to improve particle detection. These techniques increase the number of particles detected per hologram (~ 1 liter) from what was initially a few hundred per liter to what is now several thousand. With new processing hardware and increased scanning efficiency, it now takes 4 hours to scan and process a hologram compared to 24 hours earlier this year. 33 holograms were scanned with the old method. 20 holograms (each 35 GB) have since been scanned with the new method. Analysis of the NND from these scans is in progress.

Plankton samples, that were pumped from depths where bioluminescence potential was high based on both HIDE-X-BP profiles and video transect data, were analyzed. These samples were completely dominated by copepods with three species present in significant numbers. These were *Metridia lucens* (almost entirely females), *Calanus finmarchicus* and *Oithona* spp. There were also very large numbers of exoskeletons (copepod molts) in the samples. Also present was the cydippid ctenophore *Euplokamis* sp. Although destroyed by the pump, these brilliantly luminescent ctenophores were readily apparent in the video transects. The presence of this ctenophore in high abundance at relatively shallow depths (~40 m) was noteworthy because in past investigations in these waters it has always been found much deeper (> 200 m) (Widder et al., 1992; Widder 1997; Widder et al., 1999).

IMPACT/APPLICATIONS

From a strategic standpoint the existence of thin layers of intense bioluminescence could have a detrimental impact on covert naval operations and objectives. Because, these thin layers would not be detected by standard low flow (<1 l/s) bathyphotometer systems they would not be identified by real-time environmental now-casts and forecasts. Therefore, while the average bioluminescence measured by a low-flow bathyphotometer system might indicate an average bioluminescence that was deemed acceptable for a nighttime stealth mission, that mission could be seriously compromised if a boat, SDV or swimmer encountered one of these bioluminescent hot zone.

From a scientific standpoint the existence of food-rich thin layers provides a possible explanation for why measured average *in situ* food concentrations appear to be inadequate for the daily metabolic requirements of marine grazers (Mullin and Brooks 1976; Dagg 1991; Gifford 1993; Batchelder and Williams 1995; Cowles and Fessenden 1995). Although the possibility has been raised that this disparity could be accounted for by the existence of food-rich micropatches (e.g. Batchelder and Williams 1995), this solution depends on how much energy grazers must expend to locate such patches (Haury et al. 1978). Locating food-rich thin layers, especially for vertical migrators, is less problematic since the search strategy can be reduced from three dimensions to one. A grazer moving up or down through the water column has a much higher probability of encountering food-rich micropatches if those patches are spread out into very thin layers. If such thin layers are in fact a common feature of the marine environment then their existence will have a profound impact on such critical aspects of population dynamics as species-habitat associations and encounter probabilities between predators and prey.

TRANSITIONS

These data will be made readily available to NAVOCEANO and other Naval investigators as well as to the oceanographic community through publication in the open literature.

RELATED PROJECTS

During the field investigation, tests were conducted on an expendable bathyphotometer system that is currently being developed with support from ONR (see report: "A Compact Bathyphotometer" by Fucile, Widder and Brink).

REFERENCES

Batchelder HP, Williams R (1995) Individual-based modeling of the population dynamics of *Metridia lucens* in the North Atlantic. ICES J mar Sci 52: 469-482

Cowles TJ, Fessenden LM (1995) Copepod grazing and fine-scale distribution patterns during the Marine Light-Mixed Layers experiment. J Geophys Res 100: 6677-6686

Dagg MJ (1991) *Neocalanus plumchrus* (Marukawa): life in the nutritionally dilute subarctic Pacific Ocean and the phytoplankton-rich Bering Sea. In Proceedings of the Fourth International conference on Copepoda, Bull. Plank. Soc. Jap., special volume: 217-225

Gifford DJ (1993) Protozoa in the diets of *Neocalanus* spp. in the oceanic subarctic Pacific Ocean.

Prog Oceanography 32: 223-237

Haury LR, McGowan JA, Wiebe PH (1978) Patterns and processes in the time-space scales of plankton distributions. In: Steele JH (ed) Spatial patterns in plankton communities. Plenum, New York, pp 277-327

Haury LR, Yamazaki H (1995) The dichotomy of scales in the perception and aggregation behavior of zooplankton. J Plank Res 17:191-197

Katz, J., Donaghay, P.L., Zhang, J., King, S. and Russell, K. (1999) Submersible holocamera for detection of particle characteristics and motions in the ocean. Deep Sea Research I 46: 1455-1481.

MacIntyre S, Alldredge AL, Gotschalk CC (1995) Accumulation of marine snow at density discontinuities in the water column. Limnol Oceanogr 40: 449-468

Malkiel, E, Alquaddoomi, O, Katz, J (1999) Measurements of Plankton Distribution Using Submersible Holography. Measurement Science and Technology, 10: 1153-1161

Mullin MM, Brooks EH (1976) Some consequences of distributional heterogeneity of phytoplankton and zooplankton. Limnol Oceanogr 21(6): 784-796

Tao, B, Katz, J and Meneveau, C (1999) Application of HPIV data of turbulent duct flow for turbulence modeling. Proc 3rd ASME/JSME Joint Fluids Conf. July 18-22, 1999, San Francisco, CA; paper FEDSM99-7281

Widder, EA (1997) *In situ* video recordings of bioluminescence in the ostracod, *Conchoecia elegans*, and co-occurring bioluminescent zooplankton in the Gulf of Maine. In: Proceedings of the 9th International Symposium on Bioluminescence and Chemiluminescence. Eds. JW Hastings, LJ Kricka and PE Stanley. John Wiley & Sons Ltd, Sussex, UK. pp. 159-164.

Widder, EA, Bernstein S, Bracher D, Case JF, Reisenbichler KR, Torres JJ and Robison BH. (1989) Bioluminescence in Monterey Submarine Canyon: image analysis of video recordings from a midwater submersible. Mar. Biol. 100:541-551.

Widder, EA, Case JF, Bernstein SA, MacIntyre S, Lowenstine MR, Bowlby MR, and Cook. DP (1993) A new large volume bioluminescence bathyphotometer with defined turbulence excitation. Deep Sea Res. 40(3): 607-627.

Widder, EA, Greene CH and Youngbluth MJ. (1992) Bioluminescence of sound-scattering layers in the Gulf of Maine. J. Plank. Res. 14(11): 1607-1624.

Widder, E.A., S. Johnsen, S. A. Bernstein, J. F. Case, D. J. Neilson. (1999) Thin layers of bioluminescent copepods found at density discontinuities in the water column. Mar. Biol. 134: 429-437.

Widder, E.A and S. Johnsen (2000) 3D spatial point patterns of bioluminescent plankton: A map of the "minefield" J. Plank. Res. 22(3): 409-420.

PUBLICATIONS

- Widder, E.A. (1999) Bioluminescence. In: "Adaptive Mechanisms in the Ecology of Vision." Edited by: S.N. Archer, M.B.A. Djamgoz, E. Loew, J.C. Partridge & S. Vallergera. Kluwer Academic Publishers, Dordrecht, the Netherlands. Pp 555-581.
- Kocak, D.M., N.D.V. Lobo, and E.A. Widder. (1999) Computer vision techniques for quantifying, tracking and identifying Bioluminescent plankton. *IEEE J. Oceanic Engineering* 24(1):81-95.
- Johnsen, S., E.J. Balsler and E.A. Widder. (1999) Modified suckers as light organs in a deep-sea octopod. *Nature* 398:113-114.
- Johnsen, S., E.J. Balsler, E.C. Fisher, and E.A. Widder (1999) Bioluminescence in the deep-sea cirrate octopod *Stauroteuthis syrtensis* Verrill (Mollusca: Cephalopoda) *Biol. Bull.* 197: 26-39
- Widder, E.A., S. Johnsen, S. A. Bernstein, J. F. Case, D. J. Neilson. (1999) Thin layers of bioluminescent copepods found at density discontinuities in the water column. *Mar. Biol.* 134: 429-437.
- Johnsen, S. and E.A. Widder. (1999) The physical basis of transparency in biological tissue: Ultrastructure and the minimization of light scattering. *J. Theor. Biol.* 199: 181-198
- Frank, T.M. and E.A. Widder. (1999) A comparative study of the spectral sensitivities of mesopelagic crustaceans. *J. Comp. Physiol.* 185(3): 255-265.
- Malkiel, E, Alquaddoomi, O, Katz, J (1999) Measurements of Plankton Distribution Using Submersible Holography. *Measurement Science and Technology*, 10: 1153-1161
- Widder, E.A and S. Johnsen (2000) 3D spatial point patterns of bioluminescent plankton: A map of the "minefield" *J. Plank. Res.* 22(3): 409-420.
- Herring, P.J. and E.A. Widder (2001) Plankton: Light production in marine organisms, mechanisms and functions. *Encyclopaedia of Ocean Sciences*. Academic Press.
- Widder, E.A. (2000) Bioluminescence in octopods. *Yearbook of Science and Technology*. McGraw-Hill 2001:52-55.
- Widder, E.A. (2000) Bioluminescence in octopods. **Yearbook of Science and Technology**. McGraw-Hill 2001: 52-55.
- Herring, P.J. and E.A. Widder (2001) Plankton: Light production in marine organisms, mechanisms and functions. **Encyclopaedia of Ocean Sciences**. Academic Press.
- Widder, E.A. and T.M. Frank (2001) The Speed of an Isolume: A Shrimp's Eye View. **Mar. Biol.** 138:669-677.
- Johnsen, S. and E.A. Widder (2001) Ultraviolet absorption in transparent zooplankton and its implications for depth distribution and visual predation. **Mar. Biol.** 138:717-730.

Vecchione, M., C.F.E. Roper, E.A. Widder, and T.M. Frank (in press) In-situ observations of three species of large-finned deep-sea squids. **Bull. Mar. Sci.**

Widder, E.A. (in press) Bioluminescence and the pelagic visual environment. **Mar. Fresh. Behav. Physiol.** Invited review.

Frank, T.M. and E.A. Widder (in press) Effects of a decrease in downwelling irradiance on the daytime vertical distribution patterns of mesopelagic zooplankton and micronekton. **Mar. Biol.**

Daumer, K.A. and E.A. Widder (submitted) The effects of two different bioluminescent dinoflagellates (*Lingulodinium polyedrum* and *Pyrocystis noctiluca*) on the susceptibility of zooplankton to teleost predation

Widder, E.A. (submitted) Bioluminescence. **Bioscience Explained** Invited review.