

# **Global Mapping of Near-inertial and Tidal Internal Wave Propagation**

Matthew H. Alford  
Applied Physics Laboratory  
1013 NE 40<sup>th</sup> Street  
Seattle, WA 98105

phone: (206) 221-3257 fax: (206) 543-6785 email: [malford@apl.washington.edu](mailto:malford@apl.washington.edu)

Grant Number: N00014-02-1-0526  
<http://faculty.washington.edu/malford/>

## **LONG-TERM GOALS**

The long-term goal is to understand the magnitude and distribution of internal-wave propagation and internal-wave-induced mixing.

## **OBJECTIVES**

- To use detect energy-flux associated with low-mode near-inertial and tidal propagation in moored current meter data at as many locations as possible.
- To determine the degree to which energy-flux divergences (and therefore sites of enhanced dissipation) can be determined from this analysis.
- To compute time-averaged spatial maps of near-inertial, internal-tidal and mesoscale energy from TOPEX/ POSEIDON altimetry, drifters, current meters and ADCP tracks.
- To determine, within the context of Peter Muller's Internal Wave Action Model (IWAM, also supported by ONR), the extent to which these findings are consistent with existing internal-wave propagation theories.
- To use large-scale and regional atmospheric models to examine the high-latitude and small-scale reliability, interannual variability, and convergences/divergences in the wind energy flux maps computed by Alford (2001).

## **APPROACH**

To accomplish the goal stated above, one could measure microstructure at every location on the planet (impossible). The alternate approach taken here is to map the sources of internal waves already begun by Egbert and Ray (2000) for tidal input, and by Alford (2001) and Toshi (2002) for near-inertial input, and to follow their subsequent propagation. Divergences in the internal-wave energy flux field must then correspond to energy sinks, i.e., regions of enhanced mixing.

I have begun by computing the energy flux in the near-inertial and tidal bands using historical moored records of velocity and temperature. Hundreds of records have been collected over the years. Kunze et al (2002) showed that energy-flux could be computed from full-depth records of velocity and

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>30 SEP 2002</b>		2. REPORT TYPE		3. DATES COVERED <b>00-00-2002 to 00-00-2002</b>	
4. TITLE AND SUBTITLE <b>Global Mapping of Near-inertial and Tidal Internal Wave Propagation</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Applied Physics Laboratory,,1013 NE 40th Street,,Seattle,,WA, 98105</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT <b>The long-term goal is to understand the magnitude and distribution of internal-wave propagation and internal-wave-induced mixing.</b>					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>Same as Report (SAR)</b>	18. NUMBER OF PAGES <b>5</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

isopycnal displacement. The approach taken here is first to band-pass moored records of velocity and temperature to either near-inertial or tidal frequencies. Isopycnal displacement is then estimated from temperature records via  $\eta = T' / (\delta T / \delta z)$ . Time series of velocity and displacement for the first two baroclinic modes are estimated from the moored records using a least-squares fit. (Experience is showing that not all mooring geometries are equally capable of resolving the first two modes; careful sensitivity analysis will be required to address these and other sensitivity issues.) The pressure anomaly,  $p'$ , is then estimated from the full-depth modal profiles of  $\eta$  according to the methods of Kunze et al (2002), and the vector energy-flux computed from  $\mathbf{F} = \langle \mathbf{u}' p' \rangle$ , where  $\langle \rangle$  indicate averaging over a wave period.

## WORK COMPLETED

Funding for this project began May 2002. Since that time, Ph.D. student Maya Whitmont has arrived at APL/UW and has begun work analyzing historical moored records from the Oregon State Buoy group. She has examined hundreds of records for quality, and processed them into uniform format. Sixty-five records were selected for the initial flux calculation presented here. Then, I used these records to compute the fluxes in the lowest two modes in the inertial and  $M_2$  bands. A paper describing the initial findings for the near-inertial band, summarized in the next section, has been submitted to Nature (Alford 2002).


## RESULTS

The results of the first 65 moorings in the near-inertial band are shown in Figure 1. Several observations are immediately apparent:

1. Predominant propagation is toward the equator, consistent with generation by wind at the local inertial frequency and subsequent propagation to regions of lower  $f$ . This has profound implications for our understanding of the global internal-wave field, and is in contrast with the model of Fu (1981), which attempted to explain the inertial peak by *poleward* propagation of super-inertial internal waves.
2. The magnitudes are of order  $1 \text{ kWm}^{-1}$ , the same order of magnitude as measured tidal fluxes (e.g. Ray and Cartwright, 2001). Therefore, near-inertial waves appear as important as the tides in transporting energy horizontally.
3. The fluxes are of the correct order of magnitude to account for a substantial percentage of near-inertial input (shown in color). Therefore, assuming that all energy input is dissipated at the source is not justified. Propagation is, in general, away from source regions, as expected.

## IMPACT/APPLICATIONS

The observed tendency for near-inertial energy flux to be directed toward the equator is extremely important, confirming the idea of wind as a major source of internal waves. The observation of substantial fluxes in the near-inertial band is a major step toward understanding internal-wave-generated mixing. As the database of observations grows, divergences in the energy flux maps should point to regions of high dissipation, guiding future observational efforts.



**Figure 1:** (Color) Annual-mean energy-flux from the wind to near-inertial mixed-layer motions from Alford (2001). The color scale is logarithmic and is indicated at upper left. Depth-average near-inertial energy-flux vectors for modes 1 and 2 are plotted from 65 historical moored records. The length of each arrow is logarithmic, with references indicated at upper left. The few instances of poleward propagation are plotted in white. The majority of records show equatorward energy propagation, consistent with generation by midlatitude travelling storms and subsequent beta-plane propagation.

## TRANSITIONS

Maya Whitmont's Ph.D. thesis will continue these analyses, attempting to form a global view of the energy fluxes to, through and out of the internal-wave field.

Peter Muller is enthusiastic about the incorporation of these data into his ONR-funded Internal Wave Action Model (IWAM). As the reliability of this model grows, it may be useful in filling in data gaps evident in Figure 1.

## RELATED PROJECTS

None to report.

## REFERENCES

Alford, M. H., Internal swell generation: The spatial distribution of energy flux from the wind to mixed-layer near-inertial motions, *J. Phys. Oceanogr.*, 31, 2359–2368, 2001.

Dushaw, B., B. Cuornelle, P. F. Worcester, B. Howe, and D. Luther, Barotropic and baroclinic tides in the Central North Pacific Ocean determined from long-range reciprocal acoustic transmissions, *J. Phys. Oceanogr.*, 25, 631–647, 1995.

Egbert, G. D., and R. D. Ray, Significant dissipation of tidal energy in the deep ocean inferred from satellite altimeter data, *Nature*, 45 , 775–778, 2000.

Fu, L.-L., Observations and models of inertial waves in the deep ocean, *Rev. Geophys. Space Phys.*, 19, 141–170, 1981.

Garrett, C., What is the “near-inertial” band and why is it different from the rest of the internal wave spectrum?, *J. Phys. Oceanogr.*, 31 , 962–971, 2001.

Kunze, E., L. Rosenfield, G. Carter, and M. C. Gregg, Internal waves in Monterey submarine canyon, 2002, *J. Phys. Oceanogr.*, in press.

Ray, R. D., and D. E. Cartwright, Estimates of internal tide energy fluxes from topex/poseidon altimetry: Central north pacific, *Geophys. Res. Lett.*, 28, 1259–1262, 2001.

Watanabe, M., and T. Hibiya, Global estimates of the wind-induced energy flux to inertial motions in the surface mixed layer, *J. Geophys. Res.*, 29 , 80.1–80.4, 2002.

## **PUBLICATIONS**

Alford, M. H., Internal swell: near-inertial energy flux measured from historical moored records, *Nature*, submitted, 2002.