

Remote-Sensing Retrieval of Oceanic Inherent Optical Properties by Inversion of the Radiative Transfer Equation

Curtis D. Mobley
Sequoia Scientific, Inc.
Westpark Technical Center
15317 NE 90th Street
Redmond, WA 98052
phone: 425-867-2464 x 109 fax: 425-867-5505 email: mobley@sequoiasci.com

Award Number: N0001401M0268

http://www.onr.navy.mil/sci_tech/ocean/322_processes/prog_op.htm

LONG-TERM GOAL

The overall goal of this work was to develop and evaluate a new method of inverting remotely sensed ocean color data to recover the “big three” water inherent optical properties (IOPs), namely the phytoplankton absorption coefficient, the CDOM+detritus absorption coefficient, and the total constituent backscattering coefficient.

OBJECTIVES

Semianalytic radiance models [Gordon et al., 1988; Morel and Gentili, 1996] can be readily inverted by linear matrix methods [Hoge et al., 1999a,b] to provide oceanic inherent optical properties (IOPs). Such inversions are well conditioned [Hoge and Lyon, 1996] and promise a powerful method of simultaneously retrieving constituent absorption and backscattering coefficients in the upper surface layer of the world's oceans using satellite data. However, semianalytic radiance models do not provide an exact framework to account for all possible environmental and viewing conditions [Weidemann et al., 1995].

The radiative transfer equation (RTE) can provide exact inverse solutions, but the RTE is not easily inverted for many remote sensing situations [Zaneveld, 1995]. We therefore investigated the inversion of a specific form of the RTE, namely a modified version of the shape-factor formulation of Zaneveld [1995]. Some of the motivation for this work comes from the distinct need for highly accurate methods to retrieve the absorption coefficients of the chlorophyll accessory pigment phycoerythrin [Hoge et al., 1999b]. To this end the absorption coefficients of chlorophyll and CDOM must be accurately retrieved, otherwise weaker absorbing constituents (such as phycoerythrin) will be obscured.

APPROACH

Given measured water-leaving radiances or remote-sensing reflectances, it is not possible to invert the radiative transfer equation (RTE) to obtain the water column absorption and backscatter coefficients unless additional assumptions are made, or additional information is added, to constrain the inversion. Our approach to constraining the inversion is to model selected unknowns—the so called shape

Report Documentation Page

Form Approved
OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE 30 SEP 2003		2. REPORT TYPE		3. DATES COVERED 00-00-2003 to 00-00-2003	
4. TITLE AND SUBTITLE Remote-Sensing Retrieval of Oceanic Inherent Optical Properties by Inversion of the Radiative Transfer Equation				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Sequoia Scientific, Inc., Westpark Technical Center,,15317 NE 90th Street,Redmond,,WA, 98052				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT The overall goal of this work was to develop and evaluate a new method of inverting remotely sensed ocean color data to recover the ???big three??? water inherent optical properties (IOPs), namely the phytoplankton absorption coefficient, the CDOM+detritus absorption coefficient, and the total constituent backscattering coefficient.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

factors and related quantities—as well as possible in terms of known quantities, and then treat the shape factors as known in the inversion process. This reduces the number of unknowns in the inversion, at the expense of introducing uncertainties via the imperfectly modeled shape factors. Clearly, the ultimate success of the inversion depends on how well the shape factors can be modeled.

If IOPs such as the absorption and backscatter coefficients are recovered as part of an iterative inversion algorithm, then the recovered IOPs can be used to model the shape factors because both the recovered IOPs and the IOP-dependent shape factors are iteratively improved during the course of the RTE inversion. In the present modeling, we considered the absorption coefficient a , the beam attenuation coefficient c , and the backscatter coefficient b_b to be the recovered (i.e., known) IOPs.

WORK COMPLETED

A database of Hydrolight-generated synthetic data was generated from a large number of Hydrolight runs using different water IOPs, solar angles, viewing directions, wavelengths, etc. (Hydrolight is described at www.hydrolight.info; see also Mobley, 1994 and Mobley and Sundman, 2001a,b.) Each record in the database (184,800 in all) contained the input IOPs, solar angle, etc, as well as the corresponding shape factors and the water-leaving radiance. This database was used in conjunction with guidance from radiative transfer theory to develop best-fit models for the generally unknown shape factors in terms of the known quantities.

A paper (Mobley et al., 2002) describing the background theory and the shape factor models has been submitted. A companion paper showing the application of the shape factor inversion methodology to both synthetic and actual data is in preparation (Lyon et al., 2002).

No field data were acquired in this work. Therefore no data have been submitted to a national data archive.

RESULTS

The IOP-dependent shape factor models can predict the shape factors with accuracies ranging typically from two to 20%. Error analysis suggests that IOP retrievals are primarily influenced by uncertainties in the backscattering shape factor f_b (see Mobley et al., 2002 for the definition of f_b and for its model), rather than by uncertainties in the remaining shape factors and related quantities. Our model for the backscattering shape factor gives predictions that are correct to within 5% for 96% of the synthetic data points used to develop the model. Figure 1 shows our ability to predict the all-important backscatter shape factor.

The question next arises as to how well we can predict the in-air remotely sensed reflectance RSR_a (which is equivalent to the remote-sensing reflectance) using all of the shape factor models. Figure 2 gives the answer: 68% of the RSR_a predictions fall within 10% of the correct (Hydrolight computed) values, and 96% fall within 20% of correct (shown by the dashed lines in Fig. 2); 86% of the predictions are within $\pm 0.0005 \text{ sr}^{-1}$ of the correct value. The model-actual correlation coefficient is $r = 0.983$.




Fig. 1. Comparison of modeled and actual f_b values. 96% of the modeled values lie within the dashed lines, which represent values within 5% of the correct value; the model-actual correlation coefficient is $r = 0.955$.




Fig. 2. RSR_a as modeled using the shape factor models. The dashed lines are 20% error bounds; the dotted lines are $\pm 0.0005 \text{ sr}^{-1}$. The model-actual correlation coefficient is $r = 0.983$.

These results are very encouraging and warrant the continued development of the RTE shape-factor inversion algorithm and its evaluation using both synthetic (Hydrolight generated) and real data obtained at sea.

IMPACT/APPLICATION

The problem of extracting environmental information from remotely sensed ocean color spectra is fundamental to a wide range of basic and applied science problems. No single inversion technique can be expected to be superior in all situations; therefore all techniques must be evaluated. In addition to investigating a new type of inversion, part of our work is to evaluate when the shape factor technique is superior to other techniques, and when it is not. This work thus adds to the existing suite of remote sensing analysis techniques.

TRANSITIONS

The shape factor models and the underlying database have been given to Dr. Frank Hoge of NASA Goddard Space Flight Center, Wallops Flight Facility and Dr. Paul Lyon of EG&G, who are using the database and the shape factor models to simulate retrievals of water IOPs from measured water-leaving radiances.

RELATED PROJECTS

Dr. Frank Hoge of NASA is separately funded by NASA for his work on development and evaluation of the shape factor inversion technique. NASA also provided funding for the work described here via an interagency transfer.

REFERENCES

Gordon, H.R., O.B. Brown, R.H. Evans, J.W. Brown, R.C. Smith, K.S. Baker, and D.K. Clark, A semianalytic radiance model of ocean color, *J. Geophys. Res.*, *93*, 10,909-10,924, 1988.

Hoge, Frank E. and Paul E. Lyon, Satellite retrieval of inherent optical properties by linear matrix inversion of oceanic radiance models: An analysis of model and radiance measurement errors, *Jour. Geophys. Res.*, *101*, 16,631- 16,648, 1996.

Hoge, Frank E., C. Wayne Wright, Paul E. Lyon, Robert N. Swift, James K. Yungel, Satellite retrieval of inherent optical properties by inversion of an oceanic radiance model: A preliminary algorithm, *Applied Optics*, *38*, 495-504, 1999a.

Hoge, F.E., C. Wayne Wright, Paul E. Lyon, Robert N. Swift, James K. Yungel
Satellite retrieval of the absorption coefficient of phytoplankton phycoerythrin pigment: Theory and feasibility status, *Appl. Opt.*, *38*, 7431-7441, 1999b.

Lyon, P. E., F. E. Hoge, C. D. Mobley, and L. K. Sundman, 2002. Inversion of the radiative transfer equation via shape factor models. Part II: Application. In preparation for submission to *J. Geophys. Res.*

Mobley, C.D., *Light and Water, Radiative Transfer in Natural Waters*, Academic Press, New York, 1994.

Mobley, C. D. and L. K. Sundman, 2001a. Hydrolight 4.2 Users' Guide, Sequoia Scientific, Inc., Redmond, WA, 88 pages (downloadable as a pdf file at www.hydrolight.info).

Mobley, C. D. and L. K. Sundman, 2001b. Hydrolight 4.2 Technical Documentation, Sequoia Scientific, Inc., Redmond, WA, 79 pages (downloadable as a pdf file at www.hydrolight.info).

Mobley, C. D., L. K. Sundman, F. E. Hoge, and P. E. Lyon, 2002. Inversion of the Radiative Transfer Equation via Shape Factor Models. Part I: Theory and Shape Factor Models. *J. Geophys. Res.*, submitted (copy available from Mobley).

Morel, Andre and Bernard Gentili, Diffuse reflectance of oceanic waters. III. Implication of bidirectionality for the remote sensing problem, *Appl. Opt.*, 35, 4850-4862, 1996.

Weideman, Alan D., Robert H. Stavn, J.R.V. Zaneveld, and M.R. Wilcox, Error in predicting hydrosol backscattering from remotely sensed reflectance, *Jour. Geophys. Res.*, 100, 13,163 - 13,177, 1995.

Zaneveld, J. Ronald V., A theoretical derivation of the dependence of the remotely sensed reflectance of the ocean on the inherent optical properties, *J. Geophys. Res.*, 100, 13,135- 13,142, 1995.

PUBLICATIONS

Mobley, C. D. and L. K. Sundman. Effects of optically shallow bottoms on upwelling radiances: Effects of inhomogeneous and sloping bottoms. *Limnol. Oceanogr.*, in press for the "Shallow-water optics" special issue.

Mobley, C. D., H. Zhang, and K. J. Voss, 2001. Effects of optically shallow bottoms on upwelling radiances: Bidirectional reflectance distribution function effects. *Limnol. Oceanogr.*, in press for the "Shallow-water optics" special issue.

Mobley, C. D., L. K. Sundman, and E. Boss, 2002. Phase function effects on oceanic light fields, *Appl. Optics*, 41(6), 1035-1050.

Mobley, C. D., L. K. Sundman, F. E. Hoge, and P. E. Lyon, 2002. Inversion of the Radiative Transfer Equation via Shape Factor Models. Part I: Theory and Shape Factor Models. *J. Geophys. Res.*, submitted.

Voss, K. J., C. D. Mobley, L. K. Sundman, J. Ivey, and C. Mazel. The spectral upwelling radiance distribution in optically shallow waters. *Limnol. Oceanogr.*, in press for the "Shallow-water optics" special issue.