AFRL-AFOSR-UK-TR-2015-0005





# The Relationship Between Visual Sensor Equipment in Flying Insects and their Flight Performance – a 'Neurobio-Engineering' Approach

# Holger G. Krapp and \*J. Sean Humbert

# IMPERIAL COLLEGE OF SCIENCE TECHNOLOGY AND MEDICINE, DEPARTMENT OF BIOENGINEERING EXHIBITION ROAD, LONDON, SW7 2BT UNITED KINGDOM

# \*UNIVERSITY OF MARYLAND DEPARTMENT OF AEROSPACE ENGINEERING COLLEGE PARK, MARYLAND 20742

EOARD GRANT #FA8655-09-1-3083

Report Date: March 2014

Final Report from 09 September 2009 to 08 September 2013

Distribution Statement A: Approved for public release distribution is unlimited.

Air Force Research Laboratory Air Force Office of Scientific Research European Office of Aerospace Research and Development Unit 4515, APO AE 09421-4515

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. <b>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS</b> .				
1. REPORT DATE <i>(DD-MM</i> -YYYY) 16 March 2014	2. REPORT TYPE Final		3. DATES COVERED (From – To) 9 September 2009 – 8 September 2013	
4. TITLE AND SUBTITLE	1 IIIdi	5a. C0	DNTRACT NUMBER	
The Relationship Between Visual Sensor Equipment in Flying Insects and their Flight Performance – a 'Neurobio-Engineering' Approach		sects ch 5b. GR FA86	RANT NUMBER 55-09-1-3083	
		5c. PF	ROGRAM ELEMENT NUMBER	
61				
6. AUTHOR(S)			ROJECT NUMBER	
Holger G. Krapp and *J. Sean Humbert		5d. T <i>i</i>	5d. TASK NUMBER	
			ORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) IMPERIAL COLLEGE OF SCIENCE TECHNOLOGY & MEDICINE DEPT OF BIOENGINEERING, EXHIBITION RD, LONDON, SW7 2BT UNITED KINGDOM			8. PERFORMING ORGANIZATION REPORT NUMBER	
*UNIVERSITY OF MARYLAND DEPT OF AEROSPACE ENGINEERING, COLLEGE PARK, MD 20742			N/A	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSOR/MONITOR'S ACRONYM(S)	
EOARD			AFRL/AFOSR/IOE (EOARD)	
APO AE 09421-4515			11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
			AFRL-AFOSR-UK-TR-2015-0005	
12. DISTRIBUTION/AVAILABILITY STATEMENT				
Distribution A: Approved for public release; distribution is unlimited.				
13. SUPPLEMENTARY NOTES				
<ul> <li>14. ABSTRACT         The grant FA8655-09-1-3083, "The Relationship Between Visual Sensor Equipment in Flying Insects and their Flight Performance – a 'Neurobio-Engineering' Approach", that was awarded to support my research and the work of Dr. Sean Humbert, (UMD, FA9550-09-1-0075) has come to an end. In this final report on the scientific activities in my lab I will stay with the tradition to provide an overview of the work that is relevant to AFRL/AFOSR – which basically includes most, if not all projects undertaken in my lab during the funding period. Despite the fact that the seasonal availability of horseflies, robberflies and hoverflies did limit progress regarding a comprehensive comparative study of across several dipteran fly species, the following sections should demonstrate that we made some important advances regarding the biological design principles of multisensory reflex control. One of the major strategic moves was to focus on gaze stabilization as an approximation to study flight control. The rational being that head movements induced by multiple sensor systems compensate for unpredictable disturbances of the thorax flies encounter on the wing. In addition, Sean Humbert showed in his work that a head orientation which enables the alignment of head-centered sensor systems with the inertial vector significantly simplifies the measurement of state-changes required for feed-forward/feedback-based flight control. As a corollary of our joint studies we should state: Flying insects which are subject to significant inertial forces during flight and have limited a separate final report on his efforts, I will focus here on activities in my lab, adding occasional cross-references to his report where appropriate. </li> </ul>				
16. SECURITY CLASSIFICATION OF: a. REPORT b. ABSTRACT c. THI	17. LIMITATION OF ABSTRACT	18, NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON Gregg L. Abate	
UNCLAS UNCLAS UNCI	SLAD SAR	35	<b>19b. TELEPHONE NUMBER</b> (Include area code) +44 (0)1895 616021	
		· .	Standard Form 298 (Rev. 8/98)	

## **USAF Grant FA8655-09-1-3083**

Holger G Krapp

email: h.g.krapp@imperial.ac.uk

Department of Bioengineering Imperial College South Kensington Campus London SW7 2AZ

## EOARD Contract No: FA8655-09-1-3083 (HGK) US Contract No: FA9550-09-1-0075 (JSH)

"The Relationship Between Visual Sensor Equipment in Flying Insects and their Flight Performance – a 'Neurobio-Engineering' Approach" (Period of finding: October 2009 – September 2013)

# Final-Report (16 March 2014)

J. Sean Humbert Department of Aerospace Engineering University of Maryland, College Park, MD 20742 Email: <u>humbert@umd.edu</u>

Holger G. Krapp Department of Bioengineering Imperial College, London, SW7 2AZ, UK Email: <u>h.g.krapp@imperial.ac.uk</u>

## Contents:

- 1. General Summary of Achievements
  - 1.1 Publications
  - 1.2 Scientific Conferences, Workshops and Seminars
  - 1.3 Leadership
  - 1.4 Funding
  - 1.5 Group Size, Collaborations, and Scientific Impact
- 2. Report on Current Projects
  - 2.1 Behavioural performance
  - 2.2 Biomechanics of motor systems
  - 2.3 Neuronal mechanisms
  - 2.4 Modelling of gaze stabilization
- 3. Conclusion
- 4. List of Publications

# 1. General Summary of Achievements.

The grant FA8655-09-1-3083, "The Relationship Between Visual Sensor Equipment in Flying Insects and their Flight Performance – a 'Neurobio-Engineering' Approach", that was awarded to support my research and the work of Dr. Sean Humbert, (UMD, FA9550-09-1-0075) has come to an end. In this final report on the scientific activities in my lab I will stay with the tradition to provide an overview of the work that is relevant to AFRL/AFOSR – which basically includes most, if not all projects undertaken in my lab during the funding period.

Despite the fact that the seasonal availability of horseflies, robberflies and hoverflies did limit progress regarding a comprehensive comparative study of across several dipteran fly species, the following sections should demonstrate that we made some important advances regarding the biological design principles of multisensory reflex control. One of the major strategic moves was to focus on gaze stabilization as an approximation to study flight control. The rational being that head movements induced by multiple sensor systems compensate for unpredictable disturbances of the thorax flies encounter on the wing. In addition, Sean Humbert showed in his work that a head orientation which enables the alignment of head-centred sensor systems with the inertial vector significantly simplifies the measurement of state-changes required for feedforward/feedback-based flight control. As a corollary of our joint studies we should state: Flying insects which are subject to significant inertial forces during flight and have limited computing power, gaze stabilization is a functional necessity that enables high aerial manoeuvrability.

Because Sean Humbert has submitted a separate final report on his efforts, I will focus here on activities in my lab, adding occasional cross-references to his report where appropriate.

# 1.1 Publications

During the funding period of this grant from Oct 2009 until the end of Sept 2013 my group has published 13 papers in peer reviewed journals (O32-45), 3 book chapter (R4-6), and 5 peer-reviewed conference papers (C4-9). Six of the peer-reviewed papers were published in high profile journals ranked 'internationally competitive'.

Finally, I am a co-editor of last year's proceedings of the 'Living Machine Conference' on biomimetic and biohybrid systems.

We are currently working four further scientific publications one of which will be submitted within the next week (O46), with the others to follow within the next month (O47-49).

# 1.2 Scientific Workshops, Conferences, and Seminars

In total I attended 22 scientific events within the duration of this grant across a wide range of subject areas from the five-yearly specialized International Conferences on Invertebrate Vision to the novel annual Living Machine Conference which is dedicated to bio-inspired technology. At three of those conferences Dr Kit Longden, the RA on this grant, also presented his work. In addition I was invited to give seminar presentations in continental Europe and in the UK at 7 occasions.

Many of the conferences were in connection with the bio-MAV/UAS SOAR meetings, bio-inspired GNC at Eglin Air Force Base, or other events relevant to the Programme Arrangement between AFRL/AFOSR and DSTL on bio-inspired technology.

# 1.3 Poster and Oral Presentations at National and International Conferences

Members of my group presented 14 poster (A43-54) and oral contribution at 7 national and international conferences.

# 1.4 Funding

# *Current funding secured for research in my laboratory (excluding FA8655-09-1-3083):*

- Wellcome Trust research grant, PI: "Integrated reflex control" in Manduca. [Funding until end of April 2014]
- **DSTL-sponsored EPSRC Industrial CASE PhD Studentship**, PI: "The significance of (image) gaze stabilization Comparative studies on gaze control design in flying insects using behavioural, computational, and electrophysiological techniques." [Funding until end of March 2015]
- **DSTL National PhD Programme**, PI: "The significance of (image) gaze stabilization functional characterization of the neck motor system using electrical stimulation of identified neck muscles in blowflies." [Funding until end of September 2016]

# Grant proposals submitted:

- AFOSR/EOARD, PI, together with Dr Graham Taylor, PI, University of Oxford, UK, and Dr Sean Humbert, PI, University of Maryland: "*The mode sensing hypothesis*" [1<sup>st</sup> year funding currently being set up at UMD]
- French-UK PhD studentship scheme (DSTL), PI, together with Dr Stephane Viollet, CNRS, Marseille, FR: "Head-body movements in freely flying flies when chasing a dummy target"

# Grant proposals in preparation:

• AFOSR/EOARD, PI, together with Prof Doekele Stavenga, University of Groningen, NL: "Polarization Vision in Horseflies"

# 1.5 Group size, Collaborations, and Scientific Impact

## Group size:

2 RAs (postdocs), 3 postgraduate students (PhD), 1 MSc and 5 undergraduate students. I co-supervise a DSTL-funded DPhil student working in Graham Taylor's lab, University of Oxford, on the mode sensing hypothesis in *Manduca sexta*, and a PhD student in Barbara Webb's lab, University of Edinburgh, funded by the BBSRC who works on visual navigation in ants.

# **Collaborations**:

I have collaborations at Imperial College in the Department of Bioengineering with: Dr Martyn Boutelle (Biosensors), Dr Simon Schultz (Neural Coding), Dr Manos Drakakis (Low-power VLSI technology), and Dr Reiko Tanaka (Compound Control). To study the functional anatomy of insects we still collaborate with the micro-CT group at the Natural History Museum, London. Our collaboration with scientists at the Paul Scherre Institute, Swiss Light Source, CH, to obtain high-resolution 4D data from tethered flying flies on the functional organization of their fight and neck motor systems is ongoing – we are currently publishing the results of to beam time grants I got awarded in the past. Collaborations with Prof Simon Laughlin (Cambridge) and Dr Graham Taylor (Oxford) have been maintained. Work with Graham Taylor and Prof Sean Humbert (UMD, USA) will be intensified by means of the AFOSR-funded grant on the 'mode sensing hypothesis'. A new collaboration has now been established with Prof Barbara Webb to study ant navigation. Further international Collaborations include work with Prof Martin Egelhaaf, Neurobiology, Bielefeld University, Germany; Prof Fabrizio Gabbiani, Baylor College of Medicine, USA; and Mr Ric Wehling at the AF Research Laboratory, Eglin, US. A new collaboration with Prof Doekele Stavenga, University of Groningen, NL, and Dr Stephane Viollet, CNRS, Marseille, FR are in planning.

## Current Bibliometric Data and Scientific Impact:

## **Bibliometrics**:

	March 2014
Number of citations*	2137
h-factor*	25

\* = according to "Google Scholar"

## Scientific Impact/Leadership:

• *Reviewing manuscripts* for more than 30 peer review journals, including *Nature*, *Science*, and *Neuron*.

- *Reviewing grant applications* for 10 funding agencies including *RCUK*, *NSF* (*USA*), and *AFOSR* (*USA*).
- *Expert Reviewer* for *European Commission FP7* collaborative project
- Member of Academic Editorial Board of open access journal PLoS ONE
- *Financial Co-organizer* AFOSR/DSTL "*Bio MAV SOAR*" meeting, Chilworth Manor, 2011, as well as UK follow-up meetings at Imperial College, London, and University of Oxford, 2012.
- Member of *Programme Committee* of Conference on *Biomimetic and Biohybrid Systems* "Living Machines", Barcelona and London 2012 and 2013
- *Co-ordination* of DSTL-supported virtual *Centre of Excellence* on Unmanned Autonomous Systems (*UAS former "MAV"*)
- Local Co-organizer (Satellite events chair) of Conference on Biomimetic and Biohybrid Systems "Living Machines", London, 2013
- *Co-applicant/proposer* of research call initiatives in the area of sensing and actuation (ESF/NSF) and *"Robot Companions"*, Flagship proposal submitted to the European Commission.

I also took on leadership responsibility in fostering areas relevant to the remits of AFRL/AFOSR, contributing to high standards in scientific research by committing to reviewing and editorial work, as well as applying for an EPSRC Centre of Doctoral Training in *Bio-inspired Systems and Technologies*, strongly supported by DSTL.

# 2. Report on current projects:

I will outline the achievements during the funding period of this grant (FA8655-09-1-3083) in relation to the four major directions of my work:

- 2.1. *Behavioural performance* of multisensory motor control in dipteran flies
- 2.2. Biomechanics of motor systems involved in gaze and flight control
- 2.3. Neural mechanisms underlying multisensory motor control in dipteran flies
- 2.4. *Modelling* of multisensory motor control design

These four approaches result in a comprehensive description of sensorimotor control. The results obtained at different system levels and by applying a variety of experimental and theoretical approaches do inform and support each other. Altogether they enable an iterative cross-validation process that facilitates the derivation of models capturing the fundamental design principles of biological sensorimotor control. Although funding for my research over the past nine years was also provided by the the BBSRC, Human Frontier Science Program Trust, Wellcome Trust, DSTL and Imperial College resources, I will briefly summarize earlier findings (presented in earlier reports) and the development over the last couple of months followed by a conclusion.

# 2.1. Behavioural performance

## 2-degrees of freedom controller

We applied a linear systems analysis approach to the gaze stabilization system in dipteran flies to quantify the interaction between the compound eyes (mediating optic flow information), the ocelli (mediating information on fast changes in attitude) and the mechanosensory halteres (mediating rotation rates) in the high dynamic input range.

In our original studies on *Calliphora* we performed a systems identification of the gaze stabilization system which showed a surprisingly linear performance under the conditions tested (C4). An intermodulation distortion (IMD) analysis indicated that internal non-linearities in the system contribute less than 4% to the output signal.



Figure 1: 2-degree of freedom controller combining feedforward signals provided by the halteres ( $F_{ff}$ ) with feedback signals provided by the motion vision pathway (Ffb). The properties of the neck motor system are included by the transfer function of the neck motor system (Fn). For further details see text. Figure modified from (C4).

The resulting control architecture and transfer functions describing the relationship between the stimulus (thorax rotation, TR) and the systems output (head rotation, HR) are shown in Fig 1.

Recent work on the interpretation of the architecture and further analysis of the data revealed that the system including the halteres and the compound eyes can be described as a 2-degrees of freedom controller where 1 degree of freedom is given by a feed forward component ( $F_{ff}$ , halteres) and the second degree of freedom represents a feedback component ( $F_{fb}$ , compound eyes). This ingenious combination of fast feed forward signals mediated by the halteres and slow feedback signals from the compound eyes works as follows: immediate haltere-induced compensatory head roll shifts the retinal slip speed distribution caused by thorax rotation towards to lower dynamic range (subtraction of initial head roll from the thorax roll) that can then be analysed by the motion vision pathway of the compound eyes. Thus, the halteres 'enable' the visual system by eliminating those fast slip speed components the visual system would not be able to process due to its band-pass characteristics.



Figure 2: Probability density functions of retinal slip speeds under three experimental conditions and at three different thorax oscillations, 3 Hz (upper panel), 6 Hz (middle panel), and 15 Hz (bottom panel). The three traces in each panel show the distribution under condition where the head would be fixed to the thorax (HR = TR, blue), the halteres and the motion vision pathway (green), and only the motion vision pathway (red) contribute to compensatory head roll. Note that the contribution of the halteres to shift the slip speed probability density function depends on the stimulus frequency. At the highest stimulation frequency halter function is essential for enabling the processing of visual motion. If the halteres are disabled visual feedback causes a shift of the density function to higher slip speeds, which has a detrimental effect on gaze stabilization. (unpublished data).

The data presented in Figure 2 demonstrate the principle. They show probability density functions of the retinal image shifts (slip speed) the fly encounters during different experimental conditions, i.e. the frequency of occurrence plotted against slip speed. Three different oscillation frequencies of the thorax have been applied, 3 Hz, 6 Hz, and 15 Hz. The blue traces in the subpanels indicate the distribution the fly would experience if its head was was fixed to the thorax. The green traces correspond to the distribution when the head is free to move and both the compound eyes and the halteres are contributing to the compensatory head movements (+ Hs). In this case the slip speed distribution is shifted to the left, i.e. to lower slip speeds. Red traces show the shift of the distribution after the halteres were removed. At thorax oscillations of 15 Hz the significance of the feed forward component becomes most obvious (Fig 2, bottom panel). With haltere and compound eye contributions the slip speed distribution is markedly shifted to the left (green trace) while without halters (red trace) a shift of the distribution to the right is observed. These data demonstrate a cooperative effect of the two control degrees of freedom that is different from just adding up the signals generated by the compound eyes and the halteres alone.

This interpretation is in agreement with the qualitative finding that flies the halteres of which have been removed will immediately crash if thrown in the air. Only if a light weight thread is attached to their abdomen, they are able to fly again. The thread, in this case adds passive aerodynamic stability and dampens fast attitude changes resulting in high retinal slip speeds. With fast retinal slip speeds removed the visual system is sufficient to provide feedback signals for flight control.

A necessary condition for this 2-D controller to work is that the motion vision component of the controller provides stable feedback. We assessed the stability margins of the feedback controller and found that both the gain and the phase margins are unexpectedly high with 20 dB and 130 degrees, respectively. Our interpretation of those findings is that the system is specified to deal with a high variance of unpredictable input signals which, in combination with noise in the sensory pathways, potentially results in extreme output values without becoming instable.

We are currently writing up our work on the 2-degree of freedom controller for publication (O49).

## Studies on gaze stabilization in robberflies

The supply of robberflies from the US turned out to be difficult. We observed a severe reduction of the life expectancy during intercontinental transportation by air, probably because the fly respiratory system does not tolerate the massive pressure changes in the cabin of air carriers. We therefore collected two UK robberfly species in the south of England, *Dysmachus trigonus* and *Philonicus albiceps*. The former species was used in preliminary electrophysiological experiments (see below), while the latter had been used for behavioural experiments to study gaze stabilization, applying the same linear systems approach as to blowflies the results of which were reported earlier.

Robberfies are more difficult to study under tethered flight conditions. They do not assume the same stable flight position a *Calliphora* and often extend their legs to counteract external body rotations. We were nonetheless able to perform some preliminary gaze stabilization experiments on *P. albiceps* which provided us with some insight regarding general principles and species-specific adaptations.

Figure 3 shows the data obtained from at least nine robberflies the thorax of which was oscillated over the range from 1-10Hz at an amplitude  $\pm 30^{\circ}$ . The bode plots shown were obtained under four different conditions – the same that had been applies to blowflies in previous studies – in terms of sensory inputs: C1 (orange trace) all sensory inputs intact (n=11), C2 (dark blue) compound eyes and ocelli (n=9), C3 (red) compound eyes and halteres (n=11), and C4 (light blue) only compound eyes (n=9).

The most conspicuous difference in the gaze stabilization performance compared to the control condition where all sensors are intact was observed after the removal of the halteres; a similar result was found in blowflies (C4). Removing the halteres drastically increase the phase delay of the response (blue traces, bottom plot) – in particular when the ocellar input is disabled (light blue trace). The response gain shows no statistically significant differences, except for the condition where both the haltere and ocellar inputs are excluded. Any differences between the responses under the four conditions measured in the low dynamic range (1-3 Hz) are probably due to high interspecific variability. Overall, the results obtained in *P. albicepts* are comparable with those obtained in *Calliphora*. In both species haltere input reduces the phase delay of the response (orange and red traces, bottom plot).



Figure 3: Frequency response of compensatory head roll in P. albiceps. Top plot: Head roll gain against oscillation frequency for four different combinations of sensory inputs (see inset, bottom plot). Probably due to a large response variability across animals no significant differences were found for three out of four conditions. When only motion vision contributes to the response the gain is significantly reduced for an oscillation frequency of 10 Hz. Bottom plot: The head roll phase angle relative to the phase of the stimulus as a function of stimulus frequency. Haltere contribution results in a minimal phase delay over the entire dynamic range tested. There is a significant difference in phase delay between responses obtained without halteres (blue traces) and all other stimulus conditions. The largest phase delay is observed when only the motion vision pathway contributes to compensatory head roll (light blue trace). (Unpublished data).

In a direct comparison between the gain of compensatory head movements measured in blowflies and robberflies mediated only by the motion vision pathway (ocelli covered and halteres removed) we found that compensatory head roll in *Philonicus* covers a higher dynamic input range compared to *Calliphora* (Figure 4). These results could suggest that the motion vision pathway in robberflies is tuned to higher angular velocities. But it could also mean that the neck motor system of the two species has different dynamic properties. The latter aspect would be supported by the fundamental difference the anatomical constraints of the neck motor. In robberflies the head is positioned more distantly from the thorax supporting larger head movements. Further functional anatomical (see below) and behavioural studies will be required before our results can be properly interpreted.



Figure 4: Gain of compensatory head roll in robberflies (n=9) and blowflies (n=7) mediated by the motion vision pathway. The thorax of the flies was oscillations at +/- 30° at different frequencies. At high stimulus frequencies robberflies show a significantly higher gain. Error bars give SEM (unpublished data).

#### Interactions between motion vision pathway and ocelli

Since the work by Parsons et al. (O20, O34) we know that the LPTCs (VS-cells) are modulated by ocellar stimulation suggesting both pathways to use a common sensory coordinate system, likely to be set up by the LPTCs (O26).

One of the major objectives of this project was to study the interaction between the compound eye-mediated motion vision pathway and the ocellar pathway in the gaze stabilization system of different dipteran species. Although the availability of species other than our lab-bred blowflies was rather limited throughout the duration of the grant, we did achieve comparative measurements on blowflies and robberflies.

In a first set of experiments we oscillated a dark visual hemisphere as a substitute for ground around tethered flying blowflies. Under these conditions the animals had access only to visual input, but no haltere information, where the ocelli were either covered (compound eyes) or uncovered (compound eyes + ocelli). The results in terms of compensatory head roll are shown in Figure 5. We found that under these two conditions, over the range of oscillation frequency from 1 - 10 Hz there was no significant difference between the response gain observed. The phase, however, showed a clear advance of the response when the ocelli were contributing for frequencies > 3 Hz. This result suggests a non-linear integration of ocellar and motion vision signals.



Figure 5: Compensatory head roll mediated by compound eye and compound eye + ocelli information. Left panels: Linear response gain (upper panel) and response phase (lower panel) plotted as a function of pattern oscillation frequency. Note that there is no significant difference between the condition with and without ocelli regarding the response gain while the response phase becomes larger for oscillation frequencies > 3 Hz. Right panels: The simulation of the ocelli and compound eye as simple highpass (tau = 12.5ms) and lowpass filters (tau = 12.5ms), respectively, combined with an 'integrate and fire model' of the descending neuron that included a static threshold non-linearity qualitatively captures the behavioural results. (unpublished results).

We combined a phenomenological approximation of the two sensor system properties by means of a low pass (motion vision) and a high pass (ocelli) filter, with time constants in the physiologically plausible range, and a biophysically motivated leaky integrate and fire model. A biophysically motivated 'integrate and fire model' was implemented to compute the membrane potential of a descending neuron that integrates the outputs from both sensor systems (Figure 6). Like all 'real' nerves cells the descending neuron incorporates a static threshold non-linearity. Only beyond a certain threshold membrane potential does the descending neuron generate action potentials, which are necessary to produce muscle contractions in the neck motor.



Figure 6: A model that qualitatively explains the observed phase increase in compensatory head roll of Calliphora when the ocelli are disabled while the gain stays the same for compound eye and compound eye + ocelli stimulation. (unpublished).

This model qualitatively reproduces the effect observed at the behavioural level: while the gain is the same for both experimental conditions, the phase of the responses at higher stimulus frequencies is advanced if the ocelli contribute (lower right panel). This comparatively simple model suggests a static threshold non-linearly as the potential basis for the observed behavioural data.



Figure 7: First studies on the interaction of ocelli and compound eye information in the robberfly Dysmachus trigonus (left panel). In these experiments the halteres of the fly were removed and rather than oscillating the ground pattern the thorax of the animals were sinusoidally rotated around the roll axis. The main senor systems contributing to the response were therefore the compound eyes or the compound eyes + ocelli. Right panels: Similar to the experiments in Calliphora, the gain was similar under the two conditions (upper plot) but there was a difference in the response phase (lower plot). (unpublished data).

We have also started performing specific experiments on ocellar and compound eye signal integration to study gaze stabilization in UK robberfly species *Philonicus albiceps* and *Dysmachus trigonus*. The behavioural results indicate that similar non-linearities as described for blowflies above may apply to the gaze stabilization systems in robberflies. Whether or not a static threshold non-linearity at the neuronal level best explains our findings regarding ocellar and compound eye signal integration across all dipteran species is not entirely clear at this point.

Next month we will perform more experiments on *Calliphora* – and as soon as they will emerge in the wild, on robberflies, horseflies and hoverflies – to substantiate our data base. We will also refine our open loop integration model. The model will be included as a first non-linear element in our otherwise linear closed-loop simulation platform of the gaze stabilization system (cf. section 2.4) before writing up the results for publication.

Altogether, the results of our behavioural experiments on gaze stabilization in dipteran flies proved to be a worthwhile paradigm to study the relationship between sensory and motor coordinate systems. We expect further advances regarding a quantitative description of the weighting given to the various sensor systems by applying band-limited white noise stimuli with different statistical properties and by new experiments where pattern and body rotations may be combined in different ways.

# 2.2. Biomechanics of motor systems

## The neck motor system

A comprehensive understanding of the relationship between sensor and motor coordinate systems requires a functional characterization of the motor system. To this end we are combining functional anatomy based on x-ray techniques such as  $\mu$ -CT with electrical muscle stimulations and biomechanical modelling.

Based on previous research in collaboration with the Natural History Museum London, we are currently advancing methods to increase tissue contrast in  $\mu$ -CT scans to enable automatic segmentation of different tissue types. Our first results, still based on manual segmentation, provided us with a 3-dimensional reconstruction of the neck motor system in the blowfly *Calliphora* which will serve two purposes: (i) the 3D data allow us to refine our dissection methods in terms of minimizing structural damage to the system when placing electrodes for electrical muscle stimulation (ii) it will inform biomechanical models of the neck motor system, e.g. Finite Element Models, which capture the dynamic properties of the system.



Figure 8: A quantitative and qualitative comparison of the 5 preparation methods for the final staining day. A: A comparison of day 7 contrast ratios across the 5 staining protocols, for tissue types (inset). B-E: Coronal slices of ETH-PTA (B), PFA-PTA (C), ETH-Lugol (D), PFA-Iodine (E) at approximately the same depth and orientation. The slice for ETH-Iodine exhibits similarly poor contrast to that of PFA-Iodine and is omitted. (data from O48)

Figure 8 shows some of the results we obtained in a study using different fixatives (e.g. ETH = 100 ethanol, PFA = paraform aldehyde) chemical agents (e.g. iodine, PTA = phosphotungstic acid) and staining protocols to increase the tissue contrast. In these experiments we did not optimize for the highest voxel resolution, which was just below 10  $\mu$ m<sup>3</sup>, but to increase the tissue contrast ratio. Our data show that the best ratios are obtained using a combination from ETH or PFA as a fixative and PTA as a contrast agent. We are currently preparing a manuscript for publication reporting the results on our systematic study to improve the  $\mu$ -CT methodology (O48) for the functional analysis of insect motor systems.

In earlier work reported previously we applied iodine as a contrast agent to reconstruct the 3-dimensional organization of the neck motor system. As mentioned above, the results were meant to develop new dissection methods applied in the context of electrical stimulations of neck muscles to study their dynamic properties (Figure 9).



Figure 9:  $\mu$ -CT-based reconstruction of the neck motor system in Calliphora. From left to right: right, top and left view of the blowfly head. False colours mark muscles attached to cuticular structures in the thorax and the head enabling gaze stabilization. (unpublished data).

The latest advances in tissue contrast presented above will allow us to refine the current 3-dimensional anatomical model of the *Calliphora* neck motor system. We are also planning to obtain similar data from other dipteran species for a comparative study.

A key component of our work will be the specification of a biomechanical model that captures the properties of the neck motor system. We decided to apply finite element modelling (FEM) to the system as it provides a convenient tool to describe and analyse the function of complex motor system. The FEM will be informed by our 3-dimensional anatomical data – and later by electrical stimulation experiments. Currently, we base the model on assumptions regarding the local Young's modulus. Despite this initial simplification of the system's functionality, we have been successful to generate head-body movements by introducing local forces simulating muscle contraction (Figure 10).

The combined approaches of physiological characterization and modelling of the system's properties will mutually support each other. Once a sufficiently detailed model is derived and validated, a dimensionality reduction will be performed to

obtain a simplified version of a neck motor system that may be implemented in technical applications.



Figure 10: Finite model of the Calliphora neck motor system based 3-dimensional on anatomical data (ventral element view). Each describing part of a neck muscle is assigned a Young's modulus of а certain orientation. Simulating forces different muscles in the introduces relative movements between head and body. (unpublished data).

## The flight motor system

Together with Graham Taylor's group in Oxford and a group of scientists at the Paul Scherrer Institut at the Swiss Light Source we have advanced x-ray scanning techniques in terms of spatial and temporal resolution in a way that enabled us to monitor the 4-dimensional operation of the flight motor in blowflies. The animals were subjected to hard x-ray radiation while being rotated around the roll axis in tethered flight. Under these conditions the flies were able to survive for several minutes and engaged on of showed asymmetric wing trajectories in an attempt to compensate for the externally forced rotation (Figure 11).



Figure 11: Schematic illustration of Calliphora observed in a synchrotron during tethered flight. The fly responds to a forced body rotation bygenerating a differential output of the flight motor - here by increasing and decreasing lift production in the left and right wing, respectively. The x- and z-axes correspond to the longitudinal and vertical body axes of the fly. (figure taken from *O44)*.

Due to retrospective gating of images phase locked to the wing beat cycle we were able to observed the activity of the power and steering muscles during the compensatory flight manoeuvres. Figure 12 show an external view of the thorax and a view from the same perspective after virtually removing large parts of the cuticle. Phase contrast – as opposed to absorption contrast in  $\mu$ -CT – was used successfully to

resolve even the buckling of tendons of steering muscles throughout the wing beat cycle.



Figure 12: External and internal view of the Calliphora thorax. Left: External view of the thorax showing cuticular structures of the wing hinge and parts legs. Right: Virtual dissection of the thorax showing the power muscles of the flight motor (yellow, orange, and red) as well as four small identified steering muscles the activation of which could be analysed during asymmetric tethered flight in a synchrotron. (Figure from O44).

This technique allows us to analyses periodic movement patterns in any motor system of flying insects as long as an appropriate gating signal is available that is correlated with the movement. Given the high temporal and spatial resolution of the technique, in further studies we will be able to extract general principles of non-linear and linear operations in biological motor systems which will inform the development of biomechanical models. In combination with a better understanding of the neural mechanisms underlying these models will inform novel approaches to the design of small scale mechanical devices with strong potential for application in autonomous robotic systems.

On our highly successful synchrotron work, that was initiated by Daniel Schwyn, a former PhD student in the lab, we published two up-front reports in technical journals (O41 and O43) last year. A high profile publication of our results on the flight motor system is now accepted for publication in PLoS Biology (O44) with another one focusing on the ground-breaking new methodology currently being prepared (O46).

# **2.3.** *Neural mechanisms* underlying multisensory motor control in dipteran flies

# Optic flow processing depends on locomotor and nutritional state

A question of increasing interest concerns a notorious problem of biological systems that is even as significant for micro air vehicles with small payloads: limited energy supply.

Neural information processing is costly and requires sensory systems to adjust energy expenditure very tightly to the current needs. Work in our lab has shown that the activity of LPTCs in the motion vision system is increased during locomotion (O31, O36, O38).

In electrophysiological experiments where a tethered fly is able to walk on a trackball, by monitoring its forward speed and yaw rotations during stimulated with visual motion (Fig 13A, upper panel) we found that: The responses of the LPTC H2 (Fig 13A, lower panel) to temporal frequencies beyond 7 Hz of a moving patter are elevated when the fly is walking (Fig 13B, red trace) compared to those in stationary flies (Fig 13B, black trace). The increased responses are costly for the fly in terms of energy consumption but indicate a necessary adaptational mechanism to adjust its signalling range to a higher dynamic input range during locomotion. Food deprivation abolishes this adaptation in a gradual way (Fig 13C). We also found that the response increase was linearly related to walking speed in both fed (Fig 13D, grey line) and food-deprived flies (Fig 13D, black line), but that in food-deprived flies the increase was systematically reduced. In addition, the response increase was found to be inversely proportional to the number of days the flies were starved (Fig 13F). Our findings at the physiological level were corroborated by studying the vaw optomotor behaviour in flies, using wide-field motion for visual stimulation (Fig 13E, left panel). In fed flies the temporal frequency tuning of the bahaviour for frequencies from and above 4 Hz was high and stayed at a similar level over the entire dynamic input range (Fig13 E, grey trace filled circles). Flies starved for 3 days showed a significantly reduced optomotor response (Fig 13E, black trace) but recovered partly after 1 day of sucrose feeding (Fig 13E, grey trace filled triangles).



Figure 13: The temporal frequency tuning of the identified H2-cell depends on the locomotor state of Calliphora and its nutritional state. A: Schematic illustration of the experimental setup. B: Temporal frequency tuning in walking and stationary flies. C: Temporal frequency tuning in fed and food-deprived fies. D: Dependence of neural response on walking speed. E: Schematic illustration of optomotor setup (left) and optomotor responses in fed flies, food-deprived flies, and flies which were fed sucrose after 3 days of food deprivation. F: Neuronal response increment due to adaptation plotted over the number of days starved. Note that the walking-dependent adaptation of the temporal frequency tuning (panel B) is reduced due to starving. (Data from O45).

Together with other studies on the olfactory systems in starved flies, where the activity is actually increased (rather than decreased), our results suggest that under limited energy supply adaptational mechanisms in the visual system are reduced. The interpretation is that remaining energy resources are re-allocated to sensory task which increase the likelihood of survival – here: olfaction, which aids in locating food sources. This work has now been accepted for a high profile publication in Current Biology (O45).

# Comparative electrophysiology on Lobula Plate Tangential Cells in dipteran flies.

As mentioned earlier, to identify general principles and species-specific adaptations of multisensory integration and to relationship between sensory and motor coordinate systems supporting the 'Mode Sensing Hypothesis' (O42, R4) requires to study of a variety of different dipteran fly species. Ideally, those studies would include a description of the functional anatomy and modelling of the motor systems, behavioural experiments and an assessment of the neural populations involved in optic flow processing. The four families of dipteran flies we planned to include in our study are shown below. They all have different morphological and anatomical features likely to result in different aerodynamic properties, a pronounced variety of lobula plate tangential cells (LPTCs), and considerably different life styles. Together these aspects suggest specific relationships between sensory and motor coordinate systems.



Calliphoridae

Asilidae

Syrphidae

Tabanidae

Due to the limited availability of the different species, we have not yet been able to study LPTCs in all these species. Besides the detailed knowledge about the subpopulations of LPTCs in *Calliphora* – the reference species of this study – we were only able to record form a number of optic flow processing visual interneurons in robberflies so far, as I reported earlier. The results do far suggest that there are indeed species-specific differences in the response properties between *Calliphora* and the robberfly *Philonicus albiceps*. The responses of LPTCs, for instance, had a peak at higher temporal frequencies compared to *Calliphora*. We also found LPTCs in robberflies that had surprisingly small receptive fields, localized in the fronto-equatorial region of the visual field. Cells with such properties in *Calliphora* had not yet been reported although at the level of neck motor neurons (O26) similar visual receptive field sizes were found which potentially support some sort of edge-detection mechanism (e.g. Hengstenberg 1991).

Despite those some differences in the cellular equipment of the lobula plate, there is one type of LPTC that had been identified in *Calliphora* which was found in all other

species studied so far, even in the othopteran locust. A receptive field organization similar to that of the so-called H1-cell in *Calliphora* was also found in robberflies and horseflies (Figure 14).



Figure 14: H1-like LPTCs in robberflies and horseflies. The panels show the directional motion preferences and sensitivities of motion sensitive interneurons in blowfly (a), robberfly (b), and horsefly (c) plotted against azimuth and elevation within the right visual field. LPTC of the type – which has been identified as H1-cell in blowfly – respond to back-to-front wide-field motion in the right visual hemisphere. In Calliphora the H1-cell in involved in detecting rotations round the vertical axis (yaw).

We will increase our efforts to investigate LPTCs in hoverflies, robberfies, and horseflies over the summer. A dedicated rig has been set up in the lab that allows us to perform both extracellular and intracellular recordings including intracellular staining to identify the cell under study. For flexible visual stimulation we will use LED panels as described in one of my earlier reports. In addition we will be able to use another rig that includes a back-projection system to enable wide-field optic flow stimulation.

## Fly-robot interface for studying multisensory integration

A complementary approach to study multisensory integration has been developed over the last three years that utilizes a robotic fly-brain interface. As outlined in earlier reports the goal is to use neural spiking activity recorded on-board from an H1-cell to steer the robot under closed-loop conditions. To do so we had to miniaturize an entire electrophysiology set up so it fits onto a small 2-wheeled robot (Figure 15). This process has now been completed. Currently we are working on a control architecture that links the H1-cell signals to the motors driving the wheels. A challenge to overcome is that the the H1-cell increase its spike rate upon back-to-front motion (cf. Figure 14). Thus during forward motion of the robot the H1-cells in either side of the visual system would be inhibited and could not be used to control the wheels of the robot in order to avoid collisions with obstacles in the surroundings.



Figure 15: A robotic fly-brain-interface to study multisensory integration in Calliphora under closed-loop conditions. A small twowheeled robot (diameter  $\sim 10$  cm) carries a miniaturized electrophysiology setup. Micro electrodes will be positioned within the lobula plate using mini micromanipulators mounted on hollowed bases which contain shielded recording electronics. The recoding platform is isolated against the mechanical vibrations caused by the motors by a layer of foam material. The fly is fixed to a holder mounted on a central column. Signals from the H1-cell, a spiking LPTC, are extracellularly recorded and will be used to generate motor commands sent to the motors of the wheels. A/D conversion and signal processing will take place on-board. The neural signals will be used in a control architecture to avoids collision with obstacles in the environment while the robot is freely moving.

This problem will be solved by implementing a control architecture that exploits efference copies (forward models) of the command sent to the motors of the wheels forcing an undulating trajectory of the robot. The efference copie will result in a prediction of the neuronal response based on an internal model of the relationship between angular velocity of a visual pattern and the H1-cell response. The predicted spike rate will then be subtracted from the spike rate actually recorded from the H1-cell where the remaining difference is then used as a feedback signal to control the motors (Fig 16).



16: Control Figure architecture to establish collision avoidance of a robot steered by a fly. 'Robot model' and 'HImodel' refer to the relationship between command voltage and angular velocity of the robot and the relationship between the angular velocity of visual motion and the spike rate of the H1cell. (publication in preparation).

Such strategy requires (a) a forward model of H1-activity (Fig 17, left) and (b) the known input-output relationship between the control voltage sent to the wheels and the angular velocity of the robot (Fig 17, right). For the architecture to work in the

context of collision avoidance a certain response properties of the H1-cell are critical: close objects will introduce a stronger response in the neuron than objects at larger distances due to distance-dependent decreases in contrast. The other property is that the response of the H1-cell is a monotonic function of temporal frequency (Fig 17, left).

Work in of the group of Prof Egelhaaf on horizontal LPTCs suggested that so-called HS-cells may indeed be used by the fly to assess distance – although only during translational phases of flight following fast rotational body saccades. A similar strategy of 'active vision' in combination with an efference copy as described above will be implemented on the robot.



Figure 17: Angular velocity dependence of the H1-cell response in the lab environment (left). Note the monotonic relationship between positive angular velocities neuronal spike rate which will be implemented as 'H1-cell model' in the closed-loop control architecture in Fig 16. Right: Linear relationship between the command voltage and the angular velocity of the robot which will be implemented as the 'robot model' in the closed-loop control architecture shown in Fig 16. (publication in preparations).

Eventually, we will simultaneously record the activity of other LPTCs of which we know they reflect the integration from sensory modalities other than motion vision (e.g. O20, O34) in a fly that is actually moving in space. We will then compare the neuronal responses recorded when different senor systems are disabled to assess signal integration under closed loop conditions.

# 2.4 Modelling of gaze stabilization

In collaboration with Dr Reiko Tanaka (Dept Bioengineering) we have been working continuously on refinements of a closed-loop simulation platform that models the gaze stabilization system of *Calliphora*. Most modelling approaches were based on behavioural experiments (section 2.1) and assumed – in a first approximation – a linear time-invariant system (e.g. C4). One major result of this approach was the 2 degree of freedom controller discussed above.

We have recently expanded the concept of the 2 degree of freedom controller to include the stabilization of the body against unpredictable external disturbances. It is important to mention that the diagram shown in Figure 18 is purely conceptual at this point in time and has not yet been tested experimentally with respect to the flight control component. What makes it plausible in terms of its potential implementation in the fly nervous system is that the same pathways known to control the animal's gaze are also thought to be involved in flight control. The core functionality of the flight control system is therefore likely to also dependent on the 2 degrees of freedom controller that has been described earlier.



Figure 18: Control of gaze and flight by a 2 degrees of freedom controller. Upon externally caused body rotations, a fast feedforward (FF) signal initiates both, compensatory head movements ( $HR_c$ ) by activation of the neck motor (NM) system and compensatory body rotations ( $BR_c$ ) through the action of the flight motor (FM). This fast response mediated by the haltere system reduces the retinal slip speed and thus enables the processing of optic flow along the slow motion vision pathway generating a feedback signal (FB) that contributes to both,  $HR_c$  and  $BR_c$ . Any remaining misalignment between the head and the body may be measured by the prosternal and chordotonal organs providing slow feedback signals (FB, grey) primarily to the neck motor system. (unpublished work in progress).

The control diagram in Figure 18 only considers inner-loop reflexes induced by external perturbations. During voluntary movements of body to change the trajectory, motor commands are sent to both the neck and the flight motor, and efference copies may be sent to the sensors supporting the feedforward and the feedback degrees of freedom so that the sensory signal generated as a result of voluntary action does not immediately cause a counteracting stabilization reflex. Similar schemes in spirit have been proposed earlier (R3) and Chan et al 1998. But firstly, they did not include a 2 degree of freedom controller. And secondly, they are only weakly supported by experimental evidence. Further closed-loop simulations including the scheme will

have to show (i) its viability and (ii) may inform the design of further experimental studies in its support or to falsify it.

In a recent study that has been accepted for publication at a control engineering conference we took a very different approach: We propose a novel mathematical framework based on an extended Kalman filter and a controller with infinite-horizon dynamics that minimises the costs associated with muscle contraction on the one hand side and imperfect gaze stabilization on the other (C9). In closed-loop simulations the model explicitly considers inherent constraints in biological systems, i.e. ambiguity and noisiness of sensor signals, inevitable response delays along sensory pathways, and limited energy supply for both the neck motor and the nervous system. Figure 19 shows the block diagram of the system.



Figure 19: Gaze stabilization system based on a Kalman filter-type state estimator and controller providing а optimized sensoy feedback to the motor plant. The scheme involved the motion vision pathway provided by the compound eves, halteres, ocelli and neck proprioceptors which measure external disturbances,  $dX_k^{ext}$ . Small deviations of from a horizontal head orientation are tolerated due to the cost with muscle associated contraction. (from C9)

Although it is unlikely that flies have evolved an implementation of an optimized Kalman filter in their nervous system the proposed framework may as well be of interests for technical applications including autonomous micro air vehicles.

# **3.** Conclusion:

Within the period of funding we have made significant progress in all areas of research carried out in my lab and on the basis of collaborations. The collaboration with Sean Humbert turned out to be very successful with us meeting at least once or twice a year, being in regular email contract and talking on the phone whenever some immediate discussion on our collaborative work was required. We published together on 3 occasions and are still working together on a first paper where the modes of motion in *Calliphora* are related to the preferred rotation axes of its VS-cell in addition to another conference publication on a bioinspired ocellar sensor.

We also engaged on another grant proposal to AFOSR/EOARD together with Graham Taylor on the 'Mode Sensing Hypothesis' which is a key element of the implementation road map of the AFRL-DSTL programme arrangement on bio-inspired technologies. The start of the grant had been significantly delayed due to the financial situation in the US. Meanwhile funding for the first year has been set up at the University of Maryland.

One aspect regarding the scientific programme of this grant that had been underestimated was the impact of limited availability of dipteran species which cannot be bred in the lab. Although we developed strategies to acquire other species by means of wild catches and temporary breeding of hoverflies except for *Calliphora* we were quite often short of experimental animals. This was, however, compensated for by many complementary approaches and alternative aspects sensorimotor research so that a constant stream of high profile output was produced in my lab during the funding period.

Another limiting factor certainly is a comparatively small size of my research group. I mentioned in my previous reports already that the comprehensive approach I apply to insect sensorimotor control at four different system levels and across several species would definitely benefit from a group size twice as large as my own. This, however, would require me to write way more grant applications with I would like to do, but cannot because of a limited time budget.

In an ideal scientific world and to facilitate research along the merits of AFRL and AFOSR stronger support in terms of finances would be much appreciated. Having said this, it is very clear that I would not have been able to engage on a more comprehensive approach on sensimotor control without the continuous funding stream provided by AFRL/AFOSR – normally through EOARD.

Irrespective the limitations mentioned so far, we have made excellent progress in particular with work on the 3- and 4-dimensional functional characterization of the neck and flight motor systems, work that was initiated originally by Daniel Schwyn and Martina Wicklein in my group. The work on state-dependent processing which is highly relevant to the topic of energy efficient information processing was also initiated in my lab by Kit Longden and has developed into an aspect several world-leading labs are now working on. And finally, the quantitative behavioural work and modelling projects in my lab and in collaboration with others scientists has reached a continuous level of comparatively high productivity.

Holger G Krapp

London, 16 March 2014

## 4. Additional References:

Chan WP, Prete F, Dickinson MH (1998) Visual input to the fly efferent control system of the fly's gyrocope. Sience 280:289-292

Hengstenberg, R (1991) Gaze control in the blowfly *Calliphora*: a multisensory, two-stage integration process," *Seminars in the Neurosciences*, 3:19–29

Lindemann JP, Kern R, van Hateren JH, Ritter H, Egelhaaf M (2005) On the computations analyzing natural optic flow: quantitative model analysis of the blowfly motion vision pathway. J Neurosci 25:6435-48

# 5. List of Publications:

## Diploma Thesis:

**Krapp H.**: Ein Verfahren zur schnellen Bestimmung der lokalen Vorzugsrichtung und Bewegungsempfindlichkeit im rezeptiven Feld von visuellen Interneuronen der Schmeißfliege *Calliphora erythrocephala*. Diplomarbeit. Eberhard-Karls-Universität Tübingen 1992.

## **PhD Thesis:**

**Krapp H.**: Repräsentation von Eigenbewegungen der Schmeißfliege *Calliphora erythrocephala* in VS-Neuronen des dritten visuellen Neuropils. Dissertation, Eberhard-Karls-Universität, Tübingen, 1995.

## Publications in refereed journals:

- O1. **Krapp H.G.** and Hengstenberg R.: Estimation of self-motion by optic flow processing in single visual interneurons. *Nature*, **384**: 463-466, (1996)
- O2. **Krapp H.G.** and Hengstenberg R.: A fast stimulus procedure for determining local receptive-field properties of motion-sensitive visual interneurons. *Vision Research*, **37**: 225-234, (1997)
- O3. **Krapp H.G.**, Hengstenberg B., and Hengstenberg R.: Dendritic structure and receptive field organization of optic flow processing interneurons in the fly. *Journal of Neurophysiology*, **79**: 1902-1917, (1998)
- O4. Gabbiani F., **Krapp H.G.**, and Laurent G.: Computation of object approach by a widefield, motion- sensitve neuron. *Journal of Neuroscience*, **19**: 1122-1141, (1999)
- O5. Gabbiani F, Laurent G, Hatsopoulos N, and **Krapp H.G.**: The many ways of building collision-sensitive neurons. *TINS*: **22**, 437-438, (1999)
- O6. **Krapp H.G.** Neuronal matched filters for optic flow processing in flying insects. *International Review of Neurobiology*, **44**: 93-120, (2000)
- O7. Franz M.O. and **Krapp H.G.**: Wide-field, motion-sensitive neurons and optimal matched filters for optic flow. *Biological Cybernetics*, **83**: 185-197, (2000).

- O8. Petrowitz R., Dahmen H., Egelhaaf M., and **Krapp H.G.**: Arrangement of optical axes and the spatial resolution in the compound eye of the female blowfly. *Journal of Comparative Physiology A*: **186**: 737-746, (2000).
- O9. Karmeier K., Egelhaaf, M., and **Krapp H.G.**: Early visual experience and the receptive field organization of optic flow processing interneurons in the fly motion pathway. *Visual Neuroscience*, **18**: 1-8, (2001).
- O10. Krapp H.G., Hengstenberg R., and Egelhaaf M.: Binocular input organization of optic flow processing interneurons in the fly visual system. *Journal of Neurophysiology*, 85: 724-734, (2001).
- O11. Gabbiani F., **Krapp H.G.**, Koch C., and Laurent G.: Multiplicative computation in a looming sensitive neuron. *Nature*, **420**: 320-324, (2002).
- O12. Egelhaaf M., Kern R., **Krapp H.G.**, Kretzberg J., Kurtz R., and Warzecha A-K.: Neuronal encoding of behaviourally relevant visual-motion information in the fly. *TINS*: **25**, 94-100, (2002).
- O13. Karmeier K., **Krapp H.G.**, and Egelhaaf M.: Robustness of the tuning of fly visual interneurons to rotatory optic flow. *Journal of Neurophysiology*, **90**: 1626-1634 (2003).
- O14. Matheson T., **Krapp H.G.**, and Rogers S.M.: Plasticity in the visual System is correlated with a change in life-style of solitarious and gregarious locusts. *Journal of Neurophysiology*, **91**: 1-12 (2004).
- O15. Franz MO, Chahl JS, and **Krapp H.G.:** Insect inspired estimation of self-motion. *Neural Computation* **16**: 2245-2260, (2004).
- O16. Gabbiani F, **Krapp H.G.**, Hatsopoulos N., Mo C-H., Koch C. and Laurent G.: Multiplication and stimulus invariance in a looming-sensitive neuron. *Journal of Physiology Paris* **98**, 19-34. (2004).
- O17. **Krapp H.G.** and Gabbiani F.: Spatial distribution of inputs and local receptive field properties of a wide-field, looming sensitive neuron. *Journal of Neurophysiology*, **93**: 2240 2253, (2005).
- O18. Karmeier K., Krapp H.G., and Egelhaaf M: Population coding of self-motion: Applying Baysian inference to estimate the performance of visual interneurons in the fly. *Journal* of Neurophysiology, 94: 2182 – 2194, (2005).
- O19. Gabbiani F. and Krapp H.G.: Spike-frequency adaptation and intrinsic response properties of an identified, looming sensitive neuron. *Journal of Neurophysiology*, 96, 2951-2962, (2006).
- O20. Parsons M.M., **Krapp H.G**., and Simon B. Laughlin: A motion sensitive neuron responds to signals from the two visual systems of the blowfly, the compound eyes and the ocelli. *Journal of Experimental Biology*, **209**, 4464 4474, (2006).
- O21. Peron S., **Krapp H.G.**, and Gabbiani F.: Influence of electrotonic structure and synaptic mapping on the receptive field properties of a collision detection neuron. *Journal of Neurophysiology*, **97**, 159 177, (2007).
- O22. Rogers S.M., Krapp H.G., Burrows M., and Matheson T.: Compensatory plasticity at an identified pathway tunes a viso-motor pathway. *Journal of Neuroscience*, 27, 4621 – 4633, (2007).
- O23. **Krapp H.G.:** Polarization vision: How insects find their way by watching the sky. *Current Biology*, **17**, R557 R560 (2007).

- O24. Taylor G.M. and **Krapp H.G.**: Sensory systems and flight stability: What do insects measure and why? *Advances in Insect Physiology*, **34**, 231 316 (2007).
- O25. **Krapp H.G.**: Estimation of self-motion for flight and gaze stabilization. *Navigation*, **55**, 147-158 (2008).
- O26. Huston S.J. and Krapp H.G.: Visuomotor transformation in the fly gaze stabilization system. <u>PLoS Biology</u>, 6(7):e173 (2008).
- O27. Saleem A, **Krapp H.G.**, and Schultz S.R.: Spike-triggered independent component analysis: application to a fly motion-sensitive neuron. *Visual Neuroscience*, **8**(13):2, 1-16 (2008).
- O28. **Krapp H.G.**: Sensory Integration: Neuronal adaptations for robust visual self-motion adaptation. *Current Biology*, **19**(10), R413-416 (2009).
- O29. Krapp H.G.: Ocelli, Current Biology, 19(11), R435-437 (2009).
- O30. Huston S.J. and **Krapp H.G.**: Non-linear integration of visual and haltere inputs in fly neck motor neurons, *Journal of Neuroscience*, **29**,13097-13105 (2009).
- O31. Longden K.D. and **Krapp H.G.**: State-dependent receptive field properties of optic flow processing interneurons, *Journal of Neurophysiology*, **102**, 3606-3618 (2009).
- O32. Rogers S.M., Harston G., Kilburn-Toppin F., Matheson T., Burrows M., Gabbiani F., and **Krapp H.G.**: Differences in the spatiotemporal receptive field of a looming-sensitive neuron caused by phase change in the Desert Locust. *Journal of Neurophysiology*, **103**, 779-792 (2010).
- O33. **Krapp H.G.**: Sensorimotor Transformation: From visual responses to motor commands. *Current Biology*, **20**(5), R236-R239, (2010).
- O34. Parsons M.M., **Krapp, H.G.**, and Laughlin S.B.: Sensor fusion in identified visual interneurons. *Current Biology* **,20**(7), 624-628 (2010).
- O35. Hyslop A., **Krapp H.G.**, and Humbert J.S.: A control theoretic interpretation of directional motion preferences in optic flow processing interneurons. *Biological Cybernetic*, **103**(5), 353-364, DOI: 10.1007/s00422-010-0404-8 (2010).
- O36. Longden K.D. and **Krapp H.G.**: Octopaminergic modulation of temporal frequency coding in an identified optic flow-processing interneuron. *Frontiers in Systems Neuroscience*. **4**, 153. doi: 10.3389/fnsys.2010.00153 (2010).
- O37. Ejaz N., Peterson K.D., **Krapp H.G.**: An Experimental Platform to Study the Closedloop Performance of Brain-machine Interfaces. *Journal of Visualized Experiments*, **49**, <u>http://www.jove.com/index/</u>Details.stp?I D=1677, doi: 10.3791/1677 (2011).
- O38. Longden K.D. and **Krapp H.G.**: Sensory Neurophysiology: Motion vision during motor action. *Current Biology*, **21**, R650-R652 (2011).
- O39. Saleem A.B., Longden K.D., Schwyn D., Krapp H.G., Schultz S.R.: Bimodal optomotor response to plaids in blowflies: mechanisms of component selectivity and evidence for pattern selectivity. *Journal of Neuroscience*, **32**, 1634-1642 (2012).
- O40. Ejaz N., Krapp H. G. , and Tanaka R.J.: Closed-loop response properties of a visual interneuron involved in fly optomotor control. *Frontiers in Neural Circuits*, **7**, (2013) | doi: 10.3389/fncir.2013.00050
- O41. Schwyn, D.A., Moksoc, R., Walker S.M., Doube M., Wicklein M., Taylor G.K., Stampanoni M., **Krapp H.G.**: High speed x-ray imaging on the fly. *Synchrotron Radiation News*, **26**(2), 4-10, (2013) DOI:10.1080/08940886.2013.771064

- O42. Yue, XC, **Krapp, H. G.**, Drakakis, E. M.: An output code offset-free comparator for SAR ADCs based on non-linear preamplifier and CMOS inverters. *Microelectronics Journal*, **44**(5), 414-420, DOI: 10.1016/j.mejo.2013.02.011
- O43. Mokso, R., Marone, F., Irvine, S., Nyvlt, M., Schwyn, D., Mader, K., Taylor, G. K., Krapp, H. G., Skeren, M., Stampanoni, M.: Advantages of phase retrieval for fast x-ray tomographic microscopy. *Journal of Physics D-Appield Physics*, 46(49) DOI: 10.1088/0022-3727/46/49/494004
- O44. Walker S.M., Schwyn D.A., Mokso R., Wicklein M., Müller T., Doube M., Stampanoni M., Krapp H.G., Taylor G.K.,: In Vivo Time-Resolved Microtomography Reveals the Mechanics of the Blowfly Flight Motor. *PLoS Biology*, in press (2014)
- O45. Longden K.D., Muzzo T., Cook D., Schultz S.R., and **Krapp H.G.**: Nutritional state modulates the neural processing of visual motion. *Current Biology*, in press (2014)

#### **Refereed Conference Proceedings:**

- C1. **Krapp H.G.** Optic flow processing for motor control in flying insects. In: Proceedings of the 60<sup>th</sup> Annual Meeting of the Institute of Navigation, Dayton, Ohio, <u>www.ion.org</u>. (2004)
- C2. **Krapp H.G.**: Estimation and control of self-motion and gaze in flying insects. In: Proceedings of the 63<sup>th</sup> Annual Meeting of the Institute of Navigation, Cambridge, MA, <u>www.ion.org</u>. (2007)
- C3. Yue X., Peterson K., **Krapp H.G.**, and Drakakis E.M.: "A Low-Power Low-Distortion Amplifier for Fly Neural Recordings". 4th International Conference on Bioinformatics and Biomedical Engineering (iCBBE), (oral presentation), Chengdu, China. Proceedings IEEE, Ei Compendex, ISTP (2010).
- C4. Schwyn, D., Hernandes Heras F.J., Bolliger, G., **Krapp H.G.**, Tanaka, RI.: Characterization of insect gaze control systems. *18<sup>th</sup> World Congress of International Federation of Automated Control (IFAC)*, Milan, (2011)
- C5. Ejaz, N., Tanaka, R.J., Krapp, H.G.: Closed-loop performance of a proportional controller for visual stabilization using a fly-robot interface. In: IEEE Int. Conf. Rob. Biomim., pp. 1509–1515 (2011).
- C6. Ejaz, N., Tanaka, R.J., Krapp, HG.: Static versus adaptive gain control strategy for visiomotor stabilization. In: Biomimetic and Biohybrid Systems, Springer, 7375: pp. 107-119 (2012).
- C7. Gremillion G., Galfond M., Krapp H.G., Humbert S.J.: Biomimetic sensing and modeling of the ocelli visual system of flying insects. In: IEEE Int. Conf. Intellegent Robots and Systems (IROS), (2012).
- C8. Huang J.V. and Krapp H.G.: Miniaturized electrophysiology platform for fly-robot interface to study multisensory integration. In: Biomimetic and Biohybrid Systems, Springer, LANI 8064: pp. 119-130 (2013).
- C9. Sabatier Q, **Krapp H.G.**, Tanaka R.: Dynamic optimisation for fly gaze stabilisation based on noisy and delayed sensor information. European Control Conference, ECC, 2014, (accepted)

## **Book Chapters:**

- R1. Hengstenberg R., Krapp H., and Hengstenberg B.: Visual sensation of self-motions in the blowfly *Calliphora*. *In*: C. Taddei-Ferretti (ed.) Biocybernetics of Vision: Integrative and Cognitive Processes. Singapore: World Scientific, pp. 41-52, (1998).
- R2. Dahmen H., Franz M.O., and Krapp H.G.: Extracting ego-motion from optic flow: limits of accuracy and neuronal filters. *In:* (J. M. Zanker and J. Zeil, eds.) Processing visual motion in the real world -- A survey of computational, neural and ecological constraints. Berlin, Heidelberg, New York: Springer Verlag, pp. 143-168, (2001).
- R3. **Krapp H.G.** and Wicklein M.: Central processing of visual information in insects. In: Basbaum A.I, Kaneko A., Shepherd G.M., and Westheimer G. (eds) The Senses: A comprehensive reference. Vol. 1, Vision I, Masland R and Albright T.D. (eds.), San Diego, Academic Press, pp. 131-204, (2008)
- R4. **Krapp H.G.**, Taylor G.K., and Humbert S.J.: The mode-sensing hypothesis: matching sensors, actuators and flight dynamics. In: Frontiers in Sensing Biology and Engineering. Srinivasan MV and Barth F (eds.), Springer (2012)
- R5. **Krapp H.G.**: Flies, optic flow, and multisensory stabilization reflexes. In: Flow Sensing in Air and Water. Bleckmann H. et al. (eds.), Springer, Berlin, Heidelberg, pp. 215-243, (2014) DOI: 10.1007/978-3-642-41446-6\_9
- R6. **Krapp H.G.**: Optic flow processing. In: Enzyclopedia of Computational Neuroscience. Jaeger D. and Jung R (eds.), Springer, Berlin, Heidelberg, Article ID: 348397 · Chapter ID: 332 (accepted)

Manuscripts submitted/in preparation:

O46. Schwyn D.A., Mokso R., Walker S.M., Doube M., Wicklein M., Müller T., Stampanoni M., Taylor G.K., **Krapp H.G.:** Four-dimensional in-vivo x-ray micro-imaging of the internal kinematics of insect flight.

O47. Wicklein M., Schwyn D.A., Simonsen T.J., Howard L.E., Ball A., Abel R.L., and **Krapp H.G.**: Iodine staining for global and tissue specific contrast enhancement in microcomputed tomography.

O48. Swart P et al.: Micro CT preparation for imaging soft tissue in *Calliphora vicina*.

O49. Schwyn D.A., Hernandes Heras F.J., Bolliger G., Parsons M.M., Laughlin S.B., Tanaka R.J., and **Krapp H.G.**: Two-degree-of-freedom control of fly compensatory head roll.

### **Conference abstracts** (reviewed & non-reviewed):

- A1. **Krapp H.** and Hengstenberg R.: Reliability of a fast method to determine locally the preferred direction of motion sensitive neurons. In: Rhythmogenesis in Neurons and Networks. Eds. N. Elsner, D.W. Richter, Thieme, Stuttgart 1992, 306.
- A2. **Krapp H.** and Hengstenberg R.: Representation of specific optical flow fields in lobula plate neurons of the blowfly *Calliphora*. In: Gene Brain Behaviour. Eds. N. Elsner, M. Heisenberg, Thieme, Stuttgart 1993, 357.
- A3. **Krapp H.**, Hengstenberg B., and Hengstenberg R.: Correspondence of dendritic field structure, receptive field organization, and specific flow patterns in visual interneurons of the blowfly *Calliphora*. In: Proceedings of the 22<sup>nd</sup> Göttingen Neurobiology Conference, Vol. II. Eds. N. Elsner, H. Breer, Thieme, Stuttgart 1994, 447.

- A4. Krapp H. and Hengstenberg R.: Comparison between optic flow fields and response fields of visual interneurons in the lobula plate of the blowfly *Calliphora*. In: Proceedings of the 23<sup>nd</sup> Göttingen Neurobiology Conference, Eds. N. Elsner, R. Menzel, Thieme, Stuttgart 1995, 404.
- A5. Krapp H. and Hengstenberg R.: Filter neurons for specific optic flow patterns in the fly's visual system. In: Proceedings of the 4<sup>th</sup> International Congress of Neuroethology, Eds. M. Burrows, T. Matheson, P.L. Newland and H. Schuppe, Thieme, Stuttgart 1995, 264.
- A6. Hengstenberg R. and Krapp H.: Distribution of roll motion sensitivity in the eyes of *Callophora*: a comparison between neurons and behaviour. In: Proceedings of the 24<sup>th</sup> Göttingen Neurobiology Conference, Eds. N. Elsner, H.- U. Schnitzler, Thieme, Stuttgart 1996, 349.
- A7. Gabbiani F., Krapp H., Koch C. and Laurent G: Biophysical implementation of a multiplication operation in an interneuron of the locust. Soc. Neurosci. Abstr. 23: 613.17, 1997.
- A8. Krapp H.G., Gabbiani F., Koch, C. and Laurent G.: Neuronal Multiplication in the locust visual system. In: Proceeding of the 26<sup>th</sup> Göttingen Neurobiology Conference, Eds. N. Elsner, R. Wehner, Thieme, Stuttgart 1998, 410.
- A9. Franz M.O., Hengstenberg R. and Krapp H.G.: VS-neurons as matched filters for selfmotion-induced optic flow fields. In: Proceeding of the 26<sup>th</sup> Göttingen Neurobiology Conference, Eds. N. Elsner, R. Wehner, Thieme, Stuttgart 1998, 419.
- A10. Gabbiani F., **Krapp H.**, Koch C. and Laurent G: Computation of object approach by two wide field motion sensitive neurons. In: Soc. Neurosci. Abstr. **24**: 613.17, 1998.
- A11. Petrowitz R., Krapp H.G., Dahmen H. and Egelhaaf M.: The distribution of ommatidial axes in *Calliphora erythrocephala* varies with the direction of view. In: Proceeding of the 27<sup>th</sup> Göttingen Neurobiology Conference, Eds. N. Elsner, U. Eysel, Thieme, Stuttgart 1999, 439.
- A12. Krapp H.G. and Egelhaaf M.: Local preferred directions of visual wide field neurons and the compound eye geometry of the blowfly *Calliphora erythrocephala*. In: Proceeding of the 27<sup>th</sup> Göttingen Neurobiology Conference, Eds. N. Elsner, U. Eysel, Thieme, Stuttgart 1999, 440.
- A13. Karmeier K., Krapp H.G. and Egelhaaf M.: Does the receptive field organization of a blowfly visual interneuron depend on visual experience? In: Proceeding of the 27<sup>th</sup> Göttingen Neurobiology Conference, Eds. N. Elsner, U. Eysel, Thieme, Stuttgart 1999, 441.
- A14. **Krapp H.G.** and Gabbiani F.: Spatial integration of local motion stimuli by a widefield looming sensitive neuron. In: Soc. Neurosci. Abstr. **26**: 66.1, 2000.
- A15. Egelhaaf M., Petrowitz R., Dahmen H. and Krapp H.G.: Adaptation of the compound eye geometry to optic flow processing in the blowfly *Calliphora*. In: Soc. Neurosci. Abstr. 26: 368.11, 2000.
- A16. Gabbiani, F., Krapp, H.G., Koch, C, et al.: Multiplicative computation by a loomingsensitive neuron, 24th Annual International Conference of the Engineering-in-Medicineand-Biology-Society/Annual Fall Meeting of the Biomedical-Engineering-Society (EMBS 2002 BMES), pp, 1968 – 1969, 2002.
- A17. Matheson T., Basu Roy R., Copeland M., Krapp H.G. and Rogers S.M.: Plasticity in the visual system of solitarious and gregarious locusts. *Journal of Physiology* 547 P, p91PC, 2003.

- A18. Matheson T., **Krapp H.G.** and Rogers S.M.: Neuronal plasticity in the visual system of solitarious and gregarious locusts. *Comparative Biochemistry and Physiology A* **134** (3, suppl), S 73, 2003.
- A19. Matheson T., Krapp H.G. and Rogers S.M.: Adaptation to lifestyle in the visual system of solitarious and gregarious locusts. In: Proceedings of the 29<sup>th</sup> Göttingen Neurobiology Conference, Eds. N. Elsner, H. Zimmermann, Thieme, Stuttgart 2003, 483.
- A20. Karmeier K., Krapp H.G. and Egelhaaf M.: Population coding in the visual system of the blowfly: an experimental and modelling approach. In: Proceedings of the 29<sup>th</sup> Göttingen Neurobiology Conference, Eds. N. Elsner, H. Zimmermann, Thieme, Stuttgart 2003, 478.
- A21. Huston S.J. and Krapp H.G.: The visual receptive field of a fly neck motor neuron. In: Proceedings of the 29<sup>th</sup> Göttingen Neurobiology Conference, Eds. N. Elsner, H. Zimmermann, Thieme, Stuttgart 2003, 483.
- A22. Peron S.P., Krapp H.G., Laurent G., and Gabbiani F.: Role of dendritic morphology in the processing of looming stimuli: a compartmental modelling study. Program No. 491.3.
   2003 Abstract viewer/Itinerary Planner. Washington, DC: Society for Neuroscience, 2003. Online.
- A23. Card G, **Krapp H.G.**, and Laughlin SB: Motion adaptation does not affect the direction tuning of a neuron processing optic flow in the blowfly *Calliphora vicina*. In: University of Cambridge (2004) *Journal of Physiology* **555P** PC129, (2004)
- A24. Rogers SM, Harston GWJ, Kilburn-Toppin F, Matheson T, and **Krapp H.G.**: Receptive field organisation of a collision-detecting visual interneurone in solitarious and gregarious locusts. In: University of Cambridge (2004) *Journal of Physiology* **555P** PC130, (2004)
- A25. Matheson T, Rogers S, Kilburn-Toppin F, Harston G, and **Krapp H.G.**: Receptive field organization of a collision-detecting visual interneurone in solitarious and gregarious locusts. In: Proceedings of the 7<sup>th</sup> International Congress of Neuroethology, PO167, (2004)
- A26. Huston SJ and Krapp H.G.: Visual receptive field organization of sensory and motor neurons in the blowfly. In: Proceedings of the 7<sup>th</sup> International Congress of Neuroethology, PO168, (2004)
- A27. Gabbiani F and Krapp H.G.: Receptive field organization of a wide-field, looming sensitive neuron. In: Proceedings of the 7<sup>th</sup> International Congress of Neuroethology, PO182, (2004)
- A28. Gabbiani F and **Krapp H.G.**: Structure and receptive field input organization of a wide-field, looming sensitive neuron. In: Program No. 870.20. 2004 Abstract viewer/Itinerary Planner. Washington, DC: Society for Neuroscience, (2004). Online.
- A29. Karmeier K., Krapp H.G., and Egelhaaf M.: Population coding of self-motion in the blowfly visual system. In: Proceedings of the 30<sup>th</sup> Goettingen Neurobiology Conference. (2005).
- A30. **Krapp H.G.** and Huston S.J.: Encoding self-motion: From visual receptive fields to motor neuron response maps. In: Proceeding of the 30<sup>th</sup> Goettingen Neurobiology Conference. (2005).
- A31. Wüstenberg D. and **Krapp H.G.**: Self-motion estimation and flight control in blowflies and locusts: a comparative study. In: Proceeding of the 31<sup>st</sup> Goettingen Neurobiology Conference (2007).

- A32. Parsons M.M., **Krapp H.G.**, and Laughlin S.B.: Multisensory integration: Rotation detection in the visual system of the fly. In: Proceeding of the 31<sup>st</sup> Goettingen Neurobiology Conference (2007).
- A33. Huston S.J. and **Krapp H.G.**: Integration of visual and haltere inputs in fly neck motor neurons. In: Proceedings of the 8<sup>th</sup> International Congress of Neuroethology, (2007)
- A34. Longden K.D. and **Krapp H.G.**: Activity modulation of optic flow processing neurons by an octopamine agonist in the blowfly. In: Proceedings of the 8<sup>th</sup> International Congress of Neuroethology, (2007)
- A35. Rogers S.M., **Krapp H.G.**, and Matheson T.: Compensatory plasticiy at an identified synapse tunes a visuo-motor pathway in the locust. In: Proceedings of the 8<sup>th</sup> International Congress of Neuroethology, (2007).
- A36. Wüstenberg D. and **Krapp H.G.**: Optic flow processing interneurons in flies and locusts. In: Proceedings of the 8<sup>th</sup> International Congress of Neuroethology, (2007).
- A37. Saleem, A, **Krapp, HG**, Schultz, SR, Spike-triggered independent component analysis: A method for receptive field characterization. Society for Neuroscience, Page: 514.10., (2008).
- A38. Longden K.D. and **Krapp H.G.**: State-dependent coding properties of optic-flow processing cells in the blowfly visual system. Society for Neuroscience, Page: ., (2008).
- A39. Longden K.D. and **Krapp H.G.** Modulation of visual information processing in blowfly Lobula Plate Tangential Cells by an octopamine agonist. In: Proceeding of the 32<sup>nd</sup> Goettingen Neurobiology Conference (2009).
- A40. Ejaz N., Peterson K., and **Krapp H.G.** (2009) A robotic platform to study closed-loop optomotor control in the blowfly. In: Proceeding of the 32<sup>nd</sup> Goettingen Neurobiology Conference (2009).
- A41. Parsons M.M., **Krapp H.G.**, and Laughlin S.B. (2009) Convergence of compound eye and ocellar signals in Lobula Plate Tangential Cells of the blowfly *Calliphora*. In: Proceeding of the 32<sup>nd</sup> Goettingen Neurobiology Conference (2009).
- A42. Longden K.D. and **Krapp H.G.** (2009) Octopaminergic modulation of velocity coding in a blowfly optic flow-processing interneuron. Applied Vision Association Christmas Meeting, Bristol, UK (2009).
- A43. Longden K.D., and **Krapp H.G.** (2010) Velocity coding and octopamine in an identified optic flow-processing interneuron of the blowfly. Computational and Systems Neuroscience (COSYNE), 258, Salt Lake City, USA, (2010).
- A44. Longden K.D., Saleem A.B., Schultz S.R., and Krapp H.G. (2010) Responses to pattern motion of compound stimuli in an identified visual interneuron of the blowfly. 7<sup>th</sup> Forum of European Neuroscience (FENS), Amsterdam, Holland (2010).
- A45. Wicklein M., An J., Selmon A., and **Krapp H.G.** (2010) Comparing the receptive field organization of LPTCs in male and female blowflies. 7th Forum of European Neuroscience (FENS), Amsterdam, Holland (2010).
- A46. Longden K.D., Saleem A.B., Schwyn D., Schultz S.R., and **Krapp H.G.** (2010) How blowflies analyse the motion of visual patterns. <sup>9th</sup> International Congress of Neuroethology, Salamanca, Spain (2010).
- A47. Schwyn D.A., Wicklein M., Klosowski M.M., Doube M., Abel R., and **Krapp H.G.** (2010) Imaging the fly brain using micro-CT. <sup>9th</sup> International Congress of Neuroethology, Salamanca, Spain, (2010).

- A48. Wicklein M., Schwyn D.A., Abel R.L., Simonsen T. J., and **Krapp H.G.** (2011) Optic lobes in male and female blowflies determined by micro-CT imaging. Ninth Goettingen Meeting of the German Neuroscience Society, Goettingen, Germany, (2011).
- A49. Schwyn D.A., Hernández Heras F.J., Bolliger G., Parsons M.M., **Krapp H.G.**, and Tanaka R.J. (2011) Behavioural studies and modelling of gaze stabilization in the blowfly *Calliphora*. Ninth Goettingen Meeting of the German Neuroscience Society, Goettingen, Germany, (2011).
- A50. Hardcastle B.J., Schwyn D.A., Longden K.D., Bierig K., Tanaka R.J. and Krapp H.G. (2012). Ocellar contributions to gaze control: a behavioural linear systems analysis in the blowfly. *Front. Behav. Neurosci. Conference Abstract: Tenth International Congress of Neuroethology*. doi:10.3389/conf.fnbeh.2012.27.00414, University of Maryland, USA, (2012).
- A51. Longden K.D. and **Krapp H.G.** (2012) State-dependent motion vision in walking blowflies. *Tenth International Congress of Neuroethology* (oral presentation), University of Maryland, USA, (2012).
- A52. Schwyn D.A., Walker S.M., Mokso R., Wicklein M., Doube M., Stampanoni M., Taylor G.K., and Krapp H.G. (2012). Ultra-fast in vivo x-ray microtomography of blowfly wingbeat. Front. Behav. Neurosci. Conference Abstract: Tenth International Congress of Neuroethology. doi: 10.3389/conf.fnbeh.2012.27.00407, University of Maryland, USA, (2012).
- A53. Wicklein M., Schwyn D.A., Simonsen T.J., Howard L.E., Ball A., Abel R.L., and Krapp H.G. (2012) Radiocontrast micro-CT imaging optimised for differential tissue segmentation in *Calliphora vicina* (blowfly). 15<sup>th</sup> European Microscopy Congress, Manchester, UK, (2012).
- A54. Schwyn D.A., Mokso R., Walker S.M., Wicklein M., Doube M., Müller T., Taylor G.K., Stampanoni M., Krapp H.G. (2012) Ultra-fast x-ray microtomography of wing beat in tethered flies. 15<sup>th</sup> European Microscopy Congress (oral presentation), Manchester, UK, (2012).

## **Edited Books:**

- 1. **Krapp H.** and Wägenbaur T. (Eds.) Künstliche Paradiese Virtuelle Realitäten. München: Fink-Verlag, 1997, ISBN: 3-7705-3252-8
- 2. **Krapp H.** and Wägenbaur T. (Eds.) Komplexität und Selbstorganisation Chaos in den Kultur- und Naturwissenschaften. München: Fink-Verlag, 1997, ISBN: 3-7705-3252-X
- Lepora, N.F., Mura, A., Krapp, H.G., Verschure, P.F.M.J., Prescott, T.J. (Eds.) Living Machines 2013. Proceedings of the Second International Conference on Biomimetic and Biohybrid Systems. Berlin, Heidelberg: Springer-Verlag, 2013, ISBN: 978-3-642-39801-8