

# **THERMAL FRACTURING, UNDERWATER AMBIENT NOISE MEASUREMENTS AND MODELING**

Dr. Peter J. Stein and Dr. Subramaniam D. Rajan  
Scientific Solutions, Inc.  
18 Clinton Drive  
Hollis, NH 03049-6576  
pstein@mv.mv.com and rajan@gte.net  
Voice: (603) 880-3784 Fax: (603) 598-1803

Dr. James K. Lewis  
Ocean Physics Research & Development  
207 S. Seashore Ave.  
Long Beach, MS 39560  
ocnphys@aol.com  
Voice: (228) 865-0059 Fax: (228) 864-3268

Award No. N00014-97-C-0294

## **LONG TERM GOALS**

The long term goal is to understand the relationship between the environmental conditions, ice properties, ice geometry, and the thermal fracturing of ice and the resulting underwater ambient noise. By doing this we hope to determine the effect of thermal fracturing on the strength of sea ice and its possible influence on climate change.

## **OBJECTIVES**

Thermal fracturing occurs when rapid temperature changes, mainly cooling of the upper surface, stress the pack ice. Recent work conducted during the Sea Ice Mechanics Initiative (SIMI) has shown the importance of thermal fracturing. Figure 1 (top) shows data acquired during the CEAREX experiment. Plotted are the minimum and maximum underwater ambient noise levels in third octave bands. There are two separate driving mechanisms forcing the ambient noise. The low frequencies, below 100 Hz, are dominated by motion induced mechanisms. The higher frequencies are dominated by thermal fracturing activity. Thus, thermal fracturing is one of the primary source of the ambient noise which interferes with sonar systems. Strong thermal fracturing occurs one or more times per week. Work conducted during SIMI also indicates that thermal fracturing is one of the primary means of weakening the pack ice at the 100 m scale. A weakening of the pack ice implies an increase in the formation of open areas of water (leads), and hence greater heat flux between the ocean and the atmosphere. Thus, changes in the thermal fracturing rate may influence climate change in the Arctic.

Our objective is to formulate a complete model for thermal fracturing and the resulting ambient noise. Figure 1 (bottom) shows our most recent attempt at doing this. Inputs to the model are also given in Figure 1. We see good agreement in the example shown, however there are some regions of disagreement. Other data/model comparisons do not look as good. To

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>30 SEP 1997</b>		2. REPORT TYPE		3. DATES COVERED <b>00-00-1997 to 00-00-1997</b>	
4. TITLE AND SUBTITLE <b>Thermal Fracturing, Underwater Ambient Noise Measurements and Modeling</b>			5a. CONTRACT NUMBER		
			5b. GRANT NUMBER		
			5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S)			5d. PROJECT NUMBER		
			5e. TASK NUMBER		
			5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Scientific Solutions, Inc,18 Clinton Drive,Holis,NH,03049</b>			8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSOR/MONITOR'S ACRONYM(S)		
			11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>Same as Report (SAR)</b>	18. NUMBER OF PAGES <b>4</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

improve the models we need more long term ambient noise data sets where all the relevant meteorological data is also collected. This includes heat flux and snow cover data. A very complete long term meteorological data set will be collected during the NSF/ONR sponsored Surface Heat Budget of the Arctic (SHEBA) experiment. Our most recent objective has been to design, build, and deploy three autonomous measurement systems during the SHEBA experiment to collect the complimentary ambient noise data.

## **APPROACH**

The autonomous Arctic ambient noise measurement system is shown in Figure 2 prior to shipping. A watertight box houses the batteries, data acquisitions computer, signal conditioning electronics, and satellite telemetry gear. In the foreground of Figure 2 (small black object) is the hydrophone and 100 m of anti-strum cable. Mounted on a the pole is a GPS antenna, an ARGOS antenna, and four diode isolated solar panels. Four solar panels are used so one will always be facing the sun. The batteries are sized to make it through 6 months of darkness. Thus, the system should run indefinitely unless destroyed by the elements. The computer takes ambient noise data once per hour averaged over 20 seconds. The computer then calculates third octave bands and statistics of the noise sample. The ice stress is also measured once per hour. The third octave band levels, statistics, ice stress data, location, and time are telemetered back via ARGOS. In the SHEBA experiment, the main camp will be supported by a permanently stationed icebreaker frozen into the ice. To reduce contamination from ship noise, our systems will be deployed 50 km from the camp.

## **ACCOMPLISHMENTS**

Under this contract, we have designed and built three autonomous ambient noise measurement systems. The systems have been shipped to the Arctic and are being deployed as of this writing. If these systems work properly, this should give the most complete data set yet available for studying thermally induced stresses, fracturing, and underwater ambient noise.

## **IMPACT FOR SCIENCE/ SYSTEMS APPLICATIONS**

Over the past several years we have made great advances in understanding the thermal stressing of the ice, the resulting fracturing of the ice, and ambient noise. We will use the data acquired during SHEBA to further understand the fracturing mechanisms and to improve our models.


## **TRANSITIONS**

Eventually the models developed will be transitioned into operational models used by the Navy Fleet for predicting ambient noise and system performance. The models will also be incorporated in ice/ocean/atmosphere models for use in climate predictions studies. The autonomous ambient noise measurement system might also be transition into a multi-year monitoring of ambient noise. By monitoring the ambient noise we might be able to determine if there are any changes in the rate of thermal fracturing. This may yield clues as to the direction of climate change in the Arctic and the importance of thermal fracturing.


## **RELATED PROJECTS**

Scientific Solutions is also involved in developing a radiometer system for making remote unmanned radiometric measurements under harsh conditions. Eight advanced prototype units are

being deployed during the SHEBA experiment to make remote radiation measurements away from the camp and the ship contamination.



**Figure 1.** Modelled data comparison for thermal cracking noise.



**Figure 2.** Autonomous ice stress/ambient noise measurement system.