

The STRATAFORM Swathmapping Program

Larry Mayer
Ocean Mapping Group
Dept. of Geodesy and Geomatics Engineering
University of New Brunswick
Fredericton, N.B. Canada E3B 5A3
phone: (506) 453-3577 fax: (506) 453-4943 email: larry@omg.unb.ca
Award # N00014-95-0064

LONG-TERM GOALS:

To provide the fundamental mapping information necessary to support STRATAFORM's multidisciplinary effort to develop a more complete understanding of how short-term oceanographic and geological processes interact to produce the preserved geologic record on the shelf and slope portions of the continental margins..

SCIENTIFIC OBJECTIVES:

The fundamental objective of the STRATAFORM Swathmapping Program is **to provide complete (100%) bathymetric and sidescan imagery coverage of the Northern California and N.J. Margin STRATAFORM field areas**. This has allowed STRATAFORM investigators to evaluate the geologic processes of the shelf and slope over a continuum of scales. Complete coverage also has provided STRATAFORM investigators with the knowledge that their studies are based on a complete picture of morphological relationships rather than the interpolation of sparsely spaced data. In doing this, we have produced a bathymetric, geomorphological, and potentially lithological framework upon which all subsequent work can be built. Ancillary objectives are **to further develop techniques for the remote classification of seafloor materials from swathmapping data** and, **to develop techniques for the interactive 3D visualization of co-registered surficial and subbottom data**.

APPROACH:

We have chosen to use a state-of-the-art multibeam sonar system (EM-1000) for our mapping. As compared to conventional echosounders, multibeam sonars provide increased source level, lateral resolution, and a substantial increase in data density and areal coverage. Most importantly, the newer systems also provide the ability to simultaneously produce high-resolution sidescan sonar imagery. We have developed a full suite of real-time and near real-time multibeam sonar processing tools to assure that only high-quality data is collected and that this data can be processed in the field. These tools also allow for the interactive 3-D visualization of multibeam data sets and derivative products. We are also developing a range of seafloor classification algorithms with particular focus on techniques that look at the characteristics of the returned waveforms as well as the change in backscatter as a function of angle of incidence. The results of these analyses will be compared to ground truth data collected by other STRATAFORM scientists to explore the limits of extracting quantitative seafloor property information from multibeam sonar data.

WORK COMPLETED:

In 1995 and 1996, we completed mapping of the Calif. and most of the N.J. survey areas. In 1997 we completed the remaining deep-water portion of the N.J., processed these data and merged them with the shallow water data. We also processed additional multibeam (Hydrosweep) data from the Eureka area collected by Clark Alexander. We merged this data with the earlier data sets and made the new maps available to all STRATAFORM investigators. We have now focused our efforts on visualization and particularly on the question of remote seafloor classification. We have finalized interactive 3-D fly-throughs of the N. J. margin

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 1997		2. REPORT TYPE		3. DATES COVERED 00-00-1997 to 00-00-1997	
4. TITLE AND SUBTITLE The STRATAFORM Swathmapping Program				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of New Brunswick, Dept. of Geodesy and Geomatics Engineering, Fredericton, N.B. Canada E3B 5A3,				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 4	19a. NAME OF RESPONSIBLE PERSON
a REPORT unclassified	b ABSTRACT unclassified	c THIS PAGE unclassified			

and have developed a suite of automated algorithms for extracting and parameterizing the backscatter as a function of angle of incidence. We have completed a pilot study of the application of these algorithms on a well-ground truthed region of Stellwagen Bank and have now begun applying these algorithms to both STRATAFORM survey areas.

RESULTS:

The combined bathymetry and imagery of the N. J. margin show a spectacular picture of shelf processes with large scale ridge and swale features that appear to be moving across the shelf, a series of ablation features and, a large region near the shelf break that appears to be covered by old iceberg scours. The ridge and swale topography appears to be overriding the scour features though there are several large scour-like feature that also seem to cut into the ridge and swale deposits. If these large features are iceberg scours, this part of the system has been preserved since the last deglaciation. The deep water data provides direct insight into the origin of submarine canyons, with landward erosion apparently the dominant mechanism. Our parameterization of the angular backscatter response curve appears to be a sensitive indicator of changes in seafloor type or texture. In particular, the shape of the curve below 40° is especially revealing as response in the vicinity of the critical angle may be observed. The use of our 3-D visualization tools has allowed us to directly see the spatial distribution of our extracted classifiers and their relationship to seafloor morphology(Fig. 1).

IMPACT/APPLICATIONS:

The swathmapping results from the Eureka and the N. J. margins provide all STRATAFORM investigators with an unprecedented, detailed look at both the bathymetry and distribution of sediments on the shelf and slope. These results have already been used in planning the deployment of a series of long-term moorings, seismic profiling and coring cruises, as well as for planning ROV, submersible work, and ODP sites. Quantitative measurements of slope and sediment-type distributions will inevitably lead to improved models of shelf and slope development, the primary goal of the STRATAFORM program. Our initial results from the New Jersey margin indicate that pervasive ice scouring is evident further south than previously reported and may have impact on our overall understanding of the evolution of shelf stratigraphy.

TRANSITIONS:

Our maps and data have been used by numerous investigators both in and out of STRATAFORM, including several other Navy programs (e.g., SWARM, NRL). Our processing and visualization software is being used by NAVO, NRL, NOAA, USGS as well as several universities and private sector companies.

RELATED PROJECTS: NSERC Chair in Ocean Mapping

REFERENCES:

Brissette,M.B., Hughes Clarke,J.E., Bradford,J. and MacGowan,B., 1997, Detecting small seabed targets using a high frequency multibeam sonar: Geometric Models and Test Results: Proc. Oceans 97.

Dijkstra, S.J. and Mayer, L. A., 1996, "Lasso!: an interactive graphical tool for seafloor classification", Proc. IEEE-MTS Oceans '96, 1064-1070, 1996.

Du, Z, Wells, D. and Mayer, L. A., 1996, An Approach to Automatic Detection of Outliers in Multibeam Echo Sounding Data, The Hydrographic Journal, No. 79, p. 19-23.

Goff, J.A., C. L. Schuur, L. Mayer, J. E. Hughes Clarke High-resolution bathymetric survey of the New Jersey continental margin: Quantitative analysis of sand ridge morphology, EOS, v.77, no. 46., p. 329.. Dec. 1996

Goff, J.A., Mayer, L.A., Hughes Clarke, J.E., and Pratson, L.F., 1996, Swath mapping on the continental shelf and slope: the Eel River Basin, Northern California, *Oceanography*, v. 9, no. 3, p. 178- 182.

Hughes Clarke, J.E., Danforth, B.W. and Valentine, P., 1997, Areal seabed classification using backscatter angular response at 95 kHz, in Pace, N.G., Pouliquen, E., Bergem, O., and Lyons., eds., *High Frequency Acoustics in Shallow Water*, SACLANT Conference Proceedings Series, CP-45, NATO SACTLANT Research Centre, La Spezia, Italy, p. 300-312

Hughes Clarke, John E., Mayer, Larry A. and Wells, David E., 1996, Shallow-water imaging multibeam sonars: A new tool for investigating seafloor processes in the coastal zone and on the continental shelf. *Marine Geophysical Researches*, v. 18, n. 6, p. 607-629

Hughes Clarke, J.E., L.A. Mayer, J. A. Goff, H. Lee, W. Wakefield, 1996, Regional seabed characterisation through analysis of the angular response of seabed backscatter strength: application to STRATAFORM East and West Coast surveys. *EOS*, v.77, no. 46., p. 329., Dec. 1996

Mayer, L.A., Dijkstra, S., Hughes Clarke, J.E., Paton, M. and Ware, C., 1997, Interactive tools for the exploration and analysis of multibeam and other seafloor acoustic data, in Pace, N.G., Pouliquen, E., Bergem, O., and Lyons., eds., *High Frequency Acoustics in Shallow Water*, SACLANT Conference Proceedings Series, CP-45, NATO SACTLANT Research Centre, La Spezia, Italy, p. 355 - 362.

Mayer, L.A., Hughes Clarke, J.E., and Dijkstra, S., 1997, Multibeam Sonar: Applications in Fisheries Research, In: Boehlert, G.W. and Schumacher, J.D., eds., *Changing Oceans and Changing Fisheries: Environmental Data for Fisheries Research and Management*. NOAA Special Publication NOAA-TM-NMFS-SWFSC, pg. 79 - 92.

Mayer, L.A., J.E. Hughes Clarke, J. A. Goff, C. L. Schuur, D. J. P. Swift, 1996, Multibeam Sonar Bathymetry and Imagery from the New Jersey Continental Margin: Preliminary Results, *EOS*, v.77, no. 46., p. 329. Invited. Dec. 1996

Miller, J., Hughes Clarke, J.E. and Patterson, J., 1997, How Effectively Have You Covered Your Bottom?: *Hydrographic Journal*, no.83, p.3-10. http://www.omg.unb.ca/~jhc/coverage_paper.html

Paton, M., Mayer, L.A., and Ware, C., 1997, Interactive 3-D tools for pipeline route planning, *Proceedings of the IEEE-MTS Oceans 97*, pg. 1215 - 1222.

Schuur, C.L., J A Goff, J A Austin Jr, and C Fulthorpe, L A Mayer, J E Hughes Clarke, 1996, Quaternary geomorphology of the New Jersey continental shelf from acoustic swath mapping and high-resolution seismic reflection data, *EOS*, v.77, no. 46., p. 329. Dec. 1996

WebSite: <http://www.omg.unb.ca>




Fig 1. 3-D visualization of two of the twelve parameters extracted from the angular response of backscatter data from a 30 x 27 km area of Stellwagen Bank National Marine Sanctuary, Mass. Each is draped over bathymetry and displayed (in original) with color coding. Lower figure (A) represents slope of segment of backscatter curve between 75 and 30 degrees grazing ; upper figure (B) represents mean backscatter between 50 and 40 deg grazing. Parameterization of the backscatter curve may provide important insight into the distribution of seafloor properties; visualization in this mode shows the direct relationship between these parameters and the seafloor morphology