

# **Response of Particulate Optical Properties to Coastal Mixing Processes**

Heidi M. Sosik and Robert J. Olson  
Biology Department, MS 32  
Woods Hole Oceanographic Institution  
Woods Hole, MA 02534-1049

email: [hsosik@whoi.edu](mailto:hsosik@whoi.edu)      email: [rolson@whoi.edu](mailto:rolson@whoi.edu)  
phone: (508) 289-2311      phone: (508) 289-2565  
fax: (508) 457-2134      fax: (508) 457-2134  
Award #: N000149510333

## **LONG TERM GOALS**

Our long-term goals are to develop a better understanding of the relationships between upper ocean optical properties and particulate and dissolved seawater constituents, and to determine how these relationships are influenced by physical processes. Specific long term objectives include both predicting and modeling optical variability relevant for biological processes, such as phytoplankton photosynthesis, and retrieval of information about the biomass and activity of plankton from optical measurements.

## **OBJECTIVES**

Spatial and temporal variability in particulate and dissolved material is a significant source of optical variability in the upper ocean. The primary objective of the present work is to examine the interaction between physical processes and the properties, abundance, and optical significance of different particle types in coastal ocean waters. Specific project objectives are to refine individual particle measurement methods and develop approaches to using individual particle results for interpretation of both inherent and apparent bulk optical properties (IOP/AOP). The project comprises a combination of instrument development and field studies in coastal waters of the eastern US continental shelf.

## **APPROACH**

The approach we have taken employs techniques for characterizing and assessing the optical properties of particles, using both *in situ* and ship-board instrumentation and both bulk and single particle methods. Our primary tools are flow cytometry for assessing individual particle light scattering and fluorescence properties, spectrophotometry for measuring bulk dissolved and particulate absorption spectra (including separation of phytoplankton pigment absorption from the bulk absorption via methanol extraction), and spectral underwater radiometry. Our goal is to conduct flow cytometric and spectrophotometric measurements both on discrete water samples and with *in situ* instruments. *In situ* measurement provides the opportunity for relatively unperturbed sampling, with generally greater spatial resolution, while analysis of discrete water samples continues to allow more detailed characterization of optically-active seawater constituents. We have employed our sampling methods during the Coastal Mixing and Optics (CM&O) field study in continental shelf waters south of Cape Cod, MA (40° 30' N, 70° 30' W).

# Report Documentation Page

Form Approved  
OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

|  |                                   |                                    |                            |   |                                 |
|--|-----------------------------------|------------------------------------|----------------------------|---|---------------------------------|
| 1. REPORT DATE<br><b>30 SEP 1998</b>   |                                   | 2. REPORT TYPE                     |                            | 3. DATES COVERED<br><b>00-00-1998 to 00-00-1998</b> |                                 |
| 4. TITLE AND SUBTITLE<br><b>Response of Particulate Optical Properties to Coastal Mixing Processes</b>   |                                   |                                    |                            | 5a. CONTRACT NUMBER                                 |                                 |
|  |                                   |                                    |                            | 5b. GRANT NUMBER                                    |                                 |
|  |                                   |                                    |                            | 5c. PROGRAM ELEMENT NUMBER                          |                                 |
| 6. AUTHOR(S)   |                                   |                                    |                            | 5d. PROJECT NUMBER                                  |                                 |
|  |                                   |                                    |                            | 5e. TASK NUMBER                                     |                                 |
|  |                                   |                                    |                            | 5f. WORK UNIT NUMBER                                |                                 |
| 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)<br><b>Woods Hole Oceanographic Institution, Biology Department, MS<br/>32, Woods Hole, MA, 02534-1049</b> |                                   |                                    |                            | 8. PERFORMING ORGANIZATION REPORT NUMBER            |                                 |
| 9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)  |                                   |                                    |                            | 10. SPONSOR/MONITOR'S ACRONYM(S)                    |                                 |
|  |                                   |                                    |                            | 11. SPONSOR/MONITOR'S REPORT NUMBER(S)              |                                 |
| 12. DISTRIBUTION/AVAILABILITY STATEMENT<br><b>Approved for public release; distribution unlimited</b>  |                                   |                                    |                            |   |                                 |
| 13. SUPPLEMENTARY NOTES  |                                   |                                    |                            |   |                                 |
| 14. ABSTRACT   |                                   |                                    |                            |   |                                 |
| 15. SUBJECT TERMS  |                                   |                                    |                            |   |                                 |
| 16. SECURITY CLASSIFICATION OF:  |                                   |                                    | 17. LIMITATION OF ABSTRACT | 18. NUMBER OF PAGES                                 | 19a. NAME OF RESPONSIBLE PERSON |
| a REPORT<br><b>unclassified</b>  | b ABSTRACT<br><b>unclassified</b> | c THIS PAGE<br><b>unclassified</b> |                            |   |                                 |

## WORK COMPLETED

With the completion of the Coastal Mixing and Optics field experiment, effort during the last year has been focused on processing and interpretation of the observations collected in summer 1996 (R/V Seward Johnson cruise 9610) and spring 1997 (R/V Knorr cruise 150). Our primary emphasis has been on characterization of optically active particles, assessment of absorption and scattering properties of particulate and dissolved material (including size dependence for particles), and examination of apparent optical properties. We have completed description of the general differences between the summer and spring periods (Sosik et al. 1998) from both a hydrographic and optical perspective. These results have been presented in a variety of forums and we are proceeding with manuscript preparation for peer-reviewed publication.

In addition, we have been continuing to work on particle characterization, based primarily on flow cytometric measurements. During the past year, we have refined our estimates of particle size distributions and reprocessed our entire data set from CM&O. This complete reprocessing included estimation of non-phytoplankton particle concentration and size. Our efforts have also encompassed investigation of theoretical and empirical considerations for interpreting individual particle light scattering and chlorophyll fluorescence.

## RESULTS

Analysis of inherent and apparent optical properties observed during the CM&O experiment has revealed that particles played a dominant role in determining water column optical properties. This is shown clearly by comparison of variations in diffuse attenuation ( $K_d$ ) with absorption coefficients (Green et al. 1998, Figs. 1 and 2). Absorption by dissolved material ( $a_s$ ) exceeded that of particles ( $a_p$ ) at ultraviolet wavelengths; at visible wavelengths, however,  $a_s$  was usually lower and much less variable with depth and time than  $a_p$ . Interestingly,  $a_s$  was somewhat more variable under the stratified conditions present in late summer (Fig. 2) when values were consistently lower in the surface mixed layer compared to the rest of the water column (Sosik et al. 1998).

Based on flow cytometric analysis, it is possible to examine the optical role of different types of particles in greater detail (DuRand et al 1998). We have extended conventional flow cytometric measurements of marine particles, which has focused exclusively on phytoplankton, to include quantitative examination of other particles. This broad class of “non-phytoplankton” material presumably includes particles of organic detritus and of mineral origin, as well as heterotrophic organisms. We have enumerated and estimated optical properties of three types of particles: 1) picophytoplankton of the genus *Synechococcus*, 2) eukaryotic phytoplankton (~2-30  $\mu\text{m}$ ) and 3) other particles in the ~1-30  $\mu\text{m}$  size range.

Results of flow cytometric analysis show that non-phytoplankton particles dominated numerically in both late summer and spring and were less variable in space and time than the phytoplankton. In late summer before hurricane Eduoard, *Synechococcus* abundances were very high with a strong subsurface maximum present ( $> 10^5$  cells  $\text{ml}^{-1}$ ), while in spring these cells were 10-fold less abundant (DuRand et al. 1998, Sosik et al. 1998). Eukaryotic phytoplankton did not differ in abundance between the two seasons, until the bloom at the end of the spring sampling period when concentrations increased 3-4 fold (Fig. 3).

Particles were found to have substantial differences in their particle specific optical properties that affected the overall contribution to bulk water column optical properties. This is evident when relative contributions to light scattering are considered; the significance of both *Synechococcus* and the non-phytoplankton particles decreases relative to their numerical abundance (compare Figs. 3 and 4, Fig. 5). In the spring, in particular, the importance of eukaryotic phytoplankton often exceeded the other particles in surface waters, but decreased rapidly with depth.

Current work on these topics is focused on deriving absolute (rather than relative) estimates of particle scattering cross-sections and estimating backscattering contributions. Preliminary results investigating differences in spatial and temporal variability between diffuse attenuation and spectral reflectance suggest that changes in the particle size distribution and associated scattering effects are important in regulating apparent optical properties of the water column.

## **IMPACT/APPLICATIONS**


This project includes the development of improved techniques for analyzing marine particles and characterizing their optical properties. Our ability to independently quantify size distributions for phytoplankton and non-phytoplankton particles is a new contribution that will lead to better understanding of optical variability in the ocean.

## **TRANSITIONS**


We have several on-going collaborations with other investigators participating in the CM&O program. Our results from sampling on mooring-turnaround cruises are being used by Dr. Tommy Dickey's group at UCSB for interpretation of observations from moored sensors. Radiometry results from the main optics cruises have been provided to Dr. Ron Zaneveld at OSU for investigating relationships among diffuse attenuation for irradiance, remote sensing reflectance and absorption and scattering coefficients. Some results have been transferred to A. Robinson's group at Harvard for use in coupled physical – optical modeling and we have shared pigment, nutrient and size distribution results with Dr. Wilf Gardner's group at TAMU.

## **RELATED PROJECTS**


This project is closely tied with a NASA New Investigator Program award (Sosik) for investigating the regulation of local biological production of particles at the CM&O site and to explore the effects of changes in particle properties on ocean color. In addition, Olson is independently funded (DOE, NSF) to develop the in situ flow cytometer. The interpretation of flow cytometric light scattering and fluorescence distributions is also being supported by a NSF JGOFS project for work in the Arabian Sea (Olson). Due to the interdisciplinary nature of the Coastal Mixing and Optics research initiative, this project is also closely tied to several others funded by ONR; specific collaborations and exchanges are underway with C. Roesler (pigment and particle absorption), S. Pegau and R. Zaneveld (IOP/AOP relationships), T. Dickey and G. Chang (particle absorption and modeling of primary production), and W. Gardner and J. Blakely (water properties and particle size distributions).




**Figure 1. Relationships between diffuse attenuation for downwelling irradiance ( $K_d$ ) and absorption coefficients for dissolved ( $a_s$ ) and particulate ( $a_p$ ) material at three wavelengths during August-September 1996.  $K_d$  estimates were derived from irradiance measurements collected with a tethered free-fall profiler (Satlantic SPMR) and the absorption data was collected using ac-9 sensors (WetLabs) mounted on the OSU Slowdrop profiling package. Data were averaged in 10-m bins and  $K_d$  and absorption profiles collected within two-hour windows were used. Linear regression results show the importance of dissolved material at 412 nm and the strong dependence of  $K_d$  variations on  $a_p$  at 443 and 488 nm.**



**Figure 2. Relationships between diffuse attenuation for downwelling irradiance ( $K_d$ ) and absorption coefficients for dissolved ( $a_s$ ) and particulate ( $a_p$ ) material at three wavelengths during April-May 1997. Data were collected and processed as described for Fig. 1. Linear regression results show strong dependence of  $K_d$  variations on  $a_p$ , with even less contribution of dissolved material than observed under conditions of greater stratification in late summer the previous year (Fig. 1).**




*Figure 3. Time series of particle concentration observed during vertical sampling in spring 1997. Results for three general particle classes are shown, *Synechococcus* (top panel), eukaryotic phytoplankton (middle panel), and other particles in the same size range (bottom panel).*



*Figure 4. Time series of relative contributions of different particles to forward light scattering observed during vertical sampling in spring 1997. Results for three general particle classes are shown, Synechococcus (top panel), eukaryotic phytoplankton (middle panel), and other particles in the same size range (bottom panel).*





**Figure 5.** Summary of abundance and integrated forward light scattering for different types of particles sampled by flow cytometry in spring 1997. Mean +/- standard deviation is shown for three time periods and three particle types: *Synechococcus* (red), eukaryotic phytoplankton (green) and other particles in the same size range (black).

## REFERENCES

- DuRand, M. D., H. M. Sosik and R. J. Olson. 1998. Size-dependent light scattering and absorption by particles in continental shelf waters: flow cytometric analysis during the Coastal Mixing and Optics Experiment. Abstract, 1998 AGU/ASLO Ocean Sciences Meeting.
- Green, R. E. and H. M. Sosik. 1998. Vertical and temporal variability in apparent optical properties during the Coastal Mixing and Optics Experiment. Abstract, 1998 AGU/ASLO Ocean Sciences Meeting.
- Sosik, H. M., R. E. Green and R. J. Olson. 1998. Optical variability in coastal waters of the Northwest Atlantic. In: Ocean Optics XIV, S. G. Ackleson (ed.). In press.

Web sites:

<http://www.whoi.edu/science/B/sosiklab/cmo.htm>