

Development And Validation Of Depth Inversion Algorithms For Barred-Beaches Based On Nonlinear Properties Of Shoaling Waves

Stephan T. Grilli

Department of Ocean Engineering

University of Rhode Island

Narragansett, RI 02882

phone: (401) 874-6636 fax : (401) 874-6837 email: grilli@oce.uri.edu

Award Number: N00014-99-1-0439

<http://www.oce.uri.edu/~grilli>

LONG-TERM

The long-term goal of this work is to develop methods for predicting the bottom topography in nearshore regions, based on characteristics of (shoaling) surface waves, measured using remote sensing methods. We are particularly interested in the surfzone region, over barred-beaches, where large wave nonlinearity and breaking occur.

OBJECTIVES

Earlier work dealt with monotonous topographies and periodic non-breaking waves (Grilli, 1997, 1998). In this work, the objectives are to :

- (1) calculate and analyze fully-nonlinear periodic wave shoaling over barred-beaches in a two-dimensional Potential Flow Model (PFM).
- (2) implement and calibrate a spilling breaker model in the PFM shoaling region, and accurately model breaking waves in the surfzone.
- (3) compute shoaling of both non-breaking and breaking mildly irregular waves (with narrow-banded spectra), over barred-beaches.
- (4) modify/extend earlier depth-inversion-algorithms (DIA) to predict depth in the region over and beyond the bar, based on (measurable) global wave characteristics.
- (5) improve the model computational efficiency for large-size/long-term computations.
- (6) validate both shoaling calculations and depth predictions using field data (possibly from Duck (SHOWEX); John Dugan's measurements).

APPROACH

In earlier work, DIAs were developed for cylindrical beaches (Grilli, 1997, 1998). These were based on inverting a nonlinear wave celerity relationship, obtained from results of direct shoaling computations in a fully-nonlinear PFM. Since the nonlinear celerity also depends on wave steepness,

Report Documentation Page

Form Approved
OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE 30 SEP 1999		2. REPORT TYPE		3. DATES COVERED 00-00-1999 to 00-00-1999	
4. TITLE AND SUBTITLE Development And Validation Of Depth Inversion Algorithms For Barred-Beaches Based On Nonlinear Properties Of Shoaling Waves				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Rhode Island, Department of Ocean Engineering, Narragansett, RI, 02882				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a REPORT unclassified	b ABSTRACT unclassified	c THIS PAGE unclassified			

wave height or front-to-back asymmetry data was also required, in addition to wave phases, to perform depth inversion.

A similar approach is pursued in this effort. The PFM will first be extended to deal with barred-beaches and non-periodic, possibly breaking, incident waves. Direct problems will first be calculated and a parametric study of wave characteristics in the region beyond the bar will be performed. Modifications/extensions of DIAs will be made, based on these results. Shoaling calculations and depth predictions will be validated using field data.




Figure 1 : PFM computations for the shoaling of periodic waves over a barred-beach with equilibrium shape (1:50 average slope). Definition of global wave parameters (H , L , c). AB : absorbing beach. Note the wave decomposition region beyond the bar.

WORK COMPLETED

Direct computations and analyses of periodic waves, shoaling over barred-beaches, were performed in a two-dimensional PFM (Grilli and Horrillo, 1999; e.g., Fig. 1). DIAs developed earlier for monotonous topographies were applied, and discrepancies in depth prediction were observed in the region beyond the bar; analyses showed, these are due to wave decomposition phenomena occurring in this region (Grilli and Skourup, 1999).

Groundwork was done for the implementation of a spilling breaker model in the shoaling region of the potential model. This model is based on specifying an absorbing surface pressure (Grilli and Horrillo, 1997), whose work is calibrated with the energy dissipation obtained from a semi-empirical spilling breaker model.

A Volume Of Fluid (VOF) model, solving Navier-Stokes equations, was applied to the modeling of breaking surfzone waves. The VOF model was coupled to the PFM used in the shoaling zone (Guignard et al., 1999; e.g., Fig. 2). Preliminary results show good agreement with laboratory data (Guignard et al., 2000).




Figure 2 : *Coupled PFM/VOF computations for the shoaling and breaking of a solitary wave over a plane 1:15 beach. Curves a-g denote successive times.*

In a parallel, independently supported, work, a three-dimensional fully nonlinear PFM, applicable to arbitrary bottom topography, was developed (Grilli et al., 1999; e.g., Fig. 3). This model could be used to estimate effects induced on shoaling waves by three-dimensional topographic features, as compared to two-dimensional results.




Figure 3 : *Three-dimensional PFM computations for the shoaling and breaking of a solitary wave over a plane 1:15 beach with a middle ridge.*

Contacts were made with John Dugan (Arete Assoc.) who will conduct field experiments at Duck (SHOWEX), using airplane based sensors. A comparison of depth prediction using linear wave theory based and nonlinear DIAs will be made using his data.

RESULTS

Preliminary PFM computations for a barred-beach identify the need for refining DIAs in the wave modulation region beyond the bar. Computed wave front and back slope variations in this region show that wave slopes, which are obtainable from remotely sensed surface observations, may help identifying bar topographic features.

The coupled two-dimensional PFM/VOF model provides a fully nonlinear solution for wave shoaling from deep water, up to breaking in the surfzone. Breakers are well modeled by the VOF solution of Navier-Stokes equations, which also includes modeling of the air flow above the free surface, and of air pockets trapped in the water. The coupled model will make it possible to accurately calculate nonlinear wave parameter variations (celerity, height, wavelength, asymmetry), from deep water up to runup on the shore.

IMPACT/APPLICATIONS

The two- and three-dimensional models refined and developed as part of this work constitute robust and flexible nonlinear wave modeling tools, applicable to a variety of ocean and coastal engineering problems. The models were developed as so-called “numerical wave tanks”, in which many types of boundary conditions and geometry can be easily specified. In particular, it would be easy to include fixed or floating structures in the models, and to calculate wave-structure interactions. The models were based on a higher-order Boundary Element Method, which is both efficient and accurate. For more efficiency, the two-dimensional coupled model works on a bi-processor computer (Guignard et al., 2000), and the three-dimensional model has been optimized for use on a CRAY-90 multi-vector processors (Grilli et al., 2000).

TRANSITIONS

The two- and three-dimensional modeling work performed in this project is closely related to other (collaborative) wave modeling efforts the PI is involved with : (i) the generation and coastal runup of tsunamis, due to underwater landslides (Grilli and Watts, 1999); (ii) the modeling of three-dimensional breaking waves over various coastal topographies, and the study of their kinematics (Grilli et al., 2000); (iii) the three-dimensional nonlinear wave focusing, as a possible mechanism for creating very large (freak) waves.

REFERENCES

Grilli, S.T. and J., Horrillo 1997 Numerical generation and absorption of fully nonlinear periodic waves. *J. Engng. Mech.*, 123 (10), 1060-1069.

Grilli, S.T. 1998 The depth inversion problem in shallow water. In *Proc. 3rd Intl. Intl. Sympos. on Ocean Wave Measurement and Analysis* (Virginia Beach, VA, November 1997), 701-715, ASCE Publication.

Grilli , S.T. 1998 Depth inversion in shallow water based on nonlinear properties of shoaling periodic waves. *Coastal Engng.*, 35(3), 185-209.

PUBLICATIONS

Grilli, S.T. and J., Skourup 1999 Depth inversion for nonlinear waves shoaling over a barred-beach. In *Proc. 26th Intl. Conf. on Coastal Engineering* (Copenhagen, Denmark, June 1998), 603-616.

Guignard, S., Grilli, S.T., Marcer, R. and V., Rey 1999 Computation of shoaling and breaking waves in nearshore areas by the coupling of BEM and VOF methods. In *Proc. 9th Offshore and Polar Engng. Conf.* (Brest, France, May 1999), Vol. III, 304-309.

Grilli , S.T. and J., Horrillo 1999 Shoaling of periodic waves over barred-beaches in a fully nonlinear numerical wave tank. *Intl. J. Offshore and Polar Engng.* (in press)

Grilli, S.T. and Watts, P. 1999 Modeling of waves generated by a moving submerged body. Applications to underwater landslides. *Engng. Analysis Boundary Elemts.* (in press).

Grilli, S.T., Guyenne, P. and F., Dias 1999 A fully nonlinear model for three-dimensional overturning waves over arbitrary bottom. Intl. J. Numer. Methods in Fluids (submitted)

Guignard, S., Grilli, S.T. and V. Rey 2000 Computation of breaking and post-breaking wave characteristics on slopes by coupling of BEM and VOF models. Abstract accepted for 27th Intl. Conf. on Coastal Engineering (ICCE27, Sidney, Australia, July 2000).

Grilli, S.T., Guyenne, P. and F., Dias 2000 Numerical modeling of fully nonlinear 3D overturning waves over arbitrary bottom. Abstract accepted for 27th Intl. Conf. on Coastal Engineering (ICCE27, Sidney, Australia, July 2000).