

Quantitative Chemical Mass Transfer in Coastal Sediments During Early Diagenesis: Effects of Biological Transport, Mineralogy, and Fabric --- Phase III

Dawn Lavoie
Code 7431

Naval Research Laboratory
Stennis Space Center MS 39529

phone: (228) 688-4659 fax: (228) 688-5752 email: dawn.lavoie@nrlssc.navy.mil

Award #: N0001499WX30240

LONG-TERM GOAL

The long-term goal of this study is to develop a better mechanistic and quantitative understanding of the effects of biologically enhanced transport, mineralogy, sediment fabric, and particle surface chemistry on biogeochemical reactions occurring in coastal sediments. Specifically, we plan to integrate quantitative expressions of the strongly coupled effects of bioirrigation, bioturbation, mineralogy, and sediment fabric on chemical mass transfer from field and laboratory mesocosm studies using numerical modeling.

OBJECTIVES

The short term objective during FY99 was (1) to achieve a better understanding of the bio-geologic dynamics of estuarine sediments by quantitatively describing the relationship between bioturbation, sediment physical properties, and permeability and (2) to accurately compute the permeability coefficient from two-dimensional images using Effective Medium Theory (EMT), and verify results with in situ, mesocosm and laboratory permeability measurements. Our hypothesis is that bioturbation will alter sediment microfabric and thus sediment permeability.

APPROACH

The approach is an integration of field sampling, controlled experiments in laboratory mesocosms, image analysis and numerical modeling. Experimental results are measured by laboratory analyses of bulk physical properties; macro/microscopic analyses of sediment fabric utilizing x-radiography; CT scanning and transmission electron microscopy; and permeability modeling using code based on Effective Medium Theory. This work is closely coordinated with Y. Furukawa, funded under Award # NO0014-98-1-0200, who is analyzing pore fluids, bulk geochemistry and vertical distribution of ^{13}C ; and P. VanCappellan (Georgia Institute of Technology) who is incorporating our results into the RT model, STEADYSED.

FY99 tasks included:

- Characterization of burrow networks, microfabric, and permeability for sediments populated with different types of infauna in laboratory mesocosms at NRL.
- Bimonthly field study of physical properties, microfabric, and burrow networks of burrowed nearshore sediments at St. Louis Bay and Mississippi Sound, Mississippi.

Report Documentation Page

Form Approved
OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE 30 SEP 1999		2. REPORT TYPE		3. DATES COVERED 00-00-1999 to 00-00-1999	
4. TITLE AND SUBTITLE Quantitative Chemical Mass Transfer in Coastal Sediments During Early Diagenesis: Effects of Biological Transport, Mineralogy, and Fabric --- Phase III				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Research Laboratory, Code 7431, Stennis Space Center, MS, 39529				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

- Field sampling and analyses at Sapelo field site in conjunction with GIT.
- Further development of techniques to image and quantify burrow and pore networks at various spatial and temporal scales.

WORK COMPLETED

- St. Louis Bay field studies are completed. Comparison with Sapelo Island field data is continuing.
- Initial mesocosm experiments with *Schizocardium* sp. (acorn worms) are completed. Experiments with *Hemipholis elongata* and *Ophiophragmus moorei* (brittle stars) has been initiated.
- Microfabric comparison between burrow walls, ejecta mounds of *Schizocardium* sp. and matrix sediment is completed.
- Model sensitivity study is completed.

RESULTS

- ***St. Louis Bay field studies.***

Physical properties profiles from the open Sound site show the expected trends produced by sediment consolidation. Very high initial water-related values are attributed to bioturbation in the upper 12-15 cm. Physical properties trends from the grassy marsh sites lack the trends expected from consolidation processes and reflect the influence of other processes such as seasonal variation in water level and salinity, and the binding effects of marsh grasses. Trends of matrix sediment in the mesocosms (between burrows) reflect consolidation processes; water-related properties around the burrows (e.g., in ejecta mounds) are extremely high.

Permeability measured in the aquaria with a miniaturized in situ probe reflects the influence of burrows. Tanks populated with 96 worms/m² have lower permeability (two orders of magnitude) than tanks with 800 worms/m² (figure 1). The main conclusion resulting from the St. Louis Bay and mesocosm studies is that fluid flow in heavily burrowed sediment is controlled by the burrows, and that permeability through the matrix sediment is insignificant when compared with the permeability of the entire sediment volume.

- ***Modeling results.***

A sensitivity study of the EMT model to input parameters indicates that pore throat radius is the main variable controlling permeability, that is, the size of the smallest constricting passages has more effect than other variables. As the size of pore bodies increases, permeability decreases, probably because as the pore body size increases, the pore body must be filled before fluid flow is resumed. The coordination number (number of pore throats connecting pore bodies) is critical when the volume has few pore bodies. As the number of bodies increases, the influence of coordination number decreases. Use of the harmonic mean rather than the arithmetic mean produces permeability values closer to measured permeability values because the importance of the smaller pore spaces is emphasized.

- ***Microfabric analyses***

Image porosity, analyzed using Image Tool shows the influence of bioturbation on water-related properties and permeability. Large differences in void space exist between burrow walls, ejecta mounds and matrix sediment (figure 2). In situ permeability using a miniaturized

in situ probe shows parallel trends in permeability and demonstrates permeability variability on the centimeter scale.

IMPACT/APPLICATION

A better understanding and mathematical description of biologically-enhanced transport, sediment fabric and particle surface chemistry during shallow diagenesis will allow us to better model and predict the fate and transport of solutes, particles and associated pollutants. By concentrating on fine-grained sediments over the next few years, we hope to make a significant contribution understanding harbor pollution solutions. In addition, by understanding the effect of fabric changes during diagenesis, we will be able to better predict sediment physical and geacoustic properties of interest to the MCM community, (predicting mine burial in shallow coastal regions) and the acoustic community for modeling acoustic propagation.

TRANSITIONS

Techniques for quantitative characterization (2D and 3D) of sediment macro and microfabric will be transitioned to other ONR-funded programs including the High-frequency Sound Interaction in Ocean Sediments: Modeling Environmental Controls DRI. It is anticipated that results from this effort will contribute to applied environmental programs in the future.

RELATED PROJECTS

This project has leveraged the NRL 6.1 core program (Microenvironmental Studies) for support, particularly for the field effort, and will continue to do so. Microfabric results have been used in modeling efforts to predict permeability and will undoubtedly continue to benefit other programs with similar requirements, e.g., the ONR High Frequency Sound Interaction DRI will require quantitative pore space and particle geometry data for prediction of permeability and porosity.

REFERENCES/PUBLICATIONS

Publications (in press/submitted)

- Lavoie, D., Watkins, J., and Furukawa, Y. (1999) Microwave processing of sediment samples. (In: *Microwave Protocols for Microscopy*. Eds., R. T. Giberson and R. Demaree) Humana Press (due out November, 1999).
- Furukawa, Y., Bentley, S., Shiller, A., Lavoie, D., and Van Cappellen, P. The role of biologically-enhanced pore water transport in the early diagenesis of carbonate sediments from North Key Harbor, Dry Tortugas National Park, Florida. *J. Mar. Res.*
- Vaughan, C. (1999) A quantitative prediction of permeability in fine-grained estuarine sediments as a function of bioturbation. University of New Orleans. (M.S. degree expected Dec. 1999).

Published

- Furukawa, Y., Lavoie, D. and Stephens, K. (1997) Effect of biogeochemical diagenesis on sediment fabric in shallow marine carbonate sediments near the Dry Tortugas, Florida. *Geo-Marine Letters*, 17, 283-290.
- Furukawa, Y., and Lavoie, D. (1997) A geochemical investigation of

early diagenetic effects on sedimentary structure; Proceedings of the Coastal Benthic Boundary Layer Key West Workshop. NRL MR-8044; NRL/MR/7431-97-8044, 7p.

- Lavoie, D., Stephens, K., Furukawa, Y., and Lavoie, D. (1997) Geotechnical characteristics of Dry Tortugas and Marquesas Sediments; Proceedings of the Coastal Benthic Boundary Layer, Key West Workshop. NRL MR-8044; NRL/MR/7431-97-804, 9p.
- Stephens, K. P., Lavoie, D. L., Briggs, K. B., Furukawa, Y., and Richardson, M. D. (1997) Geotechnical and geoacoustic properties of sediments off South Florida: Boca Raton, Indian Rocks Beach, Lower Tampa Bay, and the Lower Florida Keys. NRL/MR/7431-97-8042, 303p.

Abstracts

- Furukawa, Y. (1999) A TEM investigation of clay-organic matter sediments. 217th American Chemical Society National Meeting and Exposition.
- Furukawa, Y. and Lavoie, D. (1999) Relationship between measured in situ permeability and diffusive transport of pore water constituents. 1999 Annual Meeting of the Mississippi Academy of Sciences. Tupelo, MS.
- Vaughan, C., D. Lavoie, A. Reed and Y. Furukawa (1999) A quantitative prediction of permeability as a result of biologically induced diagenesis. 1999 Annual Meeting of the Mississippi Academy of Sciences. Tupelo, MS.
- Furukawa, Y., Lavoie, D. and Stephens, K. (1997) Early diagenesis of biologically reworked carbonate sediments near the Dry Tortugas, Florida. *Geological Society of America Abstracts with Program*, 29.
- Furukawa, Y., Lavoie, D., Vaughan, C. and Reed, A. (1998) Reactive transport of dissolved species during early diagenesis in shallow water muddy sediments of St. Louis bay, Mississippi. American Geophysical Union 1998 Fall Meeting.




Figure 1. Comparison of permeability, measured in situ using the in situ permeameter, in the laboratory using a falling head method and modeled using EMT code, as a function of burrowing in St. Louis Bay, Mississippi Sound and laboratory aquaria.




Figure 2. Image analysis of 2D microfabric shows significant differences in porosity between sediment in the undisturbed matrix, ejecta mounds and burrow walls.