

UNCLASSIFIED

AD NUMBER

ADB173090

LIMITATION CHANGES

TO:

Approved for public release; distribution is unlimited.

FROM:

Distribution authorized to U.S. Gov't. agencies only; Proprietary Information; MAY 1993. Other requests shall be referred to Army Medical Research and Development Command, Attn: SGRD-RMI-S, Fort Detrick, Frederick, MD 21702-5012.

AUTHORITY

USAMRDC, ltr, 30 Jun 1994

THIS PAGE IS UNCLASSIFIED

(UNCLASSIFIED)



AD NUMBER

AD-B 173 090

LIMITATION CHANGES

TO Approved for Public Release
Distribution Unlimited

FROM

N/A

AUTHORITY

Carey O. Leeverett / Phyllis Rinehart,
Deputy Chief of Staff for Information
Management, Ft Detrick / S&RD-RM1-S
(70-y), Via Memo Dtd 30 Jun 94

(THIS PAGE IS UNCLASSIFIED)

AD-B173 090



AD _____

(2)

ELECTRON MICROSCOPY TO CORRELATE CELL STRUCTURE
AND BIOCHEMICAL ACTIVITY

FINAL REPORT

MASAMICHI AIKAWA

MARCH 10, 1993

Supported by

U.S. ARMY MEDICAL RESEARCH AND DEVELOPMENT COMMAND
Fort Detrick, Frederick, Maryland 21702-5012

Contract No. DAMD17-90-C-0010

Institute of Pathology
Case Western Reserve University
2085 Adelbert Road
Cleveland, Ohio 44106

Distribution authorized to U.S. Government agencies only,
proprietary information, May 5, 1993. Other requests for this
document shall be referred to Commander, U.S. Army Medical
Research and Development Command, ATTN: SGRD-RMI-S, Fort
Detrick, Frederick, Maryland 21702-5012.

The findings in this report are not to be construed as an
official Department of the Army position unless so designated
by other authorized documents

93 5 14 02 4

93-10828

DTIC
ELECTE
MAY 17 1993
S C D

REPORT DOCUMENTATION PAGE

Form Approved
OMB No 0704-0188

Public report documentation (including abstracts) should be placed in the public domain including that material containing instructions, illustrations, tables, charts, etc. The Government is authorized to reproduce and distribute reprints for Government purposes not withstanding any copyright notation that may appear hereon. This document is in the public domain in the United States of America. For all other countries, permission should be sought from the Copyright Clearance Center, Inc., 27 Congress St., Salem, MA 01970. Organizations in the U.S. who are also registered with the Copyright Clearance Center may also clear this document for intercountry distribution. For all other countries, permission should be sought from the Copyright Clearance Center, Inc., 27 Congress St., Salem, MA 01970. Organizations in the U.S. who are also registered with the Copyright Clearance Center may also clear this document for intercountry distribution. For all other countries, permission should be sought from the Copyright Clearance Center, Inc., 27 Congress St., Salem, MA 01970.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE March 10, 1993	3. REPORT TYPE AND DATES COVERED Final 15 Nov 89 - 14 Nov 92	
4. TITLE AND SUBTITLE Electron Microscopy to Correlate Cell Structures and Biochemical Activity			5. FUNDING NUMBERS DAMD17-90-C-0010 61102A 30161102BS13 DA346203	
6. AUTHOR(S) Masamichi Aikawa				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Institute of Pathology Case Western Reserve University 2085 Adelbert Road Cleveland, Ohio 44106			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING, MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research & Development Command Fort Detrick Frederick, Maryland 21702-5012			10. SPONSORING MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION AVAILABILITY STATEMENT Distribution authorized to U.S. Government agencies only, proprietary information, May 5, 1993.			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) We performed immuno-electron microscopy to study intracellular localization and fate of liposome-encapsulated malarial antigen after phagocytosis by macrophages. Liposome-encapsulate protein that is phagocytosed by macrophages can enter an intracellular compartment in which at least some of the antigenic epitopes are not degraded by lysosomal enzymes. We also localized <u>P. falciparum</u> antigen having molecular weight 175 kDa (EBA) within the parasite by immunoelectron microscopy. EBA is specifically localized in micronemes of <u>P. falciparum</u> merozoites. The CS protein, SSP2, the Duffy receptor and now EBA-175 have been identified in the micronemes, suggesting that these organelles play a critical role in the storage and release of parasite proteins that act as receptors for invasion into host cells. <u>P. coatneyi</u> produces knobs on the membrane of PRBC and these PRBC appear to sequester in the vasculature of infected rhesus monkeys. We studied the pathology of CNS of rhesus monkeys infected with <u>P. coatneyi</u> and found that cytoadherence of PRBC to endothelial cells is a consistent feature of infections with this primate parasite. Cerebral microvessels with sequestered PRBC were shown by immunohistochemistry to possess CD36, TSP and ICAM-1. Our study indicates, for the first time, that rhesus monkeys infected with <u>P. coatneyi</u> can be used as a primate model to study human cerebral malaria.				
14. SUBJECT TERMS Malaria; immunoelectron-microscopy; Chemotherapy; DFO; Cerebral malaria; RA 1			15. NUMBER OF PAGES	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

DISCLAIMER NOTICE



THIS DOCUMENT IS BEST QUALITY AVAILABLE. THE COPY FURNISHED TO DTIC CONTAINED A SIGNIFICANT NUMBER OF COLOR PAGES WHICH DO NOT REPRODUCE LEGIBLY ON BLACK AND WHITE MICROFICHE.

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

Where copyrighted material is quoted, permission has been obtained to use such material.

Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

Citations of commercial organizations and trade names in this report do not constitute an official Department of the Army endorsement or approval of the products or services of these organizations.

In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Animal Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).

For the protection of human subjects, the investigator(s) have adhered to policies of applicable Federal Law 45 CFR 46.

In conducting research utilizing recombinant DNA technology, the Investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

<i>Masanishi Akira</i>	<i>3/10/93</i>
PI Signature	Date

TABLE OF CONTENTS

Introduction 5

Body 6

Conclusions 14

References 15

Appendix 18

Figures 19

Publications 23

DTIC QUALITY ASSURANCE 3

Accession For	
NTIS CRA&I	<input type="checkbox"/>
DTIC TAB	<input checked="" type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
B-3	

INTRODUCTION:

The main objective of our work was to function as a basic core electron microscopy facility in collaboration with scientists at WRAIR. In particular we proposed to carry out studies on morphological effects of drugs on various microorganisms by electron microscopy and autoradiography in order to understand the mode of actions of these drugs.

In the past, electron microscopy has been used to investigate the mode of action of chemotherapeutic agents on a variety of microorganisms. Studies of chloroquine, primaquine, pyrimethamine, qinghaosu, pentamidine and WR-163577 have indicated that each compound produces specific changes in particular organelles (1-7). The results obtained by electron microscopy have correlated well with biochemical observations on the action of drugs. Therefore, studies of the morphological effects of drugs on various microorganisms can contribute greatly to our understanding of their mode of action.

Secondly, immuno-electron microscopy has been used to characterize protective antigens of malarial parasites. Techniques of immunology and molecular biology have been used to identify and isolate specific proteins that may be capable of producing protective immunity against malarial parasites. Immuno-electron microscopy has played an important role in identifying specific immune targets and characterizing the precise location and expression of these targets (8,9). Such work will help to develop successful vaccines against microorganisms.

BODY:

During this period, we have performed electron microscopy (standard and immuno-electron microscopy) to investigate the mode of action of antimalarials and localization and characterization of protective malarial antigens in host and parasites. Important findings are described as follows:

- a) Phagocytosis of liposomes by macrophages: intracellular fate of liposomal malaria antigen.

Liposomes continue to be useful as models for understanding the disposition patterns of materials ingested by macrophages. The fate of liposome-encapsulated protein in macrophages is of particular interest in immunology. Liposomes are highly effective in the stimulation of immune responses to encapsulated protein antigens (10). It is widely believed that the process of immunological presentation of protein antigen to cells in the immune system involves initial processing of the antigen by specialized antigen presenting cells such as macrophages and B lymphocytes. It is presumed that the ability of liposomes to enhance immune responses is due to natural targeting of liposomes to macrophages. In collaboration with Col. Carl R. Alving and his associates of the Department of Membrane biochemistry, WRAIR, we studied intracellular localization and fate of liposomes and liposome-encapsulated malarial antigen after phagocytosis by macrophages.

In the present study, we performed immunoelectron microscopy to detect the presence of all surface-associated and intracellular liposome-associated malarial antigens. After incubation of macrophages with liposomes containing malarial antigen, vacuoles appeared in the cytoplasm and the vacuoles contained numerous liposomes that were densely labelled by antibody to malarial antigen. Many of the protein-containing liposomes apparently remained intact with a considerable amount of liposome-associated antigen for at least as long as 6 hours after incubation. Liposomes were mainly located at the peripheries of the vacuoles and often seemed to adhere to old vacuolar membranes (Fig.1). Antigenic epitopes detected by the antibody were often observed in the cytoplasm next to the vacuoles. The original antigen consisted largely of repeats of ASN-ALA-ASN-PRO, and it is possible that fragments containing ASN-ALA-ASN-PRO were recognized by the antibody. However, liposomes were invariably present only within the vacuoles and liposome-like structures were rarely, if ever, present in the cytosol of associated with any other subcellular organelles.

We concluded that liposome-encapsulated protein that is phagocytosed by macrophages can enter an intracellular compartment in which at least some of the antigenic epitopes are not degraded by lysosomal enzymes. Furthermore, although liposomes do enter large vacuoles, at least some of the antigenic epitopes, as detected by a monoclonal antibody, can escape into the cytoplasm. These experiments therefore provide evidence to support the

possible existence of a pathway in which liposomal contents can bypass lysosomal degradation. The occurrence of an intracellular pathway for endocytosed or phagocytosed materials that avoids lysosomes has been previously hypothesized and such a pathway therefore might be used as a basis for delivery of non-degraded liposomal antigenic protein epitopes to the cytosol and to the cell surface.

b) A study on a primate model for human cerebral malaria

Human cerebral malaria is a pernicious manifestation of infection with *P. falciparum*. Possible factors contributing to the development of cerebral malaria include the blockage of cerebral microvessels by parasitized erythrocytes (PRBC), deposition of immune complexes in cerebral microvessels, reduced humoral or cell-mediated immune responses and action of tumor necrosis factor (TNF) (11).

Electron microscopy revealed multiple electron dense knobs protruding from the membrane of PRBC seen in the cerebral microvessels of cerebral malaria patients (12-13). These knobs attach via a parasite ligand to receptors on endothelial cells, resulting in the blockage of cerebral microvessels (Fig.2,3). Recently several investigators reported that host cell molecules such as CD36, thrombospondin (TSP) and intercellular adhesion molecule-1 (ICAM-1) may function as the endothelial cell surface receptors for PRBC. However, there is little information as to

whether these molecules actually play a role in cytoadherence of PRBC *in vivo*. We have demonstrated by immunohistochemistry the presence of CD36, TSP and ICAM-1 on the endothelial cells of cerebral microvessels of fresh-frozen, non-malarious human brain tissues (12). However, formalin-fixed and paraffin-embedded brain tissues do not stain well with antibodies against these molecules by immunohistochemical methods. Thus, the use of fresh brain tissue from cerebral malarial patients appears to be essential for studies of the potential role of these molecules in the development of cerebral malaria. However, it is difficult to obtain fresh brain tissue from cerebral malaria patients, since they often die in malaria endemic areas where no freezing facilities are available.

P. coatneyi produces knobs on the membrane of PRBC and these PRBC appear to sequester in the vasculature of infected rhesus monkeys (14). There have been no reports, however, indicating whether cytoadherence of PRBC to endothelial cells of cerebral microvessels occurs *in vivo*. Therefore, in collaboration with Col. Webster, Maj. Brown and Maj. Smith of AFRIMS, Bangkok, Thailand, we studied the pathology of the central nervous system (CNS) of rhesus monkeys infected with *P. coatneyi* in order to determine whether cytoadherence of PRBC to endothelial cells is a consistent feature of infections with this primate parasite and, if so, whether putative receptors for cytoadherence in human cerebral microvessels such as CD36, TSP and ICAM-1 are also present.

Our study demonstrated PRBC sequestration (Fig.4) and cytoadherence of knobs on PRBC to endothelial cells in cerebral microvessels of these monkeys (Fig.5). Cerebral microvessels with sequestered PRBC were shown by immunohistochemistry to possess CD36, TSP and ICAM-1. These proteins were not evident in cerebral microvessels of uninfected control monkeys. Therefore, our study indicates, for the first time, that rhesus monkeys infected with P. coatneyi can be used as a primate model to study human cerebral malaria. By using this animal model, we may be able to make strategies to develop vaccines to prevent human cerebral malaria (15).

In addition, we are planning to study morphological effects of antimalarial drugs on sequestered parasitized erythrocytes in order to understand how sequestered PRBC become detached from endothelial cells of cerebral microvessels by various antimalarials. Since this aspect of human cerebral malaria has never been studied in the past, our study will help to understand the mode of action of various antimalarial drugs on human cerebral malaria.

- c) A primate model for Human Cerebral Malaria: Effects of Artesunate (Qinghaosu derivative) on Plasmodium coatneyi-infected rhesus monkeys

In collaboration with Drs. Kyle Webster, Dennis E. Kyle, Arthur Brown, C. Dahlem Smith, C. Ockenhouse, and Kevin Corcoran of WRAIR and AFRIMS, we studied the effects of artesunate on rhesus monkeys infected with Plasmodium coatneyi (15). Sixteen rhesus monkeys were placed in four groups; group I consisted of three monkeys which were splenectomized and were treated with 3 doses (loading dose: 3.3 mg/kg; maintenance doses: 1.7 mg/kg) of artesunate; group II consisted of three monkeys which were treated with 3 doses of artesunate; group III consisted of two monkeys which were treated with one dose (3.3 mg/kg) of artesunate; group IV consisted of five untreated monkeys. Parasitemia of these groups ranged from 13.3 to 19.5% before treatment. Twenty-four hours after administration, the parasitemia was reduced to 2.2% in group I and <0.1% in group II; parasitemia was lowered to 10.6% in group III only three hours after drug administration. The rate of sequestration in the cerebral microvessels, which was 29.4% in untreated animals, was <0.1% in groups I and II (24 hours after treatment), and 2.0% in group III (3 hours after treatment). These data clearly indicate that artesunate not only reduced parasitemia, but also reduced the rate of parasitized red blood cell (PRBC) sequestration in cerebral microvessels. In an immunohistological study, endothelial-leukocyte adhesion molecule-1 (ELAM-1) became negative in group I after treatment with artesunate, although the

presence of CD36, thrombospondin (TSP), intercellular adhesion molecule-1 (ICAM-1), IgG, and C3 in the cerebral microvessels was not altered. This is the first in vivo study to show that artesunate interferes with continued PRBC sequestration in the cerebral microvessels in cerebral malaria. By using this animal model, we may be able to evaluate strategies for management of severe malaria as well as for the development of antimalarials and vaccines to prevent and/or cure cerebral malaria.

d) Morphological effects of artemisinin in Plasmodium falciparum

In collaboration with Dr. Milhous of WRAR, ultrastructural changes induced in Plasmodium falciparum by artemisinin were studied in vitro. Two hours after administration, changes were observed in parasite mitochondria, rough endoplasmic reticulum, and nuclear envelope. At four hours, in addition to the earlier changes, nuclear membranes and, to a lesser extent, plasma membranes may form myelin figures. In addition, there may be disappearance of ribosomes, and destruction of food vacuole membranes. These changes may lead to the total disorganization of the parasites. About 30% of the parasites manifested these alterations. Electron microscope autoradiography was performed after [³H]-dihydroartemisinin and [¹⁴C]-artemisinin were administered to infected erythrocytes in vitro. These drugs consistently were located in mitochondria (Fig.5) and food vacuoles (Fig.6).

e) Human vascular endothelial cell adhesion receptors

The clinical complications associated with severe and cerebral malaria occur as a result of the intravascular mechanical obstruction of erythrocytes infected with the asexual stages of Plasmodium falciparum. In collaboration with Dr. Ockenhouse of WRAIR, we report that a primary P. falciparum-infected erythrocyte (parasitized red blood cell [PRBC] isolate from a patient with severe complicated malaria binds to cytokine-induced human vascular endothelial cells, and that this adhesion is in part mediated by endothelial leukocyte adhesion molecule 1 (ELAM-1) and vascular cell adhesion molecule 1 (VCAM-1). PRBC binding to tumor necrosis factor α (TNF- α)-activated human vascular endothelial cells is partially inhibited by antibodies to ELAM-1 and ICAM-1 and the inhibitory effects of these antibodies is additive, PRBCs selected in vitro by sequential panning on purified adhesion molecules bind concurrently to recombinant soluble ELAM-1 and VCAM-1, and to two previously identified endothelial cell receptors for PRBCs, ICAM-1, and CD36. Post-mortem brain tissue from patients who died from cerebral malaria expressed multiple cell adhesion molecules including ELAM-1 and VCAM-1 on cerebral microvascular endothelium not expressed in brains of individuals who died from other causes. These results ascribe novel pathological functions for both ELAM-1 and VCAM-1 and may help delineate alternative adhesion pathways PRBCs use to modify malaria pathology.

CONCLUSIONS:

As indicated in the section "Introduction" we have collaborated with scientists from WRAIR and AFRIMS in order to investigate morphological effects of drugs, particularly antimalarials on malarial parasites and hosts.

We performed immuno-electron microscopy to study intracellular localization and fate of liposomes and liposome-encapsulated malarial antigen after phagocytosis by macrophages, since the ability of liposomes to enhance immune responses is due to natural targeting of liposomes to macrophages. Our study indicated that liposome-encapsulated protein that is phagocytosed by macrophages can enter an intracellular compartment in which at least some of the antigenic epitopes are not degraded by lysosomal enzymes.

Our attempt to make a primate model for human cerebral malaria has been successful. PRBC sequestration and cytoadherence of knobs on PRBC to endothelial cells in cerebral microvessels were demonstrated in rhesus monkeys infected with *P. coatneyi*. These cerebral microvessels with sequestered PRBC were shown to possess CD36, TSP and ICAM-1. By using this animal model, we studied morphological effects of artemisinin on sequestered PRBC within cerebral microvessels. In addition, by using this model, we may be able to develop vaccine candidates to prevent human cerebral malaria.

REFERENCES:

1. Aikawa, M. and Beaudoin, R.L. 1969. Effects of Chloroquine in the Morphology of the Erythrocytic Stages of *Plasmodium gallinaceum*. Am. J. Trop. Med. Hyg., 18:166-181.
 2. Aikawa, M. 1972. High Resolution Autoradiography of Malarial Parasites Treated with 3H-Chloroquine. Am. J. Path., 67:277-284.
 3. Macomber, F.B., Sprinz, H. and Tousimes, A.J. 1967. Morphological effects of chloroquine on *Plasmodium berghei* in mice. Nature 214:937.
 4. Beaudoin, R.L. and Aikawa, M. 1968. Primaquine induced changes in the morphology of the exoerythrocytic stages of malaria. Science, 160:1233.
 5. Aikawa, M. and Beaudoin, R.L. 1970. *Plasmodium fallax*: High-Resolution Autoradiography of Exoerythrocytic Stages Treated With Primaquine *in vitro*. Exp. Parasitol., 27:454-463.
 6. Aikawa, M. and Beaudoin, R.L., 1968. Studies on nuclear division of a malarial parasite under pyrimethamine treatment. J. Cell Biol. 39:749.
-

7. Jiang, J.B., Jacobs, G. and Aikawa, M. 1985. Qinghaosu-induced changes in morphology of malarial parasites. Am. J. Trop. Med. Hyg., 34:424-428.
 8. Atkinson, C.T., Aikawa, M., Tam, L., Hui, G. and Siddiqui, W. 1987. Ultrastructural localization of Plasmodium falciparum-polypeptides protective in Aotus monkeys. J. Parasitol., 73:1235-1240.
 9. Aikawa, M. and Atkinson, C. 1990. Immuno-electron microscopy of parasites. In: Advances in Parasitology Vol. 29 (ed. J. R. Baker), Academic Press, London p. 151-214.
 10. Alving, C.R. 1987. Liposomes, In: Biophysics to Therapeutics (ed by Ostro, M.J.), New York, Marcel Debber, p. 195-218.
 11. Aikawa, M. 1988. Human cerebral malaria. Am. J. Trop. Med. Hyg. 39:1-10.
 12. Aikawa, M., Iseki, M., Barnwell, J.W., Taylor, D., Oo, M.M. and Howard, R.J. 1990. The pathology of human cerebral malaria. Am. J. Trop. Med. Hyg., 43:30-37 (Suppl.).
-

13. Riganti, M. Pongponratn, E., Tegoshi, T., Looareesuwan, S., Punpoowong, B. and Aikawa, M. 1990. Human cerebral malaria in Thailand. Immunology Letters 25:199-206.


 14. Desowitz, R.S., Miller, L.H., Buchanan, R.D. and Permpanick, B. 1969. The sites of deep vascular schizogony in Plasmodium coatneyi malaria. Trans. Roy. Soc. Trop. Med. Hyg. 63: 198.



 15. Aikawa, M., Brown, A., Smith, C.D., Tegoshi, T., Howard, R., Hasler, T.H., Ito, Y., Perry, G., Collins, W.E., and Webster, K.I., 1992. A primate model for human cerebral malaria in P.coatneyi-infected rhesus monkeys. Am. J. Trop. Med. Hyg. 46:391-397.
-


APPENDIX:

Figure Legend

- Fig. 1 Immunolectron micrograph of macrophages. The macrophages were fixed 6 hours after incubation with liposomes containing malaria antigen and the malaria antigen was detected by specific monoclonal antibody (pf 1B2.2) to the antigen followed by the treatment with gold-labeled second antibody. Densely labeled liposomes are present within a membrane bound vacuole. X 50,000
- Fig. 2 Electron micrograph showing cytoadherence (arrow) of PRBC knobs to endothelial cells (E) of rhesus cerebral microvessels. X 56,000
- Fig. 3 Electron micrograph showing PRBC sequestration in a cerebral microvessel of rhesus monkey. X 18,000
- Fig. 4 Immunolectron micrograph showing the presence of EBA-peptide 4 in micronemes (M) of P.falciparum merozoites. No gold particles are seen associated with rhoptries (R). X 100,000
- Fig. 5&6 Autoradiography of P.falciparum treated with ^3H -dihydroartemisinin. Silver grains are associated with mitochondria (M) (Fig.5) x 64,000 and food vacuoles (F) (Fig.6). X 39,000







4



PUBLICATIONS:

1. Atkinson, C.T., Millet, P., Collins, W.E. and Aikawa, M., Localization of circumsporozoite antigen in exoerythrocytic schizonts of Plasmodium cynomolgi. Am. J. Trop. Med. Hyg. 40:131-140, 1989.
2. Atkinson, C.T., Aley, S.B., Hollingdale, M.R. and Aikawa, M., Expression of Plasmodium berghei circumsporozoite antigen on the surface of exoerythrocytic schizonts and merozoites. Am. J. Trop. Med. Hyg. 41:9-17, 1989.
3. Yamada, M., Steketee, R., Abramowsky, C., Kida, m., Wirima, J., Heymann, D., Rabbege, J., Breman, J. and Aikawa, M., P. falciparum associated placental pathology: A light and electron microscopical and immunohistochemical study. Am. J. Trop. Med. Hyg. 41:161-168, 1989.
4. Udomsangpetch, R., Carlson, J., Wahlin, B., Helmquist, G., Ozaki, L.S., Scherf, A., Mattei, D., Mercereau-Puijalon, O., Uni, S., Aikawa, M., Berzins, K. and Perlmann, P., Reactivity of the human monoclonal antibody J3G2 with repeated sequences of three distinct Plasmodium falciparum antigens. J. Immunol. 142:3620-3626, 1989.
5. Klotz, F.W., Hadley, T.J., Aikawa, M., Leech, J., Howard, R.J. and Miller, L.H., A 60 kDa Plasmodium falciparum

protein at the moving junction formed between merozoite and erythrocyte during invasion. Mol. Biochem. Parasitol. 16:177-186, 1989.

6. Barnwell, J.W., Asch, A.S., Silverstein, R.L., Nochman, R.L., Yamada, M., Aikawa, M. and Ingravallo, P., Identification of an 88 kD membrane glycoprotein (GPIVCD36) as the receptor for a cytoadherence ligand on Plasmodium falciparum-infected erythrocytes. J. Clin. Invest. 84:765-772, 1989.
7. Cochrane, A.H., Uni, S., Maracic, M., Aikawa, M. and Nussenzweig, R., A circumsporozoite-like protein is present in micronemes of mature blood stages of malaria parasites. Exptl. Parasitol. 69:351-356, 1989.
8. Torii, M., Adams, J.H., Miller, L.H. and Aikawa, M., Release of merozoite dense granules during erythrocyte invasion by Plasmodium knowlesi. Infect. Immun. 57:3230-3233, 1989.
9. Kumar, N., Koski, G., Harada, M., Aikawa, M., and Hong, Z. Induction and localization of Plasmodium falciparum stress proteins related to the heat shock protein 70 families. Molecular & Biochem. Parasitol 48:47-58, 1991.

10. Aikawa, M., Iseki, M., Barnwell, J.W., Taylor, D., Oo, M.M. and Howard, R.J., The pathology of human cerebral malaria. Am. J. Trop. Med. Hyg., 43:30-37 (Suppl.), 1990.
11. Aikawa, M., Atkinson, C.T., Beaudoin, L.M., Sedegah, M., Charoenvit, Y. and Beaudoin, R., Localization of CS- and non-CS antigens in the sporozonic stages of Plasmodium yoelii. WHO Bulletin 68 (Suppl.): 165-171, 1990.
12. Adams, J.H., Hudson, D.F., Ward, G., Torii, M., Aikawa, M., Wellems, T.E. and Miller, L.H., The Duffy receptor is concealed in the micronemes of invasive malaria merozoites. Cell 63:141-153, 1990.
13. Aikawa, M., Torii, M., Sjolander, A., Berzins, K., Perlmann, P. and Miller, L.H., Pf155/RESA is localized in dense granules of Plasmodium falciparum, Exptl. Parasitol. 71:326-329, 1990.
14. Riganti, M. Pongponratn, E., Tegoshi, T., Looareesuwan, S., Punpoowong, B. and Aikawa, M., Human cerebral malaria in Thailand. Immunology Letters 25:199-206, 1990.
15. Hollingdale, M.R., Aikawa, M., Atkinson, C.T., Ballou, W.R., Chen, G., Li, J., Meis, J.F.G.M., Siva, B., Wright, C. and

- Zhu, J., Non-CS pre-erythrocytic protective antigens. Immunology Letters 25:71-76, 1990.
16. Pongponratn, E., Riganti, M., Looareesuwan, S., Punpoowong, B. and Aikawa, M., Microvascular sequestration in parasitized erythrocytes in human falciparum malaria: A pathological study. Am. J. Trop. Med. Hyg. 44:168-175, 1991.
 17. Verma, J.N., Wasser, N.M., Wirtz, R.A., Atkinson, C.T., Aikawa, M., and Alving, C.R., Phagocytosis, intracellular localization and surface expression of liposome-encapsulated malaria antigen in macrophage. Biochemica et Biophysica Acta. 1066:229-238, 1991.
 18. Aikawa, M., Brown, A., Smith, C.D., Tegoshi, T., Howard, R., Hasler, T.H., Ito, Y., Perry, G., Collins, W.E. and Webster, K.I. A primate model for human cerebral malaria: P. coatneyi-infected rhesus monkeys. Am. J. Trop. Med. Hyg. 46:391-397, 1992.
 19. Atkinson, C.T., Bayne, M.T., Gordeuk, V.R., Brittenham, G.M., and Aikawa, M. Stage-specific ultrastructural effects of Desferrioxamine in Plasmodium falciparum in vitro. Am. J. Trop. Med. Hyg. 45:593-601, 1991.

20. Nakamura, K., Hasler, T., Morehead, K., Howard, R.J. and Aikawa, M. Plasmodium falciparum-infected erythrocyte receptor(s) for CD36 and thrombospondin are restricted to knobs on the erythrocyte surface. J. Histochem, Cytochem 40:1419-1422, 1992.

21. Atkinson, C. T., Hollingdale, M.R. and Aikawa, M. Localization of a 230 kDa parasitophorous vacuole membrane antigen of Plasmodium berghei exoerythrocytic schizonts (LSA-2) by immunoelectron and confocal laser scanning microscopy. Am. J. Trop. Med. Hyg. 46:533-537, 1992.

22. Nagatake, T., Thuc, H.V., Tegoshi, T., Rabbege, J. and Aikawa, M., Pathology of falciparum malaria in Vietnam. Am J. Trop. Med. Hyg. 47:259-264, 1992.

23. Aikawa, M., Brown, A.E., Smith, C.D., Tegoshi, T., Howard, R.J., Hasler, T.H., Ito, Y., Collins, W.E., and Webster, H.K., Plasmodium coatneyi-infected rhesus monkeys: a primate model for human cerebral malaria. Memorias do Instituto Osvald Cruz (In Press).

24. Sim, K.L., Toyoshima, T., Haynes, J.D. and Aikawa, M., Localization of the 175 kildalton erythrocyte

- binding antigen in micronemes of Plasmodium falciparum merozoites. Mol. Biochem. Parasitol 51:157-160, 1992.
25. Torii, M., Aikawa, M., Srivastava, I.K., Schmidt, M., Nakamura, K. and Perrin, L.H. Expression and subcellular localization of Plasmodium falciparum aldolase. J. Protozool Res. 2:10-15, 1992.
26. Hollingdale, M.R., Aley, S.B., Atkinson, C.T. and Aikawa, M. Malarial circumsporozoite proteins are processed during and after sporozoite invasion of hepatocytes. Allergy and Immunol. 9:173-179, 1990.
27. Ockenhouse, C.F., Tegoshi, T., Maeno, Y., Benjamin, C., Ho, M., Aikawa, M. and Lobb, R.R., Human vascular endothelial cell adhesion receptors for Plasmodium falciparum-infected erythrocytes: Role for ELAM-1 and VCAM-1, J. Exp. Med. 176:1183-1189, 1992.
28. Rawlings, D.J., Fujioka, H., Keister, D.B., Aikawa, M., and Kaslow, D.C., α -tubulin II, a male specific protein in Plasmodium falciparum. Mol. Biochem. Parasitol 56:239-250, 1992.
29. Torii, M., Nakamura, K., Sieber, K.P., Miller, L.H. and Aikawa, M., Penetration of the mosquito (Aedes aegypti)

- midgut wall by the ookinetes of Plasmodium gallinaceum. J. Protozool. 39:449-454, 1992.
30. Fujioka, H., Torri, M., Barnwell, J.W. and Aikawa, M., Localization of F-actin on the caveola-vesicle complex of the erythrocytes infected with Plasmodia vivax and P. cynomolgi. JPN. J. Parasitol. 4:361-364, 1992.
31. Maeno, Y., Steketee, R.W., Nagatake, T., Tegoshi, T., Desowitz, R.S., Winima, J.J., and Aikawa, M. Immunoglobulin complex deposits in Plasmodium falciparum-infected placentas from Malawi and Papua New Guinea. Am. J. Trop. Med. Hyg. (In press) 1992.
32. Sein, K.K., Maeno, Y., Thuc, H.V., Anh, T.K. and Aikawa, M. Differential sequestration of parasitized erythrocytes in cerebrum and cerebellum in human cerebral malaria. Am. J. Trop. Med. & Hyg. (In press) 1992.
33. Kaidoh, T., Nath, J., Okoye, V. and Aikawa, M. Novel structure in pellicular complex of Plasmodium falciparum gametocytes. J. Protozool (In press) 1993.
-

34. Tegoshi, T., Udomsangpetch, R., Brown, A., Webster, H.K. and Aikawa, M. Ultrastructure of rosette formation by Plasmodium coatneyi-infected erythrocytes of rhesus monkey. Experimental Parasitol. (In press) 1992.

FINAL REPORT
Contract No. DAMD17-90-C-0010

Personnel Paid: 100% salary & fringe - Ana Milosaveljevic
3% salary & fringe - Masamichi Aikawa