

UNCLASSIFIED

AD NUMBER
ADB263413
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies only; Proprietary Info.; Jun 2000. Other requests shall be referred to US Army Medical Research and Materiel Command, 504 Scott St., Fort Detrick, MD 21702-5012.
AUTHORITY
USAMRMC ltr, 21 Feb 2003

THIS PAGE IS UNCLASSIFIED

AD _____

Award Number: DAMD17-99-1-9134

TITLE: p19 ARF-p53 Tumor Suppressor Pathway During Oncogene-
Induced Apoptosis and Senescence

PRINCIPAL INVESTIGATOR: Jainping Jin, Ph.D.
Scott Lowe, Ph.D.

CONTRACTING ORGANIZATION: Cold Spring Harbor Laboratory
Cold Spring Harbor, New York 11724

REPORT DATE: June 2000

TYPE OF REPORT: Annual Summary

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Distribution authorized to U.S. Government agencies only (proprietary information, Jun 00). Other requests for this document shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott Street, Fort Detrick, Maryland 21702-5012.

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

20010216 061

NOTICE

USING GOVERNMENT DRAWINGS, SPECIFICATIONS, OR OTHER DATA INCLUDED IN THIS DOCUMENT FOR ANY PURPOSE OTHER THAN GOVERNMENT PROCUREMENT DOES NOT IN ANY WAY OBLIGATE THE U.S. GOVERNMENT. THE FACT THAT THE GOVERNMENT FORMULATED OR SUPPLIED THE DRAWINGS, SPECIFICATIONS, OR OTHER DATA DOES NOT LICENSE THE HOLDER OR ANY OTHER PERSON OR CORPORATION; OR CONVEY ANY RIGHTS OR PERMISSION TO MANUFACTURE, USE, OR SELL ANY PATENTED INVENTION THAT MAY RELATE TO THEM.

LIMITED RIGHTS LEGEND

Award Number: DAMD17-99-1-9134
Organization: Cold Spring Harbor Laboratory
Location of Limited Rights Data (Pages):

Those portions of the technical data contained in this report marked as limited rights data shall not, without the written permission of the above contractor, be (a) released or disclosed outside the government, (b) used by the Government for manufacture or, in the case of computer software documentation, for preparing the same or similar computer software, or (c) used by a party other than the Government, except that the Government may release or disclose technical data to persons outside the Government, or permit the use of technical data by such persons, if (i) such release, disclosure, or use is necessary for emergency repair or overhaul or (ii) is a release or disclosure of technical data (other than detailed manufacturing or process data) to, or use of such data by, a foreign government that is in the interest of the Government and is required for evaluational or informational purposes, provided in either case that such release, disclosure or use is made subject to a prohibition that the person to whom the data is released or disclosed may not further use, release or disclose such data, and the contractor or subcontractor or subcontractor asserting the restriction is notified of such release, disclosure or use. This legend, together with the indications of the portions of this data which are subject to such limitations, shall be included on any reproduction hereof which includes any part of the portions subject to such limitations.

THIS TECHNICAL REPORT HAS BEEN REVIEWED AND IS APPROVED FOR PUBLICATION.

Kathleen M. M. 11/12/01

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE June 2000	3. REPORT TYPE AND DATES COVERED Annual Summary (15 May 99 - 15 May 00)	
4. TITLE AND SUBTITLE p19 ARF-p53 Tumor Suppressor Pathway During Oncogene-Induced Apoptosis and Senescence			5. FUNDING NUMBERS DAMD17-99-1-9134	
6. AUTHOR(S) Jainping Jin, Ph.D. Scott Lowe, Ph.D.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Cold Spring Harbor Laboratory Cold Spring Harbor, New York 11724 E-MAIL: jin@cshl.org			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012			10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES This Report Contains Colored Photographs				
12a. DISTRIBUTION / AVAILABILITY STATEMENT DISTRIBUTION STATEMENT: Distribution authorized to U.S. Government agencies only (proprietary information, Jun 00). Other requests for this document shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott Street, Fort Detrick, Maryland 21702-5012.				12b. DISTRIBUTION CODE
13. ABSTRACT (Maximum 200 Words) <p>The primary objective of this project is to provide new insights into the role of the p53 and ARF tumor suppressors in cancer development and therapy. During the first year of this study, we found that different oncogenes can promote different p53 post-translational modifications. Oncogene ras promotes phosphorylation of p53 on Serine 15, however, E1A does not. These data suggested that different oncogenes induce different phenotypes through different p53 post-translational modifications. It seems that p19^{ARF} is required for p53 modifications, since p53 can not be phosphorylated on Ser 15 following Ras overexpression in ARF null cells. We also examined the impact of INK4a/ARF mutations on tumor development and therapy using the Eμ-myc transgenic mouse. Our results clearly showed that inactivation of the INK4a/ARF locus accelerated Myc-induced lymphomagenesis, leading to massively deseminated lymphomas that displayed markedly reduced apoptosis. In collaboration with C. Sherr, we tried to produce different monoclonal antibodies to p19^{ARF}. Right now, we are doing the secondary screens to determine the antibodies specificity.</p>				
14. SUBJECT TERMS Breast Cancer			15. NUMBER OF PAGES 19	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

J.J. 06-12-00

 Where copyrighted material is quoted, permission has been obtained to use such material.

 Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

 Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

X In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and use of Laboratory Animals of the Institute of Laboratory Resources, national Research Council (NIH Publication No. 86-23, Revised 1985).

X For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

 In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

 In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

N/A In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

Jianping Jin 06-12-00

PI - Signature

Date

Table of Contents

Cover.....	
SF 298.....	II
Foreword.....	III
Introduction.....	1
Body.....	1
Key Research Accomplishments.....	4
Reportable Outcomes.....	5
Conclusions.....	5
References.....	5
Appendices.....	5

Introduction

Breast cancer is the second leading cause of cancer death in North American Women. Mutations in the p53 tumor suppressor gene are the most common genetic changes found in breast cancer so far. It has been suggested that p53 plays a critical role in the development of this disease, however, the signals triggering p53 in suppressing tumor growth remain poorly defined. We found that oncogenes induce p53 through a signal transduction pathway that requires the presence of another tumor suppressor, p19^{ARF}. This pathway, when activated, directs the cells to apoptosis or senescence. The primary objective of this project is to provide new insights into the action of the tumor suppressor p53 and p19^{ARF} in cancer development. Our original studies took advantage of mouse cells with a defined genetic background to analyze the relevance of single genes in the oncogene-p19^{ARF}-p53 pathway. We propose to compare this highly controlled system to normal human breast cells in order to determine if the pathway is conserved in human cells and thus potentially relevant to breast cancer. Furthermore, by comparing and contrasting normal versus oncogene expressing cells, in these systems, we will be able to get insight to the mechanisms by which p19^{ARF} is involved in p53 activation. Finally, a better understanding of the oncogene-p19^{ARF}-p53-apoptosis pathway could allow us to exploit new strategies to enhance the chemo- and radio-sensitivity of breast cancer cells. The research progress of the first year on this project is reported as follows.

Body

1. Oncogene signaling to p53 (Aims 2 and 3)

The primary objective of this study is to provide new insights into the role of the p53 and ARF tumor suppressors in breast cancer development and therapy.

p53 promotes cell cycle arrest or apoptosis in response to different stimuli, including DNA damage and mitogenic stimulation. Following DNA damage, signaling is mediated, at least in part, by a kinase that phosphorylates p53 on Serine 15, thereby stabilizing the protein by preventing its association with its negative regulator Mdm2.

In our original proposal, we presented data showing that the adenovirus E1A oncogene activates p53 through a mechanism distinct from DNA damage, involving cancellation of Rb function and activation of the ARF tumor suppressor, leading the cells to apoptosis. ARF-null MEFs expressing E1A are incapable of activating the p53 response and are more resistant to apoptosis following serum depletion or adriamycin treatment, compared to wild type MEFs. Reintroduction of ARF restores p53 accumulation and sensitizes cells to radiation and chemotherapy.

We decided to test whether ARF induction is a common event following oncogene overexpression. Our recent data indicated that the cellular ras oncogene, by activating MAP kinase pathway, also induces p19^{ARF} and cooperates with p53 to promote premature senescence. (Lin, de Stanchina, Ferbeyre and Lowe unpublished). In this case, ARF null cells expressing ras can still induce p53 but exhibit an impaired p53 response and do not undergo premature senescence.

We have also been able to show that different oncogenes can promote different p53 post-translational modifications. In particular, we found that E1A does not promote phosphorylation of p53 on Serine 15, but Ras does it (de Stanchina and Lowe, unpublished). It seems that different oncogenic stimuli, which induce different outcomes, also are associated with different p53 modifications, raising the possibility that these differences in modifications alert p53 to the type of insult that has occurred and subtly influence the nature of the p53 response. It is interesting to note that ARF seems to be required for these functions, since in ARF null cells, p53 can not be phosphorylated on Ser 15 following Ras overexpression (de Stanchina and Lowe, unpublished).

Our current data imply that oncogenes (in part through ARF) and DNA damage cooperate to induce apoptosis or permanent cell cycle arrest and indicate that loss of ARF function may promote tumor progression and chemoresistance by disabling p53.

p53 mutations have been associated with aggressive cancers, poor prognosis and drug resistance in human patients. In principle, tumors with INK4a/ARF mutations might also display aggressive characteristics owing to extragenic defects in the p53 pathway. To test this, we examined the impact of INK4a/ARF mutations on tumor development and therapy using the E μ -myc transgenic mouse. These mice constitutively express c-myc in the B cell lineage and develop B cell lymphoma and associated leukemia. The

E μ -myc transgenic lymphoma model was used to generate genetically defined tumors with mutations in the INK4a/ARF or p53 genes. Thus it was possible to monitor the rate of formation of p53 or INK4a/ARF null lymphomas, their invasiveness and their resistance to chemotherapy *in vitro* and *in vivo*.

Our results (Schmitt et al, 1999) clearly showed that inactivation of the INK4a/ARF locus accelerated Myc-induced lymphomagenesis, leading to massively deseminated lymphomas that displayed markedly reduced apoptosis. The latency, pathology and growth characteristics were indistinguishable from Myc-induced lymphomas lacking p53. Indeed, the p53 activity was compromised in INK4a/ARF null lymphoma cells despite the fact that these cells harbor wild-type p53 genes. These results demonstrated that genetic interactions between the p53 and INK4a/ARF locus are important during tumorigenesis and predict that disruption of the INK4a/ARF locus, like p53 loss, will be a negative prognostic marker in human tumors.

2. Production of highly specific monoclonal antibodies against the N-terminal portion of ARF(Aim 1)

In tumors, mutations affecting ARF occur primarily in the region coding for the C-terminus domain (exon 2), possibly leading to the formation of truncated isoforms. These putative forms would go undetected by the only currently available anti-ARF antibodies, which are directed against the C-terminus of the protein. For this reason we proposed to raise antibodies against the amino terminal portion of ARF, which has been shown to be necessary and sufficient to induce cell cycle arrest and to interact with p53 and/or Mdm2. In collaboration with C. Sherr, who provided us with the synthetic peptides (amino acids 1-64 of the mouse protein), we produced and analyzed different monoclonal antibodies. The CSHL Monoclonal Service Facility performed all the animal work, the fusions, single cell cloning and initial screens. I am currently doing the secondary screens to determine the antibodies specificity. I will initially use the antibodies for routine characterization studies, such as western blot analysis and immunoprecipitations, to detect ARF levels in different cell lines and to investigate the presence and abundance of the truncated isoforms in tumor derived cell lines and in clinical specimens.

3. ARF regulation by E1A in cancer chemotherapy

We have shown that ARF is involved in oncogene-induced apoptosis, by stabilizing p53 level. Next question we hope to address is how ARF is regulated by oncogene. We know E1A can induce ARF expression at messenger level and N-terminal function of E1A is highly relative to E1A-induced ARF expression. With the ARF promoter reporter gene construct in hand, we will be able to see how ARF is specially activated by oncogene. We also found that E1A can induce apoptosis in mouse skin fibroblasts (MSFs) as in MEFs. N-terminal deletion of E1A is defective in E1A-induced apoptosis in MSFs. Since MSFs cells have more homogeneous genetic background, with the new microarray technology, we will try to identify new genes that regulate ARF expression in oncogene-induced ARF-p53-apoptosis pathway.

Key Research Accomplishments

1. We found that different oncogenes can promote different p53 post-translational modifications. Oncogene Ras promotes phosphorylation of p53 on Serine 15, however, E1A does not. These data suggested that different oncogenes induce different phenotypes by different p53 post-translational modifications. It seems that p19^{ARF} is required for p53 modifications.
2. We also examined the impact of INK4a/ARF mutations on tumor development and therapy using the E μ -myc transgenic mouse. Our results clearly showed that inactivation of the INK4a/ARF locus accelerated Myc-induced lymphomagenesis, leading to massively deseminated lymphomas that displayed markedly reduced apoptosis.
3. In collaboration with C. Sherr, we tried to produce and analyze different monoclonal antibodies to p19^{ARF}. Right now, we are doing the secondary screens to determine the antibodies specificity.

Reportable Outcomes

Clemens A. Schmitt, Mila E. McCurrach, Elisa de Stanchina, Rachel R. Wallace-Brodeur, and Scott W. Lowe, 1999, INK4a/ARF mutations accelerate lymphomagenesis and promote chemoresistance by disabling p53. *Gene & Development* Vol. 13, No. 20, pp. 2670-2677.

Conclusions

In conclusion, different oncogenes can promote different p53 post-translational modifications. It seems that p19^{ARF} is essential to these p53 modifications. Whether special p53 modifications are relative to different cancer development remains to be studied. In collaboration with Dr. Schmitt, Dr. de Stanchina published a paper about the mouse model of p19^{ARF}-p53-apoptosis pathway. Our results clearly showed that inactivation of the INK4a/ARF locus accelerated Myc-induced lymphomagenesis, leading to massively deseminated lymphomas that displayed markedly reduced apoptosis. We are continuing to work on this project, with a particular emphasis on how E1A induces ARF.

References

Clemens A. Schmitt, Mila E. McCurrach, Elisa de Stanchina, Rachel R. Wallace-Brodeur, and Scott W. Lowe, 1999, INK4a/ARF mutations accelerate lymphomagenesis and promote chemoresistance by disabling p53. *Gene & Development* Vol. 13, No. 20, pp. 2670-2677

Appendices

Clemens A. Schmitt, Mila E. McCurrach, Elisa de Stanchina, Rachel R. Wallace-Brodeur, and Scott W. Lowe, 1999, INK4a/ARF mutations accelerate lymphomagenesis and promote chemoresistance by disabling p53. *Gene & Development* Vol. 13, No. 20, pp. 2670-2677.

***INK4a/ARF* mutations accelerate lymphomagenesis and promote chemoresistance by disabling p53**

Clemens A. Schmitt, Mila E. McCurrach, Elisa de Stanchina, Rachel R. Wallace-Brodeur, and Scott W. Lowe

Cold Spring Harbor Laboratory, Cold Spring Harbor, New York 11724 USA

INK4a/ARF mutations accelerate lymphomagenesis and promote chemoresistance by disabling p53

Clemens A. Schmitt, Mila E. McCurrach, Elisa de Stanchina, Rachel R. Wallace-Brodeur, and Scott W. Lowe¹

Cold Spring Harbor Laboratory, Cold Spring Harbor, New York 11724 USA

The *INK4a/ARF* locus encodes upstream regulators of the retinoblastoma and p53 tumor suppressor gene products. To compare the impact of these loci on tumor development and treatment response, the *Eμ-myc* transgenic lymphoma model was used to generate genetically defined tumors with mutations in the *INK4a/ARF*, *Rb*, or *p53* genes. Like *p53* null lymphomas, *INK4a/ARF* null lymphomas formed rapidly, were highly invasive, displayed apoptotic defects, and were markedly resistant to chemotherapy *in vitro* and *in vivo*. Furthermore, *INK4a/ARF*^{-/-} lymphomas displayed reduced *p53* activity despite the presence of wild-type *p53* genes. Consequently, *INK4a/ARF* and *p53* mutations lead to aggressive tumors by disrupting overlapping tumor suppressor functions. These data have important implications for understanding the clinical behavior of human tumors.

[Key Words: *INK4a/ARF* locus; lymphomagenesis; chemo resistance; p53]

Received August 10, 1999; revised version accepted August 31, 1999.

Mutations in the *p53* tumor suppressor gene and at the *INK4a/ARF* locus are the two most frequent genetic lesions identified in human tumors (for reviews, see Haber 1997; Ruas and Peters 1998). *p53* is a sequence-specific DNA-binding protein that can induce cell-cycle arrest or apoptosis in response to pathological insults such as DNA damage and expression of mitogenic oncogenes (Kastan et al. 1991, 1992; Lowe and Ruley 1993; Hermeking and Eick 1994; Serrano et al. 1997; for reviews, see Giaccia and Kastan 1998; Prives 1998). As a consequence, inactivation of *p53* can promote oncogenic transformation and resistance to many anticancer agents (for reviews, see Giaccia and Kastan 1998; Prives 1998; Wallace-Brodeur and Lowe 1999). The *INK4a/ARF* locus encodes two tumor suppressors, designated p16^{INK4a} and p19^{ARF}. p16^{INK4a} is a cyclin-dependent kinase inhibitor that acts upstream of the retinoblastoma (*Rb*) protein to promote cell-cycle arrest (Serrano et al. 1993; for reviews, see Haber 1997; Ruas and Peters 1998). p19^{ARF} is translated in an alternative reading frame from p16^{INK4a} and activates *p53* by interfering with its negative regulator, Mdm2 (Kamijo et al. 1998; Pomerantz et al. 1998; Stott et al. 1998; Zhang et al. 1998; see also Tao and Levine 1999; Weber et al. 1999; Zhang and Xiong 1999). Consequently, *INK4a/ARF* mutations can disable both the *Rb* and *p53* tumor suppressor pathways.

Recent studies indicate that p19^{ARF} acts as an essential intermediate in oncogene signaling to *p53* (Bates et al. 1998; de Stanchina et al. 1998; Palmero et al. 1998; Pomerantz et al. 1998; Zindy et al. 1998; for review, see Sherr 1998). For example, oncogenes such as *E1A* or *c-myc* induce *ARF* message and protein in normal mouse embryo fibroblasts, which correlates with their ability to activate *p53* and promote apoptosis. In contrast, these oncogenes fail to activate *p53* in *ARF*-null cells, and promote proliferation without substantial apoptosis (de Stanchina et al. 1998; Zindy et al. 1998). Together, these studies indicate that p19^{ARF} acts as part of a *p53*-dependent fail-safe mechanism to counter hyperproliferative signals. Interestingly, p19^{ARF} is not induced by DNA damage (Kamijo et al. 1997; Stott et al. 1998) but can cooperate with DNA damaging agents to induce apoptosis in oncogene expressing cells (de Stanchina et al. 1998). These studies predict that disruption of *ARF*, or the *INK4a/ARF* locus, should cooperate with mitogenic oncogenes during tumor development, in part, by disabling *p53*.

p53 mutations have been associated with aggressive cancers, poor prognosis, and drug resistance in human patients (Schmitt and Lowe 1999; Wallace-Brodeur and Lowe 1999). In principle, tumors with *INK4a/ARF* mutations might also display aggressive characteristics owing to extragenic defects in the *p53* pathway. To test this, we examined the impact of *INK4a/ARF* mutations on tumor development and therapy using the *Eμ-myc* transgenic mouse. These mice constitutively express *c-Myc*

¹Corresponding author.
E-MAIL lowe@cshl.org; FAX (516) 367-8454.

in the B-cell lineage and develop B-cell lymphoma with associated leukemia (Adams et al. 1985; Adams and Cory 1991). This model was chosen for several reasons. First, because Myc induces p19^{ARF} and activates p53 in cultured fibroblasts (Zindy et al. 1998), *Eμ-myc* transgenic mice provide a relevant setting for comparing the impact of *INK4a/ARF* and *p53* mutations on tumor behavior. Second, *Eμ-myc* lymphomas/leukemias are easily monitored by lymph-node palpation or blood smears, a property that facilitates studies examining tumor responses to therapy. Finally, essentially pure tumor cells can be isolated from lymph nodes and studied ex vivo or expanded in genetically matched nontransgenic recipients. The tractable nature of this model is in stark contrast to human systems, which suffer from difficulties in obtaining well-characterized and comparable clinical material.

Results

Loss of the *INK4a/ARF* locus accelerates lymphomagenesis similarly to loss of *p53*

To generate lymphomas with defined alterations, we crossed the *Eμ-myc* transgenic to mice heterozygous for germ-line deletions in the *Rb* (*Rb*^{+/-}), *INK4a/ARF* (*INK4a/ARF*^{+/-}), or *p53* (*p53*^{+/-}) genes (Jacks et al. 1992, 1994; Serrano et al. 1996). Of note, the *INK4a/ARF*^{+/-} animals harbor deletions that disrupt both p19^{ARF} and p16^{INK4a}, thereby recapitulating the common gross deletions seen in human tumors (Haber 1997; Ruas and Peters 1998). The onset of *Eμ-myc* lymphomas in *Rb*^{+/-} animals was variable (Fig. 1A; green, b) and only slightly accelerated relative to that observed in the wild-type background (hereafter referred to as control) (Fig. 1A; black, a). In contrast, the onset of *Eμ-myc* lymphomas in *INK4a/ARF*^{+/-} and *p53*^{+/-} animals (Fig. 1A; blue, c, and red, d) was highly reproducible and greatly accelerated compared with controls ($P < 0.0001$ each); the timing of lymphoma development in *INK4a/ARF*^{+/-} and *p53*^{+/-} mice was virtually identical. Cell surface staining confirmed that all lymphomas were of the B-cell lineage (B220⁺; Thy1.2⁻), whereas the distribution of pre-B (IgM⁻) and B (IgM⁺) was similar between the genotypes.

These data imply that p53 and products of the *INK4a/ARF* locus limit Myc-induced lymphomagenesis. Concordantly, *Eμ-myc* lymphomas arising in the *p53*^{+/-} and *INK4a/ARF*^{+/-} animals invariably lost the wild-type *p53* or *INK4a/ARF* allele (93.8% and 88.2%, respectively) (Fig. 1B). Hence, these lymphomas were either *p53*-null (*p53*^{-/-}) or *INK4a/ARF*-null (*INK4a/ARF*^{-/-}). Virtually all control (6 out of 7), *Rb*^{+/-} (4 out of 4), and *INK4a/ARF*^{+/-} (9 out of 9) tumors retained wild-type p53 as indicated by RT-PCR and sequencing of *p53* exons 4–8 (data not shown). The one *p53* mutation identified (H190R in mouse; H193R in human) corresponds to a mutation observed in B-cell leukemias and a Burkitt's lymphoma (Beroud and Soussi 1998). Deletions of the *INK4a/ARF* locus were noted in 20% of control and *Rb*^{+/-} tumors (4 out of 20) but never in *p53*^{+/-} tumors (0 out of 10) (data not shown). In no instance did lympho-




Figure 1. Tumor development in *Eμ-myc* transgenic mice. (A) Lymphoma incidence in *Eμ-myc* transgenic mice in the wild-type (control) background ($n = 65$; black, a) and in mice heterozygous for *Rb* ($n = 39$; green, b), *INK4a/ARF* ($n = 41$; blue, c), *p53* ($n = 73$; red, d), and *INK4a/ARF*; *p53* double heterozygotes ($n = 11$; orange, e). By day 70, all *p53*^{+/-} and *INK4a/ARF*^{+/-} mice developed lymphoma, whereas >75% of *Rb*^{+/-} and control mice remained tumor free. (B) Matched normal (N) and tumor (T) DNA were isolated from tail and lymph nodes and analyzed by allele-specific PCR for the targeted gene ([m] mutated allele; [wt] wild-type allele). Shown are results from five *Eμ-myc* transgenic mice in each genetic background. Note that tumors arising in the *INK4a/ARF*^{+/-}; *p53*^{+/-} double heterozygotes invariably lost the wild-type *p53* allele but never the *INK4a/ARF* allele.

mas arising in *Rb*^{+/-} animals lose the wild-type *Rb* allele (Fig. 1B), confirming that *Rb* loss has a minimal impact on Myc-induced lymphomagenesis. Because *Rb* and p16^{INK4a} loss should produce similar phenotypes (Haber 1997; Ruas and Peters 1998), these data imply that p19^{ARF} is crucial for suppressing Myc-induced lymphomagenesis. The onset of *Eμ-myc* lymphomas is markedly accelerated in *ARF*-deficient mice (Eischen et al. 1999).

Loss of *INK4a/ARF* or *p53* promotes lymphoma spreading into visceral organs

INK4a/ARF^{+/-} and *p53*^{+/-} lymphomas were highly invasive and infiltrated into various nonlymphoid organs. For example, in mice bearing *INK4a/ARF*^{+/-} and *p53*^{+/-} lymphomas, we observed extensive periportal invasion and spreading of lymphoma cell clusters throughout the liver parenchyma and massive malignant pulmonary infiltration as consolidated aggregation of large mononuclear cells and within the distended interstitial capillaries (Fig.

2). Also, neoplastic cells accumulated in the submucosa of the urinary bladder, within the kidneys, in the serosal and mesenteric surfaces of the gastrointestinal tract, and along the meninges. In contrast, tumors in mice bearing control or *Rb*^{+/-} lymphomas showed little systemic infiltration or remained localized to the lymph nodes and blood compartment despite a similarly large tumor burden. The invasive behavior of *INK4a/ARF*^{-/-} and *p53*^{-/-} tumors was reproduced following transplantation of the tumors into syngenic recipients (data not shown) and is indicative of a highly aggressive disease.

INK4a/ARF^{-/-} lymphomas show an apoptotic defect but no genomic instability

p53 mutations can decrease cell death, increase proliferation, and produce chromosomal instability depending on context (Schmitt and Lowe 1999; Wallace-Brodeur and Lowe 1999). To determine the impact of *INK4a/ARF* and *p53* mutations on these characteristics, we examined apoptosis, mitotic index, and DNA content in control, *INK4a/ARF*^{-/-} and *p53*^{-/-} lymphomas. As revealed by histological staining and TUNEL (terminal deoxynucleotidyl transferase dUTP nick end labeling), control lymphomas contained large numbers of apoptotic cells that clustered (Fig. 3A). Apoptosis was much reduced in *INK4a/ARF*^{-/-} or *p53*^{-/-} lymphomas, and the apoptotic cells that appeared were isolated. Furthermore, primary *INK4a/ARF*^{-/-} and *p53*^{-/-} lymphoma cells explanted into culture survived much better than controls (Fig. 3B). The mitotic index (Fig. 3C) and S-phase fraction (Fig. 3D) of all lymphoma types analyzed were similar, implying that *INK4a/ARF* or *p53* mutations did not affect the proliferation rate. DNA content analysis revealed one notable difference: Whereas most of the *p53*^{-/-} tumors were aneuploid (10 out of 12), most control and *INK4a/ARF*^{-/-} tumors remained diploid (13 out of 14 and 13 out of 14, respectively) (Fig. 3D). Together, these data demonstrate that highly aggressive lymphomas can occur in the absence of chromosomal instability and imply that the aggressive nature of *INK4a/ARF*^{-/-} and *p53*^{-/-} lymphomas is due to an apoptotic defect.




Figure 2. Invasiveness of *Eμ-myc* lymphomas in liver and lung (H.E. staining, 200×). Representative examples of control, *Rb*^{+/-}, *INK4a/ARF*^{-/-}, and *p53*^{-/-} lymphomas are shown. Note the malignant embolus in the pulmonary vessel of the control—nonetheless, the lung itself remained tumor free. The relative congestion of the *Rb*^{+/-} lung is a postmortem artifact.




Figure 3. Analysis of apoptosis, proliferation, and chromosomal stability in *Eμ-myc* lymphomas. (A) Apoptosis in situ (lymph nodes) was visualized by HE staining and TUNEL. The reduced apoptotic rate observed in *INK4a/ARF*^{-/-} tumors is consistent with a similar defect observed in the ocular lens of *Rb*^{-/-}; *INK4a/ARF*^{-/-} embryos (Pomerantz et al. 1998). (B) Viability of control (●), *INK4a/ARF*^{-/-} (○), and *p53*^{-/-} (■) lymphoma cells as measured by trypan blue exclusion after explanting onto feeder cells. (C) Proliferation as estimated by the percentage of mitotic figures in HE-stained lymphoma sections. (D) DNA content analysis of primary *Eμ-myc* lymphoma. The S-phase fractions of control (30.25% ± 8.31, *n* = 9) and *INK4a/ARF*^{-/-} (29.98% ± 6.39, *n* = 10) were virtually identical, whereas sub-G1 fractions of control (3.26% ± 3.17) and *INK4a/ARF*^{-/-} (0.30% ± 0.56) lymphomas were significantly different (*P* = 0.0097). Note that sub-G1 assessment recognizes only late apoptotic cells and gives lower estimates than TUNEL. Calculations of S-phase and sub-G1 fraction in *p53*^{-/-} lymphomas were impossible due to aneuploidy. Representative profiles are shown. Note that low frequency of aneuploidy in control and *INK4a/ARF*^{-/-} lymphomas (1 of 14 and 1 of 14, respectively) is consistent with the overall *p53* mutation rate we observed in these tumors.

INK4a/ARF mutations compromise *p53* function in vivo

The remarkable similarities between *INK4a/ARF*^{-/-} and *p53*^{-/-} lymphomas suggest that these mutations disrupt

overlapping tumor suppressor functions. In agreement, *Eμ-myc* lymphomas arising in mice heterozygous for both genes (*INK4a/ARF*^{+/-}; *p53*^{+/-}) were detected at the same time as lymphomas in the *INK4a/ARF*^{+/-} and *p53*^{+/-} animals (50th percentile = 38 vs. 38 vs. 36 days to onset, respectively) [Fig. 1A; orange, e] and never displayed coincident loss of both wild-type *INK4a/ARF* and *p53* alleles (data not shown). Therefore, inactivation of both loci produces no additional advantage to *Eμ-myc* lymphomas. In cultured fibroblasts, Myc activates p53 in an ARF-dependent manner [Zindy et al. 1998]. Similarly, control *Eμ-myc* lymphomas displayed a variable but consistent increase in p53 levels and activity (as measured by expression of the p53 target p21) compared with normal splenocytes [Fig. 4A, cf. N with control]. This increase appeared dependent on the *INK4a/ARF* locus, because *INK4a/ARF*^{-/-} tumors displayed only a modest induction of p53 and virtually no increase in p21 [Fig. 4A]. This implies that *INK4a/ARF* mutations can accelerate tumor progression and impair apoptosis by compromising p53 function.

INK4a/ARF mutations reduce p53 activation following chemotherapy

The fact that p19^{ARF} can cooperate with DNA-damaging agents to induce p53 and apoptosis raises the possibility that *INK4a/ARF* mutations might compromise cancer therapy [de Stanchina et al. 1998]. To test this, we examined the impact of *INK4a/ARF* or *p53* mutations on




Figure 4. p53 levels and activity in untreated and CTX-treated *Eμ-myc* lymphomas. (A) Control (three independent tumors), *INK4a/ARF*^{-/-} (four independent tumors), and *p53*^{-/-} lymphoma lysates were probed against p53 and the p53 downstream target p21, reflecting p53's activity. Normal splenocytes (N) from nontransgenic mice were used for comparison. Tubulin (Tub) was used to verify protein loading. (B) Control, *INK4a/ARF*^{-/-}, and *p53*^{-/-} lymphoma cells were isolated from lymph nodes of untreated animals (NT) or 4 hr after CTX treatment (T1 and T2) and analyzed as above. For each tumor type, T1 and T2 were derived from separate primary tumors, whereas NT and T1 represent reconstituted lymphomas derived from the same primary tumor.

drug-induced responses in reconstituted lymphomas or following short-term culture. Reconstituted lymphomas were produced following intravenous injection of primary lymphoma cells into syngenic (nontransgenic) recipients, thereby eliminating the possibility that secondary malignancies might complicate scoring tumor responses. Importantly, these lymphomas were histopathologically identical to their respective primary tumors (data not shown). In control lymphomas, p53 and p21 levels were dramatically increased 4 hr after treatment with cyclophosphamide (CTX), an alkylating agent used to treat human leukemia and lymphoma [Fig. 4B]. p53 and p21 levels also increased in *INK4a/ARF*^{-/-} lymphomas, although this response was consistently reduced compared with controls. Therefore, in *Eμ-myc* lymphoma cells, *INK4a/ARF* mutations can reduce p53 activation by a DNA damaging agent.

INK4a/ARF mutations affect the short-term response to anticancer treatment

Loss of either *INK4a/ARF* or *p53* had a profound effect on drug-induced cell death *in vitro* and *in vivo*. In short-term cultures, *INK4a/ARF*^{-/-} or *p53*^{-/-} lymphomas displayed a marked resistance to mafosfamide (a CTX analog active *in vitro*) [Fig. 5A]. In peripheral blood, control animals harboring associated leukemias displayed a nearly 100-fold reduction in the white blood cell count (WBC) within 4 hr of CTX therapy, which coincided with a 6- to 10-fold accumulation of apoptotic cells [Fig. 5B,C]. In contrast, *INK4a/ARF*^{-/-} and *p53*^{-/-} leukemias took 12–24 hr to achieve a similar reduction. Apoptosis was not detectable, perhaps because the slow death rate allowed clearance of apoptotic cells before they could accumulate. In lymph nodes, control lymphomas displayed massive apoptosis 5 hr after CTX therapy, whereas the *INK4a/ARF*^{-/-} and *p53*^{-/-} lymphomas displayed substantially fewer dying cells [Fig. 5D].

INK4a/ARF mutations impair the long-term response to anticancer treatment

The ultimate determinant of drug-induced cell kill is tumor regression and the duration of remission. To assess long-term responses, animals harboring control, *INK4a/ARF*^{-/-}, or *p53*^{-/-} lymphomas were treated with CTX and monitored for remission and relapse by lymph node palpation and WBC. Control lymphomas responded extremely well to CTX treatment, and >70% remained in remission during the 100-day observation period [Fig. 6; black, a]. In stark contrast, *INK4a/ARF*^{-/-} and *p53*^{-/-} null tumors displayed an extremely poor response to CTX therapy: Despite initial responses, only 1 out of 14 *p53*^{-/-} and 4 out of 35 *INK4a/ARF*^{-/-} lymphomas remained in remission. *p53*^{-/-} tumors [Fig. 6; red, c] were the most relapse prone (50th percentile = 20 days in remission, *P* < 0.0001 compared with control), although the defect in the *INK4a/ARF*^{-/-} response [Fig. 6; blue, b] was also highly significant (50th percentile = 28 days in




Figure 5. *INK4a/ARF*, *p53*, and short-term response to chemotherapy. (A) Explanted control (●), *INK4a/ARF*^{-/-} (○), and *p53*^{-/-} (■) lymphoma cells were treated with mafosphamide (MAF). Viability was analyzed after 24 hr by trypan blue exclusion; each value was normalized to untreated controls and represents the mean ± S.D. of two independently derived tumors reproduced in duplicate. (B) Leukemic mice were treated with CTX, and blood samples were taken at the indicated times. Each WBC is relative to its pretreatment value and represents the mean ± S.D. of three independent leukemias. Symbols are as in A. (C) Same as in B, except that blood samples were ethanol-fixed and stained with the DNA fluorochrome DAPI to visualize the chromatin condensation characteristic of apoptotic cells. Each value reflects the percentage of cells with apoptotic morphology (of 200 cells counted) and represents the mean ± S.D. of three independent leukemias. Symbols are as in A. (D) HE staining and TUNEL of lymph nodes harboring control, *INK4a/ARF*^{-/-}, and *p53*^{-/-} lymphomas 5 hr after CTX treatment.

remission, $P = 0.0053$ compared with control). The response of *INK4a/ARF*^{+/-};*p53*^{-/-} double-mutant lymphomas (Fig. 6; orange, d) was virtually identical to the *p53*^{-/-} tumors (50th percentile = 20 days in remission), and the relapsed tumors never displayed loss of the wild-type *INK4a/ARF* allele (data not shown). Therefore, although *INK4a/ARF* mutations promote chemoresistance in the presence of wild-type *p53* genes, they confer no additional survival advantage once *p53* is mutated. These data demonstrate that *INK4a/ARF* mutations can compromise therapy, at least in part, by disabling *p53*.

Discussion

By comparing the properties of Myc-induced lymphomas in several genetic backgrounds, we provide compelling evidence that *INK4a/ARF* deletions can impact tumor

development and anticancer therapy by compromising *p53* function. Like *p53*^{-/-} tumors, *INK4a/ARF*^{-/-} lymphomas formed rapidly, were highly invasive, displayed apoptotic defects, and were markedly resistant to chemotherapy. Furthermore, *INK4a/ARF*^{-/-} lymphomas displayed attenuated *p53* activity despite the presence of wild-type *p53* genes. The profound impact of *INK4a/ARF* and *p53* mutations on Myc-induced lymphomagenesis indicates that the ARF-*p53* pathway contributes to oncogene-induced cell death in developing tumors and underscores the importance of this fail-safe mechanism in tumor suppression (also see Eischen et al. 1999; Jacobs et al. 1999). Furthermore, that *INK4a/ARF* mutations can compromise drug-induced cell death in *Em-myc* lymphomas implies that cooperative effects between oncogenes (in part via ARF) and DNA damage contribute to the remarkable drug sensitivity of some tumors.

The only substantial difference between *INK4a/ARF*^{-/-} and *p53*^{-/-} lymphomas was that the *INK4a/ARF*^{-/-} lymphomas remained diploid, whereas the *p53*^{-/-} lymphomas were aneuploid. This pattern is reminiscent of *ARF*^{-/-} and *p53*^{-/-} fibroblasts (Kamijo et al. 1997) and implies that p19^{ARF} does not control the *p53* functions involved in maintaining chromosome stability. Although we have not analyzed the secondary changes arising in *INK4a/ARF*^{-/-} tumors in detail, these data argue that invasive, chemoresistant lymphomas can arise in the absence of substantial chromosomal insta-




Figure 6. *INK4a/ARF*, *p53*, and long-term response to chemotherapy. Nontransgenic mice harboring reconstituted control ($n = 60$; black, a), *INK4a/ARF*^{-/-} ($n = 35$; blue, b), *p53*^{-/-} ($n = 14$; red, c), and *INK4a/ARF*^{-/-};*p53*^{-/-} ($n = 15$; orange, d) lymphomas were treated with CTX and monitored for tumor regression and relapse. Importantly, CTX is not affected by classic multidrug resistance mechanisms that might complicate drug delivery. Tumor shrinkage to nonpalpability within 6 days after treatment is considered 'remission' and creates the tumor-free population at time 0. Relapse was defined by recurrent palpable lymph node enlargement. Values were plotted in Kaplan-Meier survival curve format and presented as percentage of mice in remission over the time post-therapy. Note that the overall rate of treatment failure in control lymphomas (~25%–30%) is consistent with the combined frequency of *INK4a/ARF* and *p53* mutations we observe in these tumors.

bility. In turn, because *INK4a/ARF* mutations disable p53, the chromosomal instability observed in *p53*^{-/-} lymphomas appears dispensable for the aggressive behavior of these tumors.

More likely, the increased invasiveness and drug resistance of *INK4a/ARF*^{-/-} and *p53*^{-/-} lymphomas arises from an apoptotic defect. *INK4a/ARF*^{-/-} and *p53*^{-/-} lymphomas displayed decreased apoptosis in situ and ex vivo (see Figs. 3 and 5), whereas there was no obvious relationship between tumor-cell genotype and proliferation, as measured by mitotic index in vivo, DNA content analysis ex vivo, and proliferation properties in vitro (see Fig. 3; data not shown). These results stand in contrast to an earlier report indicating that *p53*-null *Eμ-myc* lymphomas do not display an apoptotic defect (Hsu et al. 1995). Although we cannot explain this discrepancy, it is worth noting that disruption of apoptosis using a *bcl-2* transgene is sufficient to accelerate Myc-induced tumorigenesis (Strasser et al. 1990). Moreover, ectopic expression of *bcl-2* in the control lymphoma cells used in this study has no effect on proliferation but renders these cells highly invasive and chemoresistant following transplantation into syngenic mice (C.A. Schmitt and S.W. Lowe, unpubl.).

Although *Eμ-myc* lymphomas harboring *INK4a/ARF* or *p53* mutations are defective in CTX-induced cell death, CTX therapy induces complete remissions irrespective of p53 status. We suggest that the p53-independent death is due to apoptosis, because *Eμ-myc* lymphomas expressing *Bcl-2* are completely nonresponsive to CTX therapy at the maximally tolerated dose (C.A. Schmitt and S.W. Lowe, unpubl.). In contrast to CTX, doxorubicin fails to induce remissions in *p53*^{-/-} lymphomas, although high doses induce p53-independent apoptosis in vitro (R.R. Wallace-Brodeur, M.E. McCurrach, and S.W. Lowe, unpubl.). Thus, the ability to achieve p53-independent killing depends on agent and dose. These data are consistent with the view that p53 is not an essential component of the apoptotic machinery but, rather, increases the probability that these agents trigger cell death (Lowe et al. 1993). In *Eμ-myc* lymphomas, this increased propensity for apoptosis can determine tumor cure or relapse.

Our data have important implications for the understanding of the clinical behavior of human tumors. First, they provide compelling evidence that disruption of apoptosis during tumor development can simultaneously select for chemoresistant cells. This pattern of coselection may explain why some tumors are de novo 'resistant' despite having no prior exposure to drug and why it is difficult to separate the impact of *p53* mutations on treatment sensitivity from its contribution to overall patient prognosis. Second, our results demonstrate that tumors with extragenic mutations in the p53 pathway can display properties of *p53* mutant tumors. This fundamental point is crucial for interpreting studies relating *p53* mutations to clinical parameters in human patients, which typically classify tumors strictly by p53 gene or protein status. Our data imply that a substantial number of p53 'normal' tumors would be misclassified

by this approach (e.g., those harboring *ARF* mutations) and may explain why some studies fail to correlate *p53* mutations with adverse clinical features (for review, see Brown and Wouters 1999).

This study provides the first evidence that *INK4a/ARF* mutations can have a negative impact on the outcome of cancer therapy and suggests that this defect arises from the failure of drugs to appropriately activate p53. Consequently, these data predict that disruption of the *INK4a/ARF* locus will contribute to chemoresistance in human tumors. As for *p53* mutations, it seems likely that the overall impact of *INK4a/ARF* disruption on chemoresistance will depend on additional factors, such as tissue type, agent, and the mutational background of the tumor (for review, see Wallace-Brodeur and Lowe 1999; also see Bunz et al. 1999). However, it is noteworthy that *p53* mutations are strongly associated with highly aggressive tumors and chemoresistance in human hematologic malignancies (e.g., see Elrouby et al. 1993; Diccianni et al. 1994; Fan et al. 1994; Wattel et al. 1994; Wilson et al. 1997), indicating that *Eμ-myc* lymphomas can recapitulate the behavior of human tumors. Therefore, we anticipate that this model will be useful for testing strategies to counter *p53* and *INK4a/ARF* mutations in hematologic malignancies and other cancers.

Materials and methods

Mouse strains and tumor monitoring

All animal protocols used in this study were approved by the Cold Spring Harbor Laboratory Animal Care and Use Committee. *Eμ-myc* transgenic mice (C57BL/6 inbred strain) and *Rb*^{-/-}, *INK4a/ARF*^{-/-}, and *p53*^{-/-} mice (C57BL/6 × 129/sv) were crossed, and the offspring was genotyped by allele-specific PCR (Jacks et al. 1992, 1994; Serrano et al. 1996). Transgenic mice of the F₁ generation (pooled from the different crosses) or transgenics being heterozygous for the named loci were monitored twice a week by palpation of the prescapular and cervical lymph nodes. Enlargements of at least 5 mm in the longest diameter were considered 'well palpable' and reflect malignant disease. For determining white blood cell status, blood smears and 20 μl of peripheral blood were obtained by tail artery bleeding. After ammoniumchloride hemolysis of the PBS-diluted and K₃-EDTA-anticoagulated blood sample, white blood cells were counted in a hemocytometer. Blood smears were fixed and stained using the Leukostat kit (Fisher Diagnostics). Mice having WBC > 3 × 10⁵/μl and being positive for lymphoblastic cells in the blood stream were considered 'leukemic'.

Histopathology

Animals harboring control, *Rb*^{+/-}, *INK4a/ARF*^{-/-}, and *p53*^{-/-} lymphomas were sacrificed when prescapular lymph nodes reached a well-palpable size. Paraffin-embedded (7 μm), 4% neutral-buffered formalin-fixed tissue sections derived from lymph nodes and lung and liver specimens were stained with hematoxylin-eosin (HE) to evaluate apoptotic nuclear morphology and invasiveness of lymphoma cells into visceral organs.

Lymphoma characterization, LOH analysis, and RT-PCR sequencing

After CO₂ euthanasia, lymph nodes were dissected, minced in

PBS, and filtered through a 35- μ m nylon mesh. Single cell suspensions of freshly harvested lymphomas were immunophenotyped by flow cytometry using antibodies directed against Thy-1.2, B220, and IgM (Pharmingen). Pre-B-cell lymphomas are Thy-1.2⁻, B220⁺, and IgM⁻, whereas mature B-cell lymphomas are Thy-1.2⁻, B220⁺, and IgM⁺. To determine the mutational status of various genes, primary lymphoma cells were subjected to short-term culturing to eliminate normal cell contamination. Loss of the remaining wild-type allele [loss of heterozygosity (LOH)] in tumors arising in mice being heterozygous for an indicated tumor suppressor locus was detected by allele-specific PCR [Jacks et al. 1992, 1994; Serrano et al. 1996]. Exons 4–8 of the *p53* gene were sequenced by dyc termination in an automated sequencer (Perkin-Elmer) after reverse transcription (SuperScript, GIBCO BRL) and PCR amplification of lymphoma cell total RNA. Finally, the gross integrity of the *INK4a/ARF* locus was assessed using PCR of exons 1 β and exon 2 in a multiplex PCR reaction harboring primers to a positive control.

Lymphoma cell culture and in vitro treatment

Single cell suspensions of freshly extracted lymphoma cells (see above) were plated on irradiated (30 Gy) feeder layer (10^6 NIH-3T3 cells/2.4-cm plate) in 45% Iscove's modified Eagle medium, 45% Dulbecco's minimal essential medium, 10% fetal bovine serum, 100 U/ml penicillin and streptomycin, 4 mM L-glutamine, and 25 μ M 2-mercaptoethanol. For in vitro drug assays, mafosfamide (cyclohexylammonium salt, a CTX analog active in vitro; a generous gift from Asta Medica, Germany) was added at 0, 0.3, 3, and 30 μ g/ml, and viability was measured (see below) 24 hr later.

Assessment of viability, cell-cycle parameters, and apoptosis

Viability of short-term cultured lymphoma cells was analyzed by trypan blue dye exclusion. For analysis of ploidy, apoptosis (as percentage of cells in sub-G₁ peak), and proliferation (as percentage of viable cells in S phase), 10^6 ethanol-fixed lymphoma cells were incubated for 30 min at room temperature in 1 ml of DNA staining solution (200 μ g of propidium iodide and 2 mg of RNase in 10 ml of PBS), and DNA content was measured at 488 nm in a flow cytometer (FACScalibur, Becton Dickinson). In situ proliferation was estimated by counting of mitotic figures (cells in anaphase or telophase) relative to cell number in HE-stained lymphoma sections (four samples each genotype, seven different fields, 200 cells each). In situ apoptosis was visualized in lymphoma sections by HE staining and a fluorescence-based TUNEL assay. TUNEL assays were performed in accordance to the manufacturer's protocol (Boehringer Mannheim). Leukemias were analyzed for apoptotic nuclear morphology by fluorescence microscopy after ethanol fixation and DAPI (4',6-Diamidino-2-phenylindole) staining of peripheral blood samples.

Western blotting analysis

Whole-cell lymphoma cell or normal splenocyte lysates were generated by lysing of extracted cells in SDS sample buffer (60 mM Tris-HCl at pH 6.8, 10% glycerol, 2% SDS, and 5% 2-mercaptoethanol). Samples corresponding to 60 μ g of protein (BioRad Bradford protein assay) were separated on a SDS-polyacrylamide gel and transferred to Immobilon-P membranes (Millipore). p53 was detected using the polyclonal antibody CM5 (Novocastra, 1:2000 dilution), p21 using the polyclonal antibody C-19 (Santa Cruz, 1:500 dilution), and α -tubulin using the monoclonal antibody B-5-1-2 (Sigma, 1:2000 dilution). Protein detection was visualized by ECL (Amersham) or Supersignal (Pierce).

Lymphoma reconstitution and in vivo treatment

Immediately after extraction, 10^6 lymphoma cells in 100 μ l of PBS were reconstituted by tail vein injection into genetically matched, nontransgenic recipient mice (two mice per individual lymphoma sample) to monitor response to treatment. Tumors derived from the *INK4a/ARF*^{-/-} and *p53*^{-/-} backgrounds were reconstituted in C57BL/6 \times 129/sv mice (Jackson Laboratories). CTX was applied as a single 300-mg/kg dose i.p. when arising tumors became well palpable.

Statistical evaluation

Tumor onset data reflect the time between birth and first-time palpability of enlarged lymph nodes; treatment response data reflect the time between remission and relapse as first-time palpability of a recurrent lymph node enlargement. Individual time values were plotted in the Kaplan-Meier population-event-time course format and compared using the log-rank (Mantel-Cox) test. Comparisons of means and standard deviations (s.d.) were performed using the unpaired *t*-test. Ploidy, cell cycle distribution, and sub-G₁ content were analyzed using the ModFit LT 2.0 software.

Acknowledgments

We thank T. Jacks for the *Rb*^{+/-} and *p53*^{-/-} mice; M. Serrano and D. Beach for the *INK4a/ARF*^{-/-} mice; A. Harris for the *E μ -myc* transgenic mice; K. Sokol for histopathology; L. Bianco and the CSHL animal facility for technical assistance; M. Ockler and J. Duffy of the CSHL Graphic Arts facility for help with the artwork; G. Ferbeyre, A. Lin, M. Soengas, and A. Samuelson for editorial advice; and M. Roussel, C. Sherr, and J. Cleveland for discussion of unpublished data. This work was supported by a Dr. Mildred Scheel Cancer Foundation fellowship (C.A.S.), a DOD Breast Cancer Research fellowship (E.d.S.), a Kimmel Scholar Award (S.W.L.), and a grant (CA13106) from the National Cancer Institute (S.W.L.).

The publication costs of this article were defrayed in part by payment of page charges. This article must therefore be hereby marked 'advertisement' in accordance with 18 USC section 1734 solely to indicate this fact.

References

- Adams, J.M. and S. Cory. 1991. Transgenic models for haemopoietic malignancies. *Biochim. Biophys. Acta* **1072**: 9–31.
- Adams, J.M., A.W. Harris, C.A. Pinkert, L.M. Corcoran, W.S. Alexander, S. Cory, R.D. Palmiter, and R.L. Brinster. 1985. The *c-myc* oncogene driven by immunoglobulin enhancers induces lymphoid malignancy in transgenic mice. *Nature* **318**: 533–538.
- Bates, S., A.C. Phillips, P.A. Clark, F. Stott, G. Peters, R.L. Ludwig, and K.H. Vousden. 1998. p14ARF links the tumour suppressors RB and p53. *Nature* **395**: 124–125.
- Beroud, C. and T. Soussi. 1998. *p53* gene mutation: Software and database. *Nucleic Acids Res.* **26**: 200–204.
- Brown, J.M. and B.G. Wouters. 1999. Apoptosis, p53, and tumor cell sensitivity to anticancer agents. *Cancer Res.* **59**: 1391–1399.
- Bunz, F., P.M. Hwang, C. Torrance, T. Waldman, Y. Zhang, L. Dillehay, J. Williams, C. Lengauer, K.W. Kinzler, and B. Vogelstein. 1999. Disruption of p53 in human cancer cells alters the responses to therapeutic agents. *J. Clin. Invest.* **104**: 263–269.

- de Stanchina, E., M.E. McCurrach, F. Zindy, S.Y. Shieh, G. Ferbeyre, A.V. Samuelson, C. Prives, M.F. Roussel, C.J. Sherr, and S.W. Lowe. 1998. E1A signaling to p53 involves the p19^{ARF} tumor suppressor. *Genes & Dev.* **12**: 2434-2442.
- Diccianni, M.B., J. Yu, M. Hsiao, S. Mukherjee, L.E. Shao, and A.L. Yu. 1994. Clinical significance of p53 mutations in relapsed T-cell acute lymphoblastic leukemia. *Blood* **84**: 3105-3112.
- Eischen, C.M., J.D. Weber, M.F. Roussel, C.J. Sherr, and J.L. Cleveland. 1999. Disruption of the ARF-Mdm2-p53 tumor suppressor pathway in Myc-induced lymphomagenesis. *Genes & Dev.* (this issue).
- Elrouby, S., A. Thomas, D. Costin, C.R. Rosenberg, M. Potmesil, R. Silber, and E.W. Newcomb. 1993. p53 gene mutation in B-cell chronic lymphocytic leukemia is associated with drug resistance and is independent of MDR1/MDR3 gene expression. *Blood* **82**: 3452-3459.
- Fan, S.J., W.S. Eldeiry, I. Bac, J. Freeman, D. Jondle, K. Bhatia, A.J. Fornace, I. Magrath, K.W. Kohn, and P.M. Oconnor. 1994. p53 gene mutations are associated with decreased sensitivity of human lymphoma cells to DNA damaging agents. *Cancer Res.* **54**: 5824-5830.
- Giaccia, A.J. and M.B. Kastan. 1998. The complexity of p53 modulation: Emerging patterns from divergent signals. *Genes & Dev.* **12**: 2973-2983.
- Haber, D.A. 1997. Splicing into senescence: The curious case of p16 and p19^{ARF}. *Cell* **91**: 555-558.
- Hermeking, H. and D. Eick. 1994. Mediation of c-myc induced apoptosis by p53. *Science* **265**: 2091-2093.
- Hsu, B., M.C. Marin, A.K. Elnaggar, L.C. Stephens, S. Brisbay, and T.J. McDonnell. 1995. Evidence that c-myc mediated apoptosis does not require wild-type p53 during lymphomagenesis. *Oncogene* **11**: 175-179.
- Jacks, T., A. Fazeli, E.M. Schmitt, R.T. Bronson, M.A. Goodell, and R.A. Weinberg. 1992. Effects of an Rb mutation in the mouse. *Nature* **359**: 295-300.
- Jacks, T., L. Remington, B.O. Williams, E.M. Schmitt, S. Halachmi, R.T. Bronson, and R.A. Weinberg. 1994. Tumor spectrum analysis in p53-mutant mice. *Curr. Biol.* **4**: 1-7.
- Jacobs, J.J.L., B. Scheijen, J.-W. Voncken, K. Kieboom, A. Berns, and M. van Lohuizen. (1999). Bmi-1 collaborates with c-Myc in tumorigenesis by inhibiting c-Myc induced apoptosis via Ink4a/ARF. *Genes & Dev.* (This issue).
- Jawhs, J.J.L., B. Scheijen, J.-W. Voncken, K. Kieboom, A. Berns, M. van Lohuizen. 1999. Bmi-1 collaborates with cMyc in tumorigenesis by inhibiting c-Myc induced apoptosis via INK4a/ARF. *Genes & Dev.* (This issue).
- Kamijo, T., F. Zindy, M.F. Roussel, D.E. Quelle, J.R. Downing, R.A. Ashmun, G. Grosfeld, and C.J. Sherr. 1997. Tumor suppression at the mouse INK4a locus mediated by the alternative reading frame product p19^{ARF}. *Cell* **91**: 649-659.
- Kamijo, T., J.D. Weber, G. Zambetti, F. Zindy, M.F. Roussel, and C.J. Sherr. 1998. Functional and physical interactions of the ARF tumor suppressor with p53 and Mdm2. *Proc. Natl. Acad. Sci.* **95**: 8292-8297.
- Kastan, M.B., O. Onyekwere, D. Sidransky, B. Vogelstein, and R.W. Craig. 1991. Participation of p53 protein in the cellular response to DNA damage. *Cancer Res.* **51**: 6304-6311.
- Kastan, M.B., Q. Zhan, W.S. el-Deiry, F. Carrier, T. Jacks, W.V. Walsh, B.S. Plunkett, B. Vogelstein, and A. Fornace Jr. 1992. A mammalian cell cycle checkpoint pathway utilizing p53 and GADD45 is defective in ataxia-telangiectasia. *Cell* **71**: 587-597.
- Lowe, S.W. and H.E. Ruley. 1993. Stabilization of the p53 tumor suppressor is induced by adenovirus E1A and accompanies apoptosis. *Genes & Dev.* **7**: 535-545.
- Lowe, S.W., H.E. Ruley, T. Jacks, and D.E. Housman. 1993. p53-dependent apoptosis modulates the cytotoxicity of anti-cancer agents. *Cell* **74**: 954-967.
- Palmero, I., C. Pantoja, and M. Serrano. 1998. p19^{ARF} links the tumour suppressor p53 to Ras. *Nature* **395**: 125-126.
- Pomerantz, J., N. Schreiber-Agus, N.J. Liegeois, A. Silverman, L. Alland, L. Chin, J. Potes, K. Chen, I. Orlow, H.W. Lee, C. Cordon-Cardo, and R.A. DePinho. 1998. The INK4a tumor suppressor gene product, p19^{ARF}, interacts with MDM2 and neutralizes MDM2's inhibition of p53. *Cell* **92**: 713-723.
- Prives, C. 1998. Signaling to p53: Breaking the MDM2-p53 circuit. *Cell* **95**: 5-8.
- Ruas, M. and G. Peters. 1998. The p16INK4a/CDKN2A tumor suppressor and its relatives. *Biochim. Biophys. Acta* **1378**: F115-F177.
- Schmitt, C.A. and S.W. Lowe. 1999. Apoptosis and therapy. *J. Pathol.* **187**: 127-137.
- Serrano, M., G.J. Hannon, and D. Beach. 1993. A new regulatory motif in cell-cycle control causing specific inhibition of cyclin D/CDK4. *Nature* **366**: 704-707.
- Serrano, M., H. Lee, L. Chin, C. Cordon-Cardo, D. Beach, and R.A. DePinho. 1996. Role of the INK4a locus in tumor suppression and cell mortality. *Cell* **85**: 27-37.
- Serrano, M., A.W. Lin, M.E. McCurrach, D. Beach, and S.W. Lowe. 1997. Oncogenic ras provokes premature cell senescence associated with accumulation of p53 and p16^{INK4a}. *Cell* **88**: 593-602.
- Sherr, C.J. 1998. Tumor surveillance via the ARF-p53 pathway. *Genes & Dev.* **12**: 2984-2991.
- Stott, F.J., S. Bates, M.C. James, B.B. McConnell, M. Starborg, S. Brookes, I. Palmero, K. Ryan, E. Hara, K.H. Vousden, and G. Peters. 1998. The alternative product from the human CDKN2A locus, p14^{ARF}, participates in a regulatory feedback loop with p53 and MDM2. *EMBO J.* **17**: 5001-5014.
- Strasser, A., A.W. Harris, M.L. Bath, and S. Cory. 1990. Novel primitive lymphoid tumours induced in transgenic mice by cooperation between myc and bcl-2. *Nature* **348**: 331-333.
- Tao, W. and A.J. Levine. 1999. P19^{ARF} stabilizes p53 by blocking nucleo-cytoplasmic shuttling of Mdm2. *Proc. Natl. Acad. Sci.* **96**: 6937-6941.
- Wallace-Brodeur, R.R. and S.W. Lowe. 1999. Clinical implications of p53 mutations. *Cell Mol. Life Sci.* **55**: 64-75.
- Wattél, E., C. Preudhomme, B. Hecquet, M. Vanrumbeke, B. Quesnel, I. Dervite, P. Morel, and P. Fenaux. 1994. p53 mutations are associated with resistance to chemotherapy and short survival in hematologic malignancies. *Blood* **84**: 3148-3157.
- Weber, J.D., L.J. Taylor, M.F. Roussel, C.J. Sherr, and D. Bartsch. 1999. Nucleolar Arf sequesters Mdm2 and activates p53. *Nature Cell Biol.* **1**: 20-26.
- Wilson, W.H., J. Teruya-Feldstein, T. Fest, C. Harris, S.M. Steinberg, E.S. Jaffe, and M. Raffeld. 1997. Relationship of p53, bcl-2, and tumor proliferation to clinical drug resistance in non-Hodgkin's lymphomas. *Blood* **89**: 601-609.
- Zhang, Y. and Y. Xiong. 1999. Mutations in human ARF exon 2 disrupt its nucleolar localization and impair its ability to block nuclear export of MDM2 and p53. *Mol. Cell* **3**: 579-591.
- Zhang, Y., Y. Xiong, and W.G. Yarbrough. 1998. ARF promotes MDM2 degradation and stabilizes p53: ARF-INK4a locus deletion impairs both the Rb and p53 tumor suppression pathways. *Cell* **92**: 725-734.
- Zindy, F., C.M. Eischen, D.H. Randle, T. Kamijo, J.L. Cleveland, C.J. Sherr, and M.F. Roussel. 1998. Myc signaling via the ARF tumor suppressor regulates p53-dependent apoptosis and immortalization. *Genes & Dev.* **12**: 2424-2433.



DEPARTMENT OF THE ARMY
US ARMY MEDICAL RESEARCH AND MATERIEL COMMAND
504 SCOTT STREET
FORT DETRICK, MARYLAND 21702-5012

REPLY TO
ATTENTION OF:

MCMR-RMI-S (70-1y)

21 Feb 03

MEMORANDUM FOR Administrator, Defense Technical Information
Center (DTIC-OCA), 8725 John J. Kingman Road, Fort Belvoir,
VA 22060-6218

SUBJECT: Request Change in Distribution Statement

1. The U.S. Army Medical Research and Materiel Command has reexamined the need for the limitation assigned to technical reports written for this Command. Request the limited distribution statement for the enclosed accession numbers be changed to "Approved for public release; distribution unlimited." These reports should be released to the National Technical Information Service.

2. Point of contact for this request is Ms. Kristin Morrow at DSN 343-7327 or by e-mail at Kristin.Morrow@det.amedd.army.mil.

FOR THE COMMANDER:

A handwritten signature in black ink, appearing to read "Phyllis M. Rinehart", is written over the typed name and title.

PHYLLIS M. RINEHART
Deputy Chief of Staff for
Information Management

Encl

ADB263458	ADB282838
ADB282174	ADB233092
ADB270704	ADB263929
ADB282196	ADB282182
ADB264903	ADB257136
ADB268484	ADB282227
ADB282253	ADB282177
ADB282115	ADB263548
ADB263413	ADB246535
ADB269109	ADB282826
ADB282106	ADB282127
ADB262514	ADB271165
ADB282264	ADB282112
ADB256789	ADB255775
ADB251569	ADB265599
ADB258878	ADB282098
ADB282275	ADB232738
ADB270822	ADB243196
ADB282207	ADB257445
ADB257105	ADB267547
ADB281673	ADB277556
ADB254429	ADB239320
ADB282110	ADB253648
ADB262549	ADB282171
ADB268358	ADB233883
ADB257359	ADB257696
ADB265810	ADB232089
ADB282111	ADB240398
ADB273020	ADB261087
ADB282185	ADB249593
ADB266340	ADB264542
ADB262490	ADB282216
ADB266385	ADB261617
ADB282181	ADB269116
ADB262451	
ADB266306	
ADB260298	
ADB269253	
ADB282119	
ADB261755	
ADB257398	
ADB267683	
ADB282231	
ADB234475	
ADB247704	
ADB258112	
ADB267627	