

UNCLASSIFIED

AD NUMBER
ADB263737
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies only; Proprietary Info.; Oct 2000. Other requests shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott St., Fort Detrick, MD 21702-5012.
AUTHORITY
USAMRMC ltr, 23 Aug 2001

THIS PAGE IS UNCLASSIFIED

AD _____

Award Number: DAMD17-95-1-5070

TITLE: Pregnancy-Induced Changes in Body Fat, Physical Fitness
and Energy Requirements in Military and Civilian Women

PRINCIPAL INVESTIGATOR: Nancy F. Butte, Ph.D.

CONTRACTING ORGANIZATION: Baylor College of Medicine
Houston, Texas 77030-3498

REPORT DATE: October 2000

TYPE OF REPORT: Final

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Distribution authorized to U.S. Government agencies only (proprietary information, Oct 00). Other requests for this document shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott Street, Fort Detrick, Maryland 21702-5012.

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

20010301 089

NOTICE

USING GOVERNMENT DRAWINGS, SPECIFICATIONS, OR OTHER DATA INCLUDED IN THIS DOCUMENT FOR ANY PURPOSE OTHER THAN GOVERNMENT PROCUREMENT DOES NOT IN ANY WAY OBLIGATE THE U.S. GOVERNMENT. THE FACT THAT THE GOVERNMENT FORMULATED OR SUPPLIED THE DRAWINGS, SPECIFICATIONS, OR OTHER DATA DOES NOT LICENSE THE HOLDER OR ANY OTHER PERSON OR CORPORATION; OR CONVEY ANY RIGHTS OR PERMISSION TO MANUFACTURE, USE, OR SELL ANY PATENTED INVENTION THAT MAY RELATE TO THEM.

LIMITED RIGHTS LEGEND

Award Number: DAMD17-95-1-5070

Organization: Baylor College of Medicine

Location of Limited Rights Data (Pages):

Those portions of the technical data contained in this report marked as limited rights data shall not, without the written permission of the above contractor, be (a) released or disclosed outside the government, (b) used by the Government for manufacture or, in the case of computer software documentation, for preparing the same or similar computer software, or (c) used by a party other than the Government, except that the Government may release or disclose technical data to persons outside the Government, or permit the use of technical data by such persons, if (i) such release, disclosure, or use is necessary for emergency repair or overhaul or (ii) is a release or disclosure of technical data (other than detailed manufacturing or process data) to, or use of such data by, a foreign government that is in the interest of the Government and is required for evaluational or informational purposes, provided in either case that such release, disclosure or use is made subject to a prohibition that the person to whom the data is released or disclosed may not further use, release or disclose such data, and the contractor or subcontractor or subcontractor asserting the restriction is notified of such release, disclosure or use. This legend, together with the indications of the portions of this data which are subject to such limitations, shall be included on any reproduction hereof which includes any part of the portions subject to such limitations.

THIS TECHNICAL REPORT HAS BEEN REVIEWED AND IS APPROVED FOR PUBLICATION.

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE October 2000	3. REPORT TYPE AND DATES COVERED Final (25 Sep 95 - 24 Sep 00)
----------------------------------	--------------------------------	---

4. TITLE AND SUBTITLE Pregnancy-Induced Changes in Body Fat, Physical Fitness and Energy Requirements in Military and Civilian Women	5. FUNDING NUMBERS DAMD17-95-1-5070
---	--

6. AUTHOR(S) Nancy F. Butte, Ph.D.

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Baylor College of Medicine Houston, Texas 77030-3498 E-MAIL: nbutte@bcm.tmc.edu	8. PERFORMING ORGANIZATION REPORT NUMBER
--	--

9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012	10. SPONSORING / MONITORING AGENCY REPORT NUMBER
---	--

11. SUPPLEMENTARY NOTES

12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; Distribution unlimited	12b. DISTRIBUTION CODE
---	------------------------

13. ABSTRACT (Maximum 200 Words) This study addresses the impact of pregnancy-induced changes in body composition and physical fitness on postpartum return to duty readiness. Physical activity, weight, FFM and fat mass, fitness, strength, and iron status was measured in women with low to high pre-pregnancy BMI through a complete reproductive cycle. Physical activity was quantified by the doubly-labeled water method. Body composition was measured by hydrodensitometry, total body water, ⁴⁰ K counting, prompt-gamma activation, and DXA. Data obtained prior to pregnancy strongly indicate a need to revise military equations used to predict body fat. Postpartum weight retention at 6 mo averaged 2.7 ± 3.9 kg, and was greater in the high BMI group (4.0 ± 4.4 kg) than the low and normal BMI groups (2.4 ± 3.5 kg). BMI had increased by approximately one unit in all groups. FM increased by 2.1 ± 4.1 kg FM at 6 mo postpartum, with means of 2.6 ± 3.7, 1.7 ± 4.1 and 3.0 ± 5.0 kg in the low, normal and high BMI groups, respectively. There was evidence of deconditioning at 6 wk postpartum with near recovery at 6 mo postpartum. The effect of gestational weight gain, fat accretion and deconditioning would affect a woman's ability to return to weight, body fat and fitness standards in the 6 mo following delivery.

14. SUBJECT TERMS Women's Health	15. NUMBER OF PAGES 42
	16. PRICE CODE

17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited
---	--	---	---

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

_____ Where copyrighted material is quoted, permission has been obtained to use such material.

_____ Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

_____ Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

N/A In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Resources, National Research Council (NIH Publication no. 86-23, Revised 1985).

X NFB For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal law 45 CFR 46.

N/A In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the national Institutes of health.

N/A In conduct of research utilizing recombinant DNA technology, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

N/A In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

Janay Buttz
PI - Signature

10/26/00
Date

TABLE OF CONTENTS

1. FRONT COVER	1
2. SF 298, REPORT DOCUMENTATION	2
3. FOREWORD	3
4. TABLE OF CONTENTS	4
List of Figures	6
List of Tables	6
5. INTRODUCTION	7
A. Subject, Hypothesis and Technical Objectives of the Research	7
B. Background of Previous Work	8
6. METHODS	13
<i>Study Design</i>	13
<i>Subjects</i>	14
<i>Methodology</i>	14
<i>Anthropometry</i>	14
<i>Hydrodensitometry</i>	15
<i>Total Body Potassium (TBK)</i>	15
<i>Dual Energy X-ray Absorptiometry (DEXA)</i>	15
<i>Total Body Nitrogen (TBN)</i>	15
<i>Muscle and Nonmuscle Mass</i>	16
<i>Body Composition Models</i>	16
<i>Total Body Water (TBW)</i>	16
<i>Total Energy Expenditure</i>	16
<i>Room Respiration Calorimetry</i>	16
<i>Physical Activity</i>	17
<i>Physical Fitness</i>	17
<i>Strength</i>	17
<i>Infant Outcome</i>	17
<i>Biochemical Analysis</i>	17
<i>Statistical Analysis</i>	17

7. RESULTS AND DISCUSSION	18
Data Presentation	18
Subject Description	18
Reproductive History	18
Pregnancy Outcome	19
Iron Status	19
Anthropometry	21
Body Composition	23
Figures 1-6	27
Fat-free Mass Compartment	29
Energy Expenditure and Substrate Utilization by Room Respiration Calorimetry ..	30
Total Energy Expenditure by Doubly-labeled Water Method	32
Fitness and Strength	33
8. CONCLUSIONS	36
9. REFERENCES	36
10. BIBLIOGRAPHY OF PUBLICATIONS	41
11. LIST OF PERSONNEL	41

LIST OF FIGURES

Figure 1,4. Prediction of fat mass percent from Army equation vs. Fuller-4 component model (prepregnancy)	27
Figure 2,5. Prediction of fat mass percent from Air Force equation vs. Fuller-4 component model (prepregnancy)	27
Figure 3,6. Prediction of fat mass percent from Navy equation vs. Fuller-4 component model (prepregnancy)	27
Figure 1,4. Prediction of fat mass percent from Army equation vs. Fuller-4 component model (postpartum)	28
Figure 2,5. Prediction of fat mass percent from Air Force equation vs. Fuller-4 component model (postpartum)	28
Figure 3,6. Prediction of fat mass percent from Navy equation vs. Fuller-4 component model (postpartum)	28

LIST OF TABLES

Table 1. Body fat standards for women as a percent of body weight	8
Table 2. Recommended ranges of gestational weight gain by pre-pregnancy BMI	9
Table 3. Expected gestational weight gain and postpartum weight retention in military women	13
Table 4. Study design	14
Table 5. Fat mass prediction equations for the U.S. Army, Navy, and Air Force service women	15
Table 6. Reproductive history	18
Table 7. Pregnancy outcome	19
Table 8. Iron status	20
Table 9. Anthropometry	21
Table 10. Fat mass (FM) estimated by deuterium dilution, hydrodensitometry, total body potassium, DEXA, and skinfold thicknesses, and multi-component models	23
Table 11. Body fat mass (FM) predicted from military equations and body composition models at the pre-pregnancy and 24 wk postpartum	26
Table 12. Fat-free mass (FFM) compartment determined by total body potassium, DEXA, and prompt-gamma activation	29
Table 13. Room respiration calorimetry	30
Table 14. Substrate utilization estimated from 24-h respiration calorimetry	32
Table 15. Total energy expenditure (TEE) by doubly-labeled water method	33
Table 16. Physical fitness and strength	34

5. INTRODUCTION

A. Subject, Hypothesis and Technical Objectives of the Research

The level of fatness associated with optimal physical fitness in women is less than the level of fatness associated with optimal fetal growth and survival. Women with low pre-pregnancy energy stores are at greater risk for fetal loss, premature delivery and intrauterine growth retardation. However, a low pre-pregnancy body mass index (BMI) can be compensated for by a greater pregnancy weight gain. In this study the effect of physical activity level on changes in physical fitness, weight, FFM, fat mass, energy expenditure will be evaluated through a complete reproductive cycle in sixty-eight women stratified by pre-pregnancy BMI. Measurements will be done prior to pregnancy to establish pre-pregnancy nutritional status and physical fitness. During pregnancy measurements will be made at 8 wk (first trimester), 22 wk (second trimester), and 36 wk (third trimester). Postpartum measurements will be made at 2 wk to establish postpartum baseline body composition; at 6 wk when military and many civilian women return to work; and at 24 wk when military women must meet weight and physical fitness standards. We will investigate if military policies requiring women to return to weight and physical fitness standards by 6 mo postpartum are physiologically reasonable. The appropriateness of military body weight retention standards and the accuracy of the military equations used to predict body fat will be assessed in postpartum women.

Hypothesis

Moderate levels of *physical activity* will maintain *physical fitness* and limit *excess fat deposition* during pregnancy without jeopardizing fetal growth if dietary intake is not restricted, and facilitate *fat mobilization* and conserve *fat-free mass* during postpartum weight loss.

Technical Objectives

1. *The effect of physical activity level on pregnancy-induced and postpartum changes in weight, FFM and fat mass will be compared in military and civilian women with low to high pre-pregnancy BMI. Changes in FFM and fat mass will be computed from measurements of body volume, total body water, potassium, nitrogen and bone mineral through a complete reproductive cycle (pre-pregnancy-6 mo postpartum).*
2. *The effect of physical activity level on pregnancy-induced and postpartum changes in physical fitness will be compared in military and civilian women. Submaximal and maximal aerobic capacity will be measured through a complete reproductive cycle. The impact of body composition (specifically FFM, muscle mass and body fat), and iron status on physical performance will be assessed.*
3. *The effect of physical activity level on the energy requirements of physically-active military and civilian adult women will be determined prior to pregnancy, during pregnancy and postpartum. Energy requirements will be estimated from rates of energy expenditure and energy deposition/mobilization.*

B. Background of Previous Work

Military weight and body fat standards

The rationale for weight and body fat standards for accession and retention in the military as stated in Army regulation 600-9 (1) is "to insure that all personnel are able to meet the physical demands of their duties under combat conditions and present a trim military appearance at all times." Although similar, weight and body fat standards differ somewhat among the military services. If an individual exceeds the acceptable weight for height standard, body fat is assessed using anthropometric measures. The Army uses height, weight, and circumferences of neck, forearm, wrist, and hips in women to calculate body fat (2). The Navy uses height, weight and circumferences of neck, waist and hips in women (3). The Marines use height, weight, flexed biceps, forearm, neck, waist and thigh circumferences for women (4). The Air Force uses height, weight and forearm circumference (this equation is no longer used) (5).

The weight and body fat standards for accession and retention differ for men and women. Between 1960 and 1983 the maximum weight limits for women were lowered by 15-20 pds (AR 40-501)(6). The tables for women are considerably more restrictive relative to the national population. The military standard levels for body fatness are lower in men than women, acknowledging the fact that women biologically have higher body fat.

Table 1. Body fat standards for women as a percent of body weight (7)

Age	<u>17-20y</u>	<u>21-27y</u>	<u>28-39y</u>	<u>>40y</u>
Army	30%	32%	34%	36%
Navy	-----36%-----			
Air Force	-----28%-----		-----34%-----	

However, accession and retention weight criteria are stricter for women than men. For enrollment into the Army men may be 37% above the desirable weight based on the 1959 Metropolitan Life Insurance Tables, while women can be only 6% above. For retention in the Army men can be 14% above desirable weight, and women only 5% above. Consequently, 29% of women Army recruits are rejected in contrast to 3% of male recruits (8). Military weight standards for women are set at an upper BMI limit of 24. Therefore, military women enter pregnancy with a low to normal BMI.

Performance standards

For retention in the military, personnel are evaluated regularly not only for weight compliance, but also for aerobic fitness (7). Physical fitness tests for women consist of a 1.5-2-mile run, push-ups and sit-ups or curl-ups. Although the tasks of military women are increasingly diverse, the military contends that all individuals need to maintain a certain level of physical fitness to preserve the combat readiness of the services. If an individual fails to meet the body fat or physical fitness standards, he/she is assigned a program of diet and exercise. Individuals who do not lose sufficient weight or body fat are discharged from the military.

Pregnancy-induced changes in body weight

In the Institute of Medicine (9) appointed a Subcommittee to review the effect of gestational weight gain on maternal and child health and to make recommendations for optimal weight gain during pregnancy. Gestational weight gains associated with optimal infant outcome were found to be a function of maternal pre-pregnancy BMI. Several epidemiologic studies demonstrated that the effect of a given weight gain is greatest in thin women and least in overweight women. Pre-pregnancy BMI is also a determinant of fetal growth above and beyond its effect on gestational weight gain. Women with low pre-pregnancy BMIs tend to have smaller infants than those of their heavier counterparts. Low gestational weight gain is associated with a higher risk of intrauterine growth retardation and subsequent poor somatic and neurobehavioral development. Increased perinatal mortality among infants born to women with low weight gain especially those with low pre-pregnancy weight for height was seen in the Collaborative Perinatal Project (10) and the 1980 National Fetal Mortality Survey (11).

The Subcommittee concluded that desirable gestational weight gains should be based on pre-pregnancy BMI and should include the mean weight gain for women delivering full-term infants with birth weights in the optimal range of 3 to 4 kg (9).

Table 2. Recommended ranges of gestational weight gain by pre-pregnancy BMI (9)

<u>Weight for height category</u>	<u>Recommended total gain (kg)</u>
Low (BMI<19.8)	12.5-18.0
Normal (BMI=19.8-26.0)	11.5-16.0
High (BMI>26.0-29.0)	7.0-11.5

Although gestational weight gain is correlated with birth weight, which component(s) of gestational weight gain is(are) critical for optimal fetal growth is unknown. Fetal growth may be influenced more by specific changes in fat-free mass (FFM), fat or water than by total gestational weight gain. In a study of 56 Swedish women total weight gain (14.8 kg), but not fat mass accretion (4-5 kg), was positively correlated with birth weight (12). However, their estimates of fat mass based on skinfolds were relatively imprecise. In another study of 115 Scottish women the lack of correlation between maternal fat accretion, estimated from weight retention 2-3 wk postpartum, and birth weight was confirmed (13). These results should be interpreted cautiously, since fat accretion was not measured directly and only women with normal pre-pregnancy BMI (22.5) were studied. In women with low pre-pregnancy BMI fat accretion may be more critical to fetal growth. There may be a threshold for maternal fat gain below which fetal growth is compromised. Further studies on the composition of weight gain in women stratified by BMI and its impact on pregnancy outcome are needed.

Pregnancy-induced changes in body composition

The amount of FFM and fat accreted at any given gestational weight gain is poorly defined. Theoretical estimations of protein and fat deposition in the maternal (uterus, breasts, blood, adipose tissue, extracellular fluid) and fetal (fetus, placenta, amniotic fluid) compartments have been

calculated for a 12.5 kg weight gain (14). Total body water increases continuously through pregnancy, primarily in extracellular fluids; therefore, the hydration of FFM increases substantially through pregnancy (15). Protein accretion rates estimated by recent nitrogen balance studies agree with Hytten's theoretical values (16). Therefore, it seems unlikely that protein is stored in excess of the amounts accounted for by the fetus and maternal reproductive tissues. The suggestion that protein is stored in muscle early in pregnancy, based on urinary 3-methylhistidine excretion data (17), has been refuted by others who attributed the higher urinary excretion of 3-methylhistidine to decreased renal tubular reabsorption and higher skeletal muscle turnover (16). Our measurements of total body nitrogen and potassium retention in pregnancy will resolve the amount and partitioning of protein deposited.

During pregnancy fat is deposited primarily in the subcutaneous adipose of the lower and upper trunk, and thighs (18-20), however, the amount of fat deposited is uncertain. Because of the increased hydration of FFM during pregnancy, standard techniques used to estimate fat deposition are invalid during pregnancy. Corrections were not applied in earlier dilution studies (15,21), but were in a later study (22), in which the mean fat gain was 2.77 ± 3.23 kg. Based on underwater weighing corrected for changes in the density of FFM, fat mass accretion was 2.7 ± 2.2 kg (23). Based on combined measurements of TBW and TBK, mean fat gains were 1.87 ± 2.23 kg (24) and 5.8 ± 4.0 kg (25). Recent values of fat gain differ from Hytten's original estimate of 3.3 kg fat. Technical errors undoubtedly contribute to the high variability in fat gain, but gestational weight gain, which is positively correlated with fat gain, is also variable in healthy women (26). Because of wide variation, changes in body composition can only be estimated from serial measurements, preferably with pre-pregnancy baseline values.

Postpartum weight and body fat loss

Postpartum weight loss is influenced by total gestational weight gain, age, parity, pre-pregnancy BMI, and feeding mode (27). The 1988 National Maternal and Infant Survey indicated a strong positive association between gestational weight gain and postpartum weight retention 10 to 18 months postpartum (28). Excessive weight retention was more common in black women in every weight gain category. Greater weight losses were observed in women with lower pre-pregnancy weights (20). Lactation facilitates weight loss in most women. Lactating women averaged 0.7 kg/mo weight loss during the first 4-6 mo postpartum (29,30). This weight loss was compatible with successful lactation and was associated with a decrease in body fat from 28.0 to 26.3%. Acceleration of weight loss may compromise lactation performance. Short-term energy restriction (1591 kcal/d for 1 wk resulting in a weight loss of 1.18 kg/wk) was associated with a fall in subsequent milk output and infant weight gain (31).

Accelerated weight loss may result in an undesirable loss of muscle mass in postpartum women (32). Whenever body weight is reduced, both FFM and fat mass contribute to the weight loss. For any given change in weight, the ratio $\Delta\text{FFM}/\Delta\text{WT}$ is inversely related to initial body fat. However, exercise-training has been shown to enhance FFM preservation during diet restriction. Moderate exercise training reduced the amount of body weight lost as FFM compared to dietary restriction alone (33). Exercise training also can induce a greater energy deficit and mobilization of fat. We

therefore hypothesize that dietary restriction combined with moderate levels of physical activity may preserve FFM in postpartum women.

When pregnant women perform nonweight-bearing exercise such as stationary cycling, oxygen consumption (VO_2 ($\text{l} \cdot \text{min}^{-1}$)) is either unchanged or only slightly increased at any given submaximal work rate compared to nonpregnant women (34). During submaximal weight-bearing exercise (eg. walking, running, treadmill exercise) maternal VO_2 ($\text{l} \cdot \text{min}^{-1}$) is significantly increased approximately in proportion to maternal weight gain. VO_2 expressed in $\text{ml} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ is similar to or slightly reduced during pregnancy compared to nonpregnant state at the same speed and grade of walking. Limited data indicate that maximal VO_2 ($\text{l} \cdot \text{min}^{-1}$) during cycle ergometry and treadmill exercise is not altered by pregnancy compared to the postpartum period (35,36). However, in the only study in which aerobic capacity was assessed prior to pregnancy, $\text{VO}_{2\text{max}}$ ($\text{ml} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$) during cycle ergometry was higher pre-pregnancy compared to 4 to 8 wk postpartum (37). The decrement in $\text{VO}_{2\text{max}}$ was attributed to increased body weight and decreased physical activity through pregnancy and the postpartum period. Whether this deconditioning effect is an inevitable consequence of pregnancy or whether moderate exercise throughout pregnancy can ameliorate the decline in aerobic capacity is uncertain. The effect of pregnancy-induced changes in body composition on physical fitness has not been evaluated through a reproductive cycle.

Moderate exercise under the ACOG Guidelines (38) poses minimal risk to the mother and her fetus. Such exercise programs may maintain aerobic fitness and control gestational weight gain (34). Strenuous exercise, on the other hand, may result in inadequate weight gain and give rise to smaller (300-500 g less) infants.

Effect of iron status on work capacity and performance

Maximal aerobic capacity is linearly related to hemoglobin concentration in humans (39). In iron deficiency without anemia skeletal muscle function is impaired by a decrease in mitochondrial iron-dependent enzymes of the electron transport chain and cytochromes. Iron deficiency induces a greater dependence on anaerobic glucose utilization, with lactic acidemia as a consequence. These defects reduce endurance and submaximal work performance. This is of practical importance since most human work is performed at submaximal levels (40% of $\text{VO}_{2\text{max}}$). In iron-deficient women without overt anemia total O_2 uptake and total energy expenditure were decreased, and post-exercise lactate concentration was increased in response to a progressive, graded aerobic capacity test on a cycle ergometer (40). Peak oxygen consumption was not impaired.

Postpartum women are "at risk" for iron deficiency or anemia. Iron losses during pregnancy include loss to the fetus and placenta, blood loss at delivery, and basal losses totalling approx. 840 mg (9). Blood loss during a cesarean delivery is almost twice that of a vaginal delivery. While we do not expect to find overt anemia in pregnant military women under medical surveillance, marginal iron deficiency may be present.

Pregnancy-induced changes in energy requirements

The energy requirement of pregnancy is a topic of considerable uncertainty as reflected in the lack of consensus in international recommendations: +1.20 MJ/d for all trimesters (41); +1.25 MJ/d for the last 2 trimesters (42); +0.80 MJ/d for the third trimester (43).

The energy requirement of pregnancy entails the energy deposited in maternal and fetal tissues and their associated increase in energy expenditure. Basal metabolic rate (BMR) steadily rises through pregnancy, due primarily to the products of conception and to a lesser extent to increased maternal cardiac and respiratory work. By late pregnancy VO_2 is 16-32% above nonpregnant values. Serial measurements of BMR through pregnancy indicate considerable interindividual variation in metabolic response (22,25,44,45), which was correlated to pre-pregnancy fatness, weight gain and fat gain (22). BMR actually declined in the first trimester in some women, suggesting increased metabolic efficiency. The lean women tended to be energy-sparing and the fatter women energy-profligate.

Total energy expenditure (TEE) using room calorimetry (46,47) and the doubly-labeled water method (22,48) has been measured longitudinally in only a few pregnant women. Near term TEE increased 1.5-2 MJ/d. Although changes in BMR accounted for most the increment in TEE, the level of physical activity contributed significantly to the variability in TEE.

The energy requirement of pregnancy will depend on the woman's gestational weight gain and the level of physical activity maintained throughout pregnancy. If a moderate level of physical activity and therefore physical fitness are maintained, it is of paramount importance that dietary intake is sufficient to meet maternal and fetal needs. In women with low pre-pregnancy BMI, if dietary intake is not sufficient to replenish maternal stores, fetal growth will be suboptimal. However, recommendations for dietary intake cannot be made, since the energy requirements of physically active pregnant women with low to normal pre-pregnancy BMI have not been quantified.

Nutritional implications for military women during reproduction

The nutritional problems of military women are similar to physically active civilian women participating in recreational sports. Inadequate intakes of iron and calcium place these weight-conscious women at risk for anemia and osteoporosis (49). Friedl (50) reported that 36.6% of West Point cadets were at risk of developing iron deficiency. Inadequate iron nutrition may impact physical performance through its effect on O_2 transport and oxidative metabolism (39). The nutritional status of military women may be further jeopardized by the need to meet military body weight and fat standards.

Because pregnant military women are required to meet weight and physical fitness standards by 6 mo postpartum, they may restrain food intake and possibly jeopardize fetal growth. Postpartum military women may seriously restrict food intake to accelerate weight loss resulting in loss of fat and FFM, and possibly compromising milk production, if breast-feeding. Military women enter pregnancy with low to normal BMIs, since the military retention weight standards are equivalent to a BMI of 24. We calculated desirable gestational weight gains for optimal infant outcome and postpartum weight loss for women with low and normal pre-pregnancy BMIs (Table 3) based on the IOM recommendations for weight gain; an immediate weight loss associated with the baby, placenta and amniotic fluid of 4.85 kg (14); diuresis resulting in an additional 3.0 kg loss by day 15 postpartum; further weight loss at an average rate of 0.7 kg/mo.

Table 3. Expected gestational weight gain and postpartum weight retention in military women

	<u>Low BMI</u>	<u>Normal BMI</u>
Weight gain (kg)	12.5-18.0	11.5-16.0
Weight retention day 1 postpartum (kg)	7.6-13.1	6.6-11.2
Weight retention day 15 postpartum (kg)	4.6-10.1	3.6-8.2
Weight retention day 42 postpartum (kg)	3.6-9.1	2.7-7.2
Weight retention day 180 postpartum (kg)	0.4-5.9	0.6-4.0

If military women gain at the recommended levels and lose at a reasonable rate postpartum, they will return to active duty at day 42 postpartum with significant excess weight. At 6 mo when military women are expected to meet weight standards, many women will be 0.5-6.0 kg above their pre-pregnancy weight. In order to achieve the weight standards by 6 mo, military women must restrict dietary intake to accelerate weight loss.

The level of physical activity maintained by pregnant military women is uncertain. Although military "pregnancy profiles" exempt women from jobs requiring heavy physical work and eliminates physical fitness testing, these physically-fit women may voluntarily continue to exercise. Or they may become relatively inactive and physical fitness may decline due to deconditioning. Upon returning to work at 6 wk postpartum, military women are expected to perform at their pre-pregnancy work capacity. Whether pre-pregnancy physical fitness has been regained by 6 wk postpartum is uncertain. In only one study was aerobic capacity assessed prior to pregnancy and 4-8 wk postpartum; a decline in aerobic capacity was observed postpartum and attributed to weight gain and decreased physical activity during pregnancy (37). Whether this detraining effect is inevitable during pregnancy or whether exercising throughout pregnancy ameliorates the decline is unknown.

Application of the body fat equations to postpartum women is questionable. Although the prediction equations for body fat have been cross-validated for men and women, they have not been evaluated for postpartum women (3). The Army and Navy equations may overestimate total body fat since these equations use hip circumference, a site of predominate fat deposition in pregnant women. The Air Force equation may underestimate total body fat since it uses forearm circumference which changes little during pregnancy.

Information relevant to the Military Recommended Dietary Allowances (51) for energy intakes of active duty and reserve military women will be provided by this study. Energy requirements of physically active women will be defined prior to, during and after pregnancy which may assist in the design of feeding strategies and food rations for military women.

6. METHODS

Study Design

The effect of physical activity level on changes in physical fitness, weight, FFM, fat mass, and energy expenditure will be evaluated through a complete reproductive cycle in sixty-eight women. Women will be studied prior to conception, at each trimester, and at 2, 6 and 24 wk postpartum.

Table 4. Study design

Study variables	Pregnancy				Postpartum		
	0	8wk	22wk	36wk	2wk	6wk	24wk
Anthropometry	x	x	x	x	x	x	x
Hydrodensitometry	x	x	x	x	x		x
TBK	x	x	x	x	x		x
DEXA	x				x	x	x
TBN	x				x		x
TBW		x			x		
TEE/TBW	x		x	x			x
24-h calorimetry	x	x	x	x			x
Physical fitness	x	x	x	x		x	x
Iron status	x	x	x	x		x	x

Subjects

Sixty-eight women will be enrolled, stratified by pre-pregnancy BMI. An upper BMI limit of 24.0 was designated in accordance with military weight standards. However, we are enrolling an additional 34 women with BMI>24, to describe women in the military in noncompliance of weight standards. Subjects will be healthy, physically-active, nonsmoking, ages 18-39 years, parity not greater than 4, no chronic medications or alcohol/drug abuse. Health history should be unremarkable (i.e., normotensive, glucose tolerant, nonanemic and euthyroid).

Methodology

Anthropometry. Maternal body weight to the nearest 0.1 kg will be measured with a digital balance (Scale-Tronix, Dallas TX). Height to the nearest 1 mm will be measured with a stadiometer (Holtain, Ltd, Crymmych, Pembs, UK). The circumferences of the head, chest, upper arm, forearm, wrist, neck, abdomen, thigh, and calf will be measured to the nearest 1 mm in duplicate with a metal tape. These sites include those currently used in military equations to predict body fat. Skinfold thicknesses will be measured to the nearest 0.5 mm in duplicate with a Lange skinfold caliper (Cambridge Scientific Industries, Cambridge, MD) at the following sites: triceps, biceps, subscapular, thigh, and suprailiac.

The U.S. Army, Navy and Air Force equations used to predict the body fat of women will be compared against other body composition models.

Table 5. Fat mass prediction equations for the U.S. Army, Navy, and Air Force servicewomen(7)

U.S. Army

$$\text{FM}(\%) = 105.3 \times \log_{10}(\text{weight}) - 0.20 \times \text{waist} - 0.533 \times \text{neck} - 1.574 \times \text{forearm} + 0.173 \times \text{hip} - 0.515 \times \text{height} - 35.6$$

U.S. Navy

$$\text{Density} = -0.35 \times \log_{10}(\text{abdomen I} + \text{hip} - \text{neck}) + 0.221 \times \log_{10}(\text{height}) + 1.296$$

$$\text{FM}(\%) = 100 \times [(4.95/\text{density}) - 4.5]$$

U.S. Air Force

$$\text{FFM}(\text{kg}) = 1.619 \times \text{forearm} + 0.311 \times \text{height} - 47.76$$

$$\text{FM}(\%) = 100 \times (\text{weight} - \text{FFM})/\text{weight}$$

Weight is in kg; height and body circumferences are in cm.

Note: As of 1996 the Air Force adopted the Navy equations for prediction of fat mass.

Hydrodensitometry. An underwater weighing system utilizing "force cube" transducers (Precision Biomedical Systems, Inc., State College, PA) will be used for body density measurements (52). Each subject will be requested to urinate, defecate, shampoo, shower, change to a swimming suit, and then submerge herself and exhale maximally while her body weight in the water is being measured. Residual lung volume will be measured using the simplified nitrogen washout method (53). Body density will be calculated from body weights in and out of the water and residual lung volume.

Total Body Potassium (TBK). Total body ^{40}K content of each subject will be measured using the CNRC low-background whole-body counter. The CNRC counting system consists of a total of 30 NaI(Tl) detectors (each 10 cm x 10 cm x 45 cm) for a total detection volume of 135,000 cc. The detectors are arranged into two arrays positioned above and below the bed with the subject in a supine position. The subject will lie supine for 15 minutes while the body's natural gamma ray signal is recorded. The gamma signal is directly proportional to the amount of potassium in the body. The precision of the ^{40}K counting is <1% in adults.

Dual Energy X-ray Absorptiometry (DEXA). Composition of the total body and major subregions will be measured using a Hologic QDR 2000W system (Hologic, Inc., Madison, WI). Subregions include the head, thoracic spine, ribs, lumbar spine, pelvis, arms, and legs. The whole body scan takes approximately 15 minutes with the subject lying supine. The low dose (<0.01 mSv) has allowed for IRB approval of research measurements in healthy individuals. Over a 2 y period the average precision for bone mineral density using the spine phantom was $\pm 0.6\%$.

Total Body Nitrogen (TBN). Prompt-Gamma Activation Analysis will be used to measure TBN. The subject is placed in a very weak neutron beam: neutrons interact with body tissues and generate a gamma signal that can be detected external to the body. This technique induces activities of interest that are very short-lived (less than 1 μsec) which requires the detection system to be included in the irradiator assembly. Two shielded, collimated AmBe sources provide a bilateral beam through which the subject is scanned. Four large volume NaI(Tl) detectors with custom designed neutron/gamma shielding are positioned at 90° to both the bed and sources. The scan time is about 15 min; the dose is less than 0.3

mSv (comparable to a chest x-ray). In pigs, the *in vivo* precision was determined to be 3.5% for body nitrogen.

Muscle and Nonmuscle Mass. In the nonpregnant women TBK and TBN will be used to estimate the relative amounts of protein in muscle and nonmuscle components of the body and the mass of each component (54). The principle underlying this approach is that the K/N ratio for muscle is higher than that of nonmuscle tissues (3.03 vs 1.33 meq/g). Assuming the values of 30 g N/kg and 91 meq K/kg muscle, and 36 g N/kg and 48 meq K/kg nonmuscle, total muscle mass is equal to $(K-1.33N)/51.0$ and total nonmuscle mass is equal to $(3.03 N-K)/61.2$.

Body Composition Models. The Fuller (55) four-compartment model that combines total body water (TBW), body density and DEXA measurements will be used to compute fat-free mass (FFM) and fat mass (FM). A constant ratio of bone mineral to non-osseous mineral (0.8191:0.1809) is assumed. The equation used is: $FM(kg) = 2.747 \text{ body volume} - 0.710 \text{ TBW} + 1.460 \text{ total body bone ash} - 2.050 \text{ weight}$. Gestational values for total body bone ash will be linearly interpolated from pre-pregnancy and day 15 postpartum measurements. Regional fat deposition during pregnancy and mobilization postpartum will be studied through all reproductive phases using anthropometry and DEXA. Subregions described by DEXA include the head, thoracic spine, ribs, lumbar spine, pelvis, arms, and legs.

Total Body Water (TBW). After collection of baseline saliva samples, 40 mg $^2\text{H}_2\text{O}/\text{kg}$ in the form of water will be administered orally at 8 and 36 wk of gestation, and 2 and 24 wk postpartum. The ^2H abundance in saliva samples will be measured by gas-isotope-ratio mass spectrometry. TBW will be calculated from the elevation of ^2H abundance in the 4-h and 6-h postdose samples. TBW will be determined from $^2\text{H}_2^{18}\text{O}$ dilution prior to pregnancy and at 22 wk gestation, and at 24 wk postpartum (see TEE).

Total Energy Expenditure. Total energy expenditure (TEE) over a 14-day period will be calculated from the fractional turnover rates of ^2H and ^{18}O following oral ingestion of 100 mg/kg $^2\text{H}_2\text{O}$ and 125 mg ^{18}O as water (56). Isotope dilution spaces will be used to compute TBW. Baseline saliva samples will be collected from each subject. Subsequently, one daily saliva sample will be collected by each subject at home for the next 14 days. The ^2H and ^{18}O abundances of the saliva samples will be measured by gas-isotope-ratio mass spectrometry. Carbon dioxide production (VCO_2) will be calculated from the dilution spaces and fractional turnover rates of ^2H and ^{18}O using the multipoint slope-intercept method of calculation. Fractionated insensible water losses will be calculated from ventilatory volume and body surface area, both expressed as functions of CO_2 production. Respiratory quotient will be computed from the food quotient based on dietary records (57). TEE will be calculated using the Weir equation (58).

Room Respiration Calorimetry. Oxygen consumption (VO_2) and VCO_2 of mothers will be measured and monitored continuously in a room-sized indirect calorimeter for 24-h (59). Energy expenditure will be calculated from VO_2 and VCO_2 . Performance tests with N_2 and CO_2 infusions demonstrated that the accuracy of individual measurements of VO_2 and VCO_2 were 3%. System response to a step change exceeded 90% in 4 min. Sleeping metabolic rate (SMR), and basal metabolic rate (BMR) will be extracted from specific time periods. BMR will be measured for 40 minutes while the subject lies quietly 30 min after awaking and voiding. Fat and carbohydrate net utilization will be computed using 24-h excretion rates of urinary nitrogen according to Livesey (60).

The heart rate of the subject will be monitored and recorded continuously by telemetry (Dynascope 3300 Telemetry System, Fukuda Denshi America). During the 24-h period in the calorimeter, subjects

will be asked to adhere to a schedule of feeding, sleeping and exercise times. Energy intake will be controlled according to body weight. Subjects will walk twice for 10 minute each at 2.5 and 3.5 mph, no grade, on a treadmill.

Physical Activity. Physical activity level, our major independent variable, will be determined by combining the doubly-labeled water method with respiration calorimetry. Physical activity level will be quantified in terms of MJ/d as the difference between TEE and BMR. Physical activity level will be described qualitatively using the Minnesota Leisure Time Activity Scale (61) combined with the Health Insurance Plan (HIP) Questionnaire for assessment of occupational activity (62).

Physical Fitness. Maximal VO_2 will be measured in nonpregnant women who will undergo a stepwise increase in exercise intensity until volitional fatigue is achieved (3-Corval 400, LODE B.V., Groningen, Holland and 2-Combi Cycle EX80, Combi, Co., LTD. Tokyo). Subjects will cycle for 4 min at 50 watts for steady state determination of VO_2 , and then the workload will be increased by 25 watts every minute. VO_2 consumption will be measured continuously via the open circuit technique with a metabolic cart (SensorMedics 2900; Yorba Linda, CA). Heart rate and blood pressure will be monitored. Steady-state heart rate and VO_2 values will be used to estimate VO_{2max} .

Submaximal VO_2 also will be measured in the room respiration calorimeter, unimpeded by a mouthpiece apparatus. In the morning and afternoon women will walk for 20 minute at 2.5 mph, no grade, on a treadmill (905E, Precor, Bothell, WA); heart rate will not be allowed to exceed 140 bpm in pregnant women.

Strength. Prior to the 1 RM strength assessments, the subject will practice using the equipment to become familiar with the proper exercise techniques and to prevent injuries. The upper and lower body 1 RM strength tests will be done utilizing the Cybex Smith Press and the Cybex Latissimus Pulldown (upper body), and the Cybex Modular Leg Press and the Cybex Leg Extension (lower body). The 1 RM is defined as the maximum amount of weight that can be lifted successfully one time only. Starting with a weight used in the practice session, the subjects will attempt lifts with gradually increasing weight (10% at first, decreasing to 5 and 2.5% as difficulty becomes evident). Successive attempts will be made with a 90-s rest between attempts until failure occurs. These measurements will be done at the Texas Children's Hospital Wellness Center within the Department of Pediatrics.

Infant Outcome. Birthweight, length and gestational age will be recorded from medical records.

Biochemical Analysis. A 12-h fasting blood sample will be obtained for the following analyses. A Complete Blood Count and Differential including hemoglobin and hematocrit will be performed by Smith Kline Beecham Clinical Laboratories. Standard techniques of flow cytometry, automated cytochemistry and microscopy are used. Serum iron and total iron binding capacity will be measured using atomic absorption spectrophotometry (Perkin-Elmer Corp., Norwalk, Conn). Serum ferritin will be determined by ^{125}I -radioimmunoassay (Diagnostic Products Corp., CA). To correct calorimetry data, 24-h urinary nitrogen will be determined by the Kjeldahl method (Tecator, Höganäs, Sweden).

Statistical Analysis. Analysis of variance and covariance with repeated measures will be used to test our hypothesis. The grouping factor will be pre-pregnancy BMI stratification. The within factors will be pregnancy status (antepartum or postpartum), and time (levels= 0, 8, 22, 36 wk gestation; 2, 6 and 24 wk postpartum). The major independent variable to be tested is physical activity level. The dependent variables to be analyzed under this model will be weight, FFM, fat mass, energy expenditure, and physical fitness. Covariates to be included in the model are age, parity, iron status and infant feeding

mode. Pre-pregnancy BMI will be treated also as a continuous variable in a separate analysis. We will also use multiple regression analysis to investigate relationships between the various dependent variables. Statistical analyses will be performed using BMDP (63) and Minitab (64) statistical packages. Microsoft ACCESS will be used for data management.

7. RESULTS and DISCUSSION

Data Presentation:

Prepregnancy data will be presented on 117 women enrolled into the study. The body composition, energy expenditure, energy requirements, physical fitness and strength will be examined in this data set of healthy women. The effects of physical activity level on pregnancy-induced changes in weight, body composition, physical fitness and energy requirements, and assessment of postpartum return to weight and physical fitness standards will be examined in the women who completed normal pregnancies. The sample has been subdivided according to the women's prepregnancy BMI. The low BMI group is defined as BMI < 19.8, the normal BMI group as BMI > 19.8 and < 26.0, and the high BMI group as BMI > 26.0. Military-eligible women fall into the low and normal BMI groups, while women in the high BMI group exceed military weight standards and would not be eligible for military service. However, since at any given time the weights of many women in the military exceed weight standards, it is important to recognize that some women enter pregnancy "out of standard," compounding the difficulty of losing weight postpartum.

Subject Description:

A total of 117 women completed the baseline measurements prior to pregnancy. Seventy-six women became pregnant of whom 65 women completed their pregnancies at term. All but four of these 65 women have completed the entire protocol through 6 mo postpartum. Eleven women were discontinued because of twin/triple births, miscarriages, or other complications. Forty-one women discontinued the study because of fertility problems, change of family plans, relocation or lost to follow-up.

Reproductive History:

N		65
Maternal age	y	30.7 ± 4.2
Menarche age	y	12.9 ± 1.4
Menses interval	d	29.6 ± 5.4
Menses duration	d	4.7 ± 1.1
Gravidity		1.0 ± 1.2
Parity		0.5 ± 0.6
Education	y	16.8 ± 2.3
Ethnicity	W/B/H/A	50/7/6/2

*Mean ± SD

Pregnancy Outcome:

Of the 61 deliveries, all were term, healthy infants, except for five preterm infants born between 35-36 weeks of gestation. Birthweight and gestational age tended to be lower in the low BMI group (3.3 ± 0.5 kg, 38.5 ± 1.9 wk) compared with the normal BMI group (3.5 ± 0.4 kg, 39.1 ± 1.3 wk) and the high BMI group (3.6 ± 0.4 kg, 39.4 ± 1.9 wk).

		Male	Female	All
N		30	31	61
Gestational age	wk	$39.2 \pm 1.4^*$	38.9 ± 1.6	39.1 ± 1.5
Maternal weight gain	kg	15.5 ± 4.3	15.5 ± 5.4	15.5 ± 4.9
Birth weight	kg	3.6 ± 0.4	3.5 ± 0.5	3.5 ± 0.5
Birth length	cm	51.5 ± 2.1	50.5 ± 2.2	51.0 ± 2.2
Apgar score 1 min		8.0 ± 0.8	8.2 ± 0.6	8.1 ± 0.7
Apgar score 5 min		8.9 ± 0.5	9.1 ± 0.3	9.0 ± 0.4

*Mean \pm SD

Iron Status:

Iron status is assessed throughout the reproductive cycle, because of its potential effect on pregnancy outcome, work capacity, and physical fitness. Iron metabolism and erythrocyte indices are altered by the physiological changes of normal pregnancy. Serum iron decreases with increasing duration of pregnancy due to plasma volume expansion. Serum transferrin increases by ~ 2.4 fold due to estrogen-stimulated hepatic protein synthesis; iron saturation accordingly decreases. Serum ferritin decreases as iron stores are used for expansion of erythrocyte mass. Because of these changes in iron metabolism, it is difficult to distinguish anemia due to iron deficiency from dilutional anemia. To provide a more sensitive indicator of iron deficiency in pregnancy, we are using an EIA assay for serum transferrin receptor (Ramco Laboratories Inc.). In contrast to ferritin, serum transferrin receptor is not elevated with inflammation or infection. Any elevation of serum transferrin receptor during pregnancy is due to maternal iron deficiency. Serum ferritin, transferrin and transferrin receptor will be monitored throughout the reproductive cycle to evaluate the impact of iron status on pregnancy outcome and postpartum recovery. The data also will be analyzed for evidence of an effect of iron deficiency on basal energy expenditure and submaximal work performance.

Prior to pregnancy, there is no evidence of iron deficiency or anemia in these women, based on normal values for hemoglobin, hematocrit, serum iron, iron binding capacity and percent iron saturation. Serum ferritin, an indicator of iron stores, and transferrin receptor, an index of tissue iron need, also fell within the normal limits for all women before pregnancy. The transferrin receptor initially decreases in pregnancy due to hemodilution and then increases in the third trimester in all groups. The increasing levels of transferrin receptor with advancing pregnancy and postpartum indicate the appearance of mild iron deficiency in some women.

Table 8. Iron status							
Low BMI Group		Pre-pregnancy n=17	8 wk pregnancy n=14	22 wk pregnancy n=18	36 wk pregnancy n=13	6 wk postpartum n=12	24 wk postpartum n=13
Serum iron	mcg/dl	92.76 ± 29.14*	118.29 ± 24.53	119.17 ± 35.13	94.62 ± 35.39	78.83 ± 25.16	109.69 ± 32.30
Iron binding capacity	mcg/dl	300.53 ± 38.81	301.36 ± 30.56	374.78 ± 50.82	421.85 ± 51.16	296.42 ± 53.28	332.46 ± 57.24
Iron saturation	%	31.88 ± 12.81	39.50 ± 9.09	32.61 ± 11.19	22.54 ± 8.34	27.92 ± 12.68	33.69 ± 11.54
White blood cell count	x10 ⁹ /L	5.53 ± 1.38	8.21 ± 1.66	9.73 ± 1.44	9.01 ± 1.61	6.28 ± 1.09	5.74 ± 1.37
Red blood cell count	x10 ¹² /L	4.41 ± 0.31	4.22 ± 0.31	3.64 ± 0.26	3.73 ± 0.34	4.20 ± 0.33	4.50 ± 0.28
Hemoglobin	g/dl	13.64 ± 0.78	13.15 ± 0.87	11.78 ± 0.75	11.97 ± 1.09	12.89 ± 0.75	13.74 ± 0.78
Hematocrit	%	40.30 ± 2.54	38.81 ± 2.78	34.41 ± 2.40	35.23 ± 3.60	38.08 ± 2.46	40.62 ± 2.39
Mean corpuscular volume	fL	91.11 ± 3.08	92.06 ± 3.15	94.52 ± 3.04	94.56 ± 3.78	90.76 ± 4.64	90.28 ± 3.07
Mean corpuscular hemoglobin	pg	30.88 ± 0.87	31.21 ± 1.27	32.41 ± 1.18	32.12 ± 1.19	30.78 ± 1.84	30.61 ± 1.21
Mean corpuscular hemoglobin concentration	%	33.89 ± 0.55	33.89 ± 0.58	34.26 ± 0.62	33.99 ± 0.67	33.88 ± 0.69	33.85 ± 0.48
Ferritin	ng/ml	42.12 ± 26.90	66.43 ± 33.58	25.53 ± 14.99	18.54 ± 8.09	55.50 ± 36.41	49.23 ± 32.04
Transferrin receptor	mg/ml	4.54 ± 0.90	4.05 ± 1.07	4.30 ± 1.12	5.87 ± 1.77	5.27 ± 1.27	5.17 ± 1.11
Normal BMI Group		Pre-pregnancy n=32	8 wk pregnancy n=31	22 wk pregnancy n=34	36 wk pregnancy n=32	6 wk postpartum n=34	24 wk postpartum n=33
Serum iron	mcg/dl	102.59 ± 26.32	121.29 ± 24.55	112.68 ± 34.78	114.84 ± 47.83	92.74 ± 31.02	105.64 ± 33.88
Iron binding capacity	mcg/dl	309.91 ± 39.42	312.52 ± 36.56	385.68 ± 54.17	453.28 ± 76.14	326.71 ± 74.43	329.03 ± 43.20
Iron saturation	%	33.31 ± 8.59	38.97 ± 7.34	29.26 ± 8.62	26.00 ± 11.68	29.41 ± 10.34	32.33 ± 10.79
White blood cell count	x10 ⁹ /L	5.62 ± 1.59	7.93 ± 2.23	9.55 ± 2.67	10.02 ± 2.87	5.39 ± 1.05	5.59 ± 1.17
Red blood cell count	x10 ¹² /L	4.43 ± 0.28	4.22 ± 0.34	3.77 ± 0.29	3.94 ± 0.36	4.22 ± 0.29	4.49 ± 0.32
Hemoglobin	g/dl	13.72 ± 0.75	13.18 ± 0.89	12.05 ± 0.88	12.44 ± 1.05	12.88 ± 0.83	13.68 ± 0.75
Hematocrit	%	40.38 ± 2.31	38.55 ± 2.65	35.18 ± 2.46	36.63 ± 3.00	38.06 ± 2.28	40.07 ± 2.86
Mean corpuscular volume	fL	91.33 ± 3.48	91.43 ± 3.53	93.26 ± 3.83	93.02 ± 4.53	90.32 ± 4.42	89.06 ± 5.72
Mean corpuscular hemoglobin	pg	31.00 ± 1.19	31.28 ± 1.18	31.95 ± 1.47	31.57 ± 1.75	30.58 ± 1.69	30.59 ± 1.55
Mean corpuscular hemoglobin concentration	%	33.93 ± 0.56	34.20 ± 0.59	34.25 ± 0.79	33.92 ± 0.69	33.82 ± 0.65	33.97 ± 0.55
Ferritin	ng/ml	49.19 ± 39.54	79.26 ± 55.79	37.62 ± 44.53	18.20 ± 6.82	56.76 ± 36.16	40.97 ± 21.58
Transferrin receptor	mg/ml	4.56 ± 1.27	4.30 ± 1.24	4.45 ± 1.48	5.53 ± 1.88	5.36 ± 2.01	5.40 ± 1.55
High BMI Group		Pre-pregnancy n=10	8 wk pregnancy n=9	22 wk pregnancy n=9	36 wk pregnancy n=9	6 wk postpartum n=11	24 wk postpartum n=9
Serum iron	mcg/dl	111.70 ± 19.69	119.22 ± 17.93	97.11 ± 28.04	93.11 ± 25.80	65.91 ± 26.27	106.78 ± 53.65
Iron binding capacity	mcg/dl	325.20 ± 43.84	334.33 ± 38.27	374.44 ± 44.71	424.33 ± 28.04	318.64 ± 52.19	359.56 ± 69.04
Iron saturation	%	34.20 ± 5.83	36.11 ± 6.39	25.78 ± 6.42	21.78 ± 5.80	20.55 ± 9.33	31.33 ± 17.94
White blood cell count	x10 ⁹ /L	6.62 ± 2.04	8.54 ± 1.79	10.51 ± 3.01	10.44 ± 1.97	6.46 ± 1.28	6.11 ± 1.36
Red blood cell count	x10 ¹² /L	4.46 ± 0.36	4.41 ± 0.33	3.89 ± 0.23	4.01 ± 0.33	4.34 ± 0.42	4.48 ± 0.37
Hemoglobin	g/dl	13.60 ± 1.01	13.39 ± 0.98	12.06 ± 0.67	12.36 ± 1.15	12.77 ± 1.31	13.38 ± 1.20
Hematocrit	%	40.64 ± 3.57	39.06 ± 3.12	35.38 ± 1.92	36.61 ± 3.65	38.22 ± 4.02	40.29 ± 3.30
Mean corpuscular volume	fL	89.31 ± 2.43	88.42 ± 1.90	91.90 ± 2.87	91.23 ± 3.14	88.02 ± 3.41	89.96 ± 4.13
Mean corpuscular hemoglobin	pg	29.89 ± 0.46	30.39 ± 0.66	31.05 ± 0.94	30.86 ± 1.23	29.46 ± 1.13	29.89 ± 0.82
Mean corpuscular hemoglobin concentration	%	33.47 ± 0.90	34.35 ± 0.44	34.04 ± 0.35	33.81 ± 0.58	33.44 ± 0.57	33.27 ± 1.36
Ferritin	ng/ml	47.70 ± 22.20	60.13 ± 29.08	32.91 ± 28.61	23.00 ± 16.92	49.82 ± 43.06	34.13 ± 27.38
Transferrin receptor	mg/ml	4.64 ± 1.31	3.88 ± 1.00	4.16 ± 0.80	5.58 ± 2.25	7.16 ± 2.57	7.73 ± 2.72

*Mean ± SD

Anthropometry

The anthropometric measurements reflect a wide array of body sizes. Prior to conception, the mean weight was 49.8 ± 3.8 kg for the low BMI group, 59.6 ± 6.1 kg for the normal BMI group, and 77.3 ± 10.2 kg for the high BMI group. Anthropometric measurements will be used to monitor gestational weight gain/postpartum weight loss, and site-specific deposition/postpartum mobilization of subcutaneous body fat. Total gestational weight gain at delivery was 15.5 ± 4.9 kg. Maternal weight gain tended to be higher in the high (17.7 ± 5.9 kg) and normal BMI groups (15.2 ± 4.8 kg) than in the low BMI group (14.7 ± 4.0 kg). Postpartum weight retention at 6 mo averaged 2.7 ± 3.9 kg, but was greater in the high BMI group (4.0 ± 4.4 kg) than the low and normal BMI groups (2.4 ± 3.5 kg). BMI had increased by approximately one unit in all groups.

Low BMI Group		Pre-pregnancy	8 wk pregnancy	22 wk pregnancy	36 wk pregnancy	2 wk postpartum	6 wk postpartum	24 wk postpartum
		n=18	n=17	n=18	n=15	n=14	n=14	n=14
Weight	kg	49.83 \pm 3.82*	51.78 \pm 5.34	57.42 \pm 4.72	62.97 \pm 4.70	54.59 \pm 4.75	54.22 \pm 5.26	51.36 \pm 4.52
Height	cm	162.27 \pm 5.32	162.38 \pm 5.22	162.62 \pm 5.42	163.40 \pm 5.09	161.66 \pm 5.70	161.60 \pm 5.57	161.61 \pm 5.44
BMI	kg/m ²	18.90 \pm 0.73	19.61 \pm 1.51	21.71 \pm 1.46	23.66 \pm 1.60	20.89 \pm 1.54	20.76 \pm 1.66	19.65 \pm 1.26
Head circumference	cm	53.89 \pm 1.31	53.95 \pm 1.25	53.88 \pm 1.23	54.33 \pm 1.09	54.13 \pm 1.36	54.26 \pm 1.23	54.00 \pm 1.39
Neck circumference	cm	29.86 \pm 1.35	29.78 \pm 1.09	30.36 \pm 1.14	31.01 \pm 1.07	30.53 \pm 1.37	30.51 \pm 1.33	30.16 \pm 1.36
Chest circumference	cm	81.87 \pm 2.44	85.42 \pm 3.75	90.30 \pm 3.74	92.29 \pm 3.37	90.14 \pm 3.83	89.51 \pm 2.95	85.76 \pm 3.14
Waist circumference	cm	63.56 \pm 2.77	67.43 \pm 4.56	76.32 \pm 4.78	87.30 \pm 3.69	72.29 \pm 4.91	71.34 \pm 4.86	67.16 \pm 4.56
Hip circumference	cm	88.49 \pm 3.14	89.48 \pm 4.49	93.96 \pm 3.90	96.00 \pm 3.87	93.26 \pm 4.31	92.74 \pm 4.04	89.34 \pm 4.39
Thigh circumference	cm	44.48 \pm 2.74	44.13 \pm 3.70	46.06 \pm 2.72	47.05 \pm 2.35	45.23 \pm 2.20	44.86 \pm 2.46	44.27 \pm 2.83
Calf circumference	cm	32.97 \pm 1.80	33.08 \pm 2.03	33.61 \pm 1.78	34.11 \pm 1.53	32.56 \pm 1.58	32.69 \pm 1.81	32.56 \pm 1.87
Wrist circumference	cm	13.85 \pm 0.45	13.77 \pm 0.59	13.89 \pm 0.55	13.99 \pm 0.39	13.79 \pm 0.60	13.91 \pm 0.55	14.64 \pm 2.75
AF Forearm circumference	cm	21.88 \pm 0.96	21.71 \pm 1.07	21.98 \pm 1.05	22.49 \pm 0.76	22.04 \pm 0.91	22.01 \pm 0.88	21.94 \pm 1.05
Army forearm circumference	cm	22.10 \pm 1.01	21.98 \pm 1.10	22.21 \pm 1.00	22.63 \pm 0.76	22.46 \pm 0.83	22.11 \pm 0.87	22.16 \pm 1.16
Upper arm circumference	cm	23.89 \pm 1.59	24.46 \pm 1.73	24.98 \pm 1.76	25.33 \pm 1.84	24.46 \pm 2.31	24.66 \pm 1.57	24.52 \pm 1.62
Triceps	mm	11.74 \pm 3.60	13.80 \pm 4.23	14.55 \pm 4.90	14.88 \pm 4.91	13.06 \pm 5.22	13.42 \pm 5.44	13.88 \pm 5.16
Biceps	mm	3.63 \pm 1.16	4.16 \pm 1.44	4.31 \pm 1.52	4.32 \pm 1.80	4.18 \pm 1.76	3.86 \pm 1.31	3.52 \pm 1.17
Subscapular	mm	8.49 \pm 2.21	9.72 \pm 3.22	12.78 \pm 4.69	13.56 \pm 5.95	14.49 \pm 6.78	12.66 \pm 5.42	9.32 \pm 2.89
Suprailiac	mm	8.73 \pm 3.51	12.86 \pm 7.09	18.45 \pm 7.91	20.77 \pm 8.99	16.36 \pm 8.38	14.84 \pm 7.16	9.36 \pm 5.35
Thigh	mm	18.47 \pm 7.04	22.62 \pm	23.26 \pm 8.56	23.81 \pm 7.63	21.85 \pm 8.83	21.60 \pm 8.25	18.95 \pm 8.09
Sagittal diameter	cm	12.97 \pm 0.71				15.08 \pm 1.56	15.02 \pm 1.39	13.61 \pm 0.80

Normal BMI Group		Pre-pregnancy	8 wk pregnancy	22 wk pregnancy	36 wk pregnancy	2 wk postpartum	6 wk postpartum	24 wk postpartum
		n=35	n=31	n=34	n=32	n=34	n=35	n=34
Weight	kg	59.62 \pm 6.13*	60.60 \pm 6.66	65.56 \pm 7.69	72.38 \pm 8.64	65.16 \pm 8.43	64.07 \pm 8.83	62.04 \pm 8.22
Height	cm	163.85 \pm 5.60	164.21 \pm 5.60	164.31 \pm 5.52	163.60 \pm 5.55	163.50 \pm 5.49	163.51 \pm 5.55	163.66 \pm 5.81
BMI	kg/m ²	22.17 \pm 1.52	22.43 \pm 1.73	24.24 \pm 2.18	26.99 \pm 2.36	24.33 \pm 2.53	23.91 \pm 2.55	23.11 \pm 2.30
Head circumference	cm	54.94 \pm 1.41	54.79 \pm 1.13	55.12 \pm 1.69	55.09 \pm 1.45	55.28 \pm 1.54	55.23 \pm 1.47	55.13 \pm 1.42

Table 9. Anthropometry ... (cont)

Neck circumference	cm	31.45 ± 1.44	31.61 ± 1.63	31.43 ± 1.74	32.06 ± 1.79	32.15 ± 1.59	32.11 ± 1.67	31.85 ± 1.61
Chest circumference	cm	88.39 ± 4.13	90.54 ± 4.28	94.91 ± 4.79	97.57 ± 5.32	96.08 ± 5.32	95.45 ± 5.32	92.09 ± 5.23
Waist circumference	cm	70.70 ± 3.74	72.34 ± 4.09	81.03 ± 4.84	93.41 ± 5.53	78.69 ± 4.67	76.87 ± 4.47	74.19 ± 6.36
Hip circumference	cm	96.36 ± 4.57	97.33 ± 5.15	99.97 ± 5.86	102.43 ± 6.71	100.84 ± 6.98	100.20 ± 7.19	97.45 ± 6.27
Thigh circumference	cm	49.55 ± 3.26	48.84 ± 3.53	49.77 ± 5.05	51.59 ± 4.18	50.20 ± 4.25	50.05 ± 4.30	50.11 ± 4.33
Calf circumference	cm	35.62 ± 2.37	35.31 ± 2.43	35.76 ± 2.55	36.84 ± 2.55	35.69 ± 2.66	35.55 ± 2.90	35.66 ± 2.69
Wrist circumference	cm	14.47 ± 0.60	14.46 ± 0.64	14.44 ± 0.69	14.63 ± 0.75	14.55 ± 0.71	14.57 ± 0.66	14.46 ± 0.74
AF Forearm circumference	cm	23.29 ± 1.14	23.30 ± 1.22	23.34 ± 1.51	23.70 ± 1.64	23.48 ± 1.52	23.44 ± 1.55	23.35 ± 1.51
Army forearm circumference	cm	23.76 ± 1.30	23.50 ± 1.27	23.54 ± 1.52	24.07 ± 1.77	23.81 ± 1.64	23.63 ± 1.53	23.74 ± 1.92
Upper arm circumference	cm	26.64 ± 1.73	26.76 ± 1.95	26.95 ± 2.29	27.57 ± 2.31	27.10 ± 2.56	27.18 ± 2.53	27.41 ± 2.42
Triceps	mm	17.15 ± 4.72	17.04 ± 4.28	18.50 ± 6.20	19.01 ± 6.94	17.81 ± 7.34	18.19 ± 7.59	19.41 ± 7.37
Biceps	mm	5.91 ± 2.28	6.10 ± 2.32	7.34 ± 3.57	7.42 ± 3.20	6.51 ± 2.98	6.33 ± 2.67	7.17 ± 3.57
Subscapular	mm	15.34 ± 6.11	15.59 ± 5.84	18.49 ± 8.15	20.54 ± 9.01	19.56 ± 7.97	19.32 ± 7.89	16.90 ± 7.75
Suprailiac	mm	17.65 ± 8.23	19.32 ± 8.76	26.11 ± 9.09	29.50 ± 8.87	25.65 ± 10.45	24.70 ± 10.70	20.44 ± 10.22
Thigh	mm	26.54 ± 7.58	25.55 ± 7.07	30.44 ± 10.53	32.86 ± 11.24	30.98 ± 11.76	30.81 ± 11.58	27.98 ± 9.26
Sagittal diameter	cm	14.83 ± 1.14				17.46 ± 3.77	16.39 ± 1.72	15.35 ± 1.42
High BMI Group		Pre-pregnancy	8 wk pregnancy	22 wk pregnancy	36 wk pregnancy	2 wk postpartum	6 wk postpartum	24 wk postpartum
		n=12	n=10	n=12	n=11	n=11	n=11	n=8
Weight	kg	77.27 ± 10.15*	81.83 ±	85.80 ± 10.45	93.84 ± 10.59	84.66 ± 10.88	83.86 ± 11.37	77.98 ± 6.69
Height	cm	163.67 ± 6.14	164.68 ± 5.97	163.59 ± 6.48	164.13 ± 6.74	163.68 ± 6.34	163.56 ± 6.42	164.04 ± 7.22
BMI	kg/m ²	28.77 ± 2.56	30.14 ± 3.48	32.02 ± 2.96	34.82 ± 3.22	31.54 ± 3.10	31.29 ± 3.28	29.01 ± 2.39
Head circumference	cm	55.61 ± 0.76	55.74 ± 0.88	55.79 ± 0.68	55.93 ± 0.65	55.88 ± 0.62	55.89 ± 0.79	55.75 ± 0.66
Neck circumference	cm	34.22 ± 1.49	34.73 ± 2.02	34.33 ± 1.50	34.98 ± 1.49	34.98 ± 1.15	34.70 ± 1.58	34.43 ± 1.39
Chest circumference	cm	103.62 ± 8.61	105.24 ± 8.82	108.82 ± 6.73	111.39 ± 7.27	108.47 ± 7.37	109.27 ± 8.30	102.23 ± 6.21
Waist circumference	cm	85.68 ± 8.26	89.30 ± 9.63	95.08 ± 6.04	106.95 ± 6.97	92.40 ± 7.28	91.40 ± 8.25	85.86 ± 5.54
Hip circumference	cm	110.02 ± 6.71	112.24 ± 7.40	113.93 ± 7.60	116.29 ± 7.69	110.96 ± 13.21	113.22 ± 7.86	107.81 ± 5.34
Thigh circumference	cm	55.60 ± 2.73	56.48 ± 3.41	58.34 ± 3.11	59.91 ± 3.52	57.84 ± 3.68	58.05 ± 3.42	56.18 ± 2.94
Calf circumference	cm	38.13 ± 1.25	38.49 ± 1.36	39.37 ± 1.23	40.40 ± 1.65	38.97 ± 1.65	38.76 ± 1.99	38.87 ± 1.67
Wrist circumference	cm	15.09 ± 0.55	15.15 ± 0.67	15.29 ± 0.52	15.61 ± 0.55	15.42 ± 0.59	15.48 ± 0.49	15.22 ± 0.48
AF Forearm circumference	cm	25.77 ± 1.33	26.00 ± 1.49	26.38 ± 1.37	26.78 ± 1.32	26.31 ± 1.20	26.38 ± 1.40	26.18 ± 0.99
Army forearm circumference	cm	25.96 ± 1.33	26.40 ± 1.49	26.65 ± 1.43	27.53 ± 1.90	27.12 ± 1.37	26.63 ± 1.42	27.63 ± 3.36
Upper arm circumference	cm	33.27 ± 3.50	34.03 ± 3.69	33.93 ± 3.86	34.37 ± 3.65	32.84 ± 4.43	33.91 ± 3.81	33.26 ± 2.29
Triceps	mm	29.66 ± 5.90	29.77 ± 6.35	32.38 ± 6.65	32.52 ± 9.11	29.35 ± 7.47	30.12 ± 7.03	30.14 ± 4.99
Biceps	mm	15.30 ± 8.30	15.68 ± 6.92	15.33 ± 7.25	16.37 ± 8.89	14.26 ± 8.02	13.53 ± 6.17	13.45 ± 6.23
Subscapular	mm	26.52 ± 9.35	28.30 ± 9.89	28.90 ± 8.63	30.06 ± 9.77	28.44 ± 9.61	27.12 ± 8.03	24.58 ± 10.06
Suprailiac	mm	33.39 ± 5.21	33.80 ± 5.89	37.00 ± 5.97	40.58 ± 7.38	36.99 ± 5.83	36.44 ± 7.89	32.75 ± 11.18
Thigh	mm	40.16 ± 7.62	41.77 ± 7.21	47.63 ± 7.70	49.45 ± 8.05	46.95 ± 9.69	44.88 ± 8.51	41.43 ± 6.86
Sagittal diameter	cm	18.18 ± 1.87				21.14 ± 2.04	21.20 ± 2.31	19.56 ± 2.21

*Mean ± SD

Body Composition

Standard 2-component body composition models based on deuterium dilution, hydrodensitometry, or total body potassium, while applicable prior to conception, are invalid during pregnancy due to the expansion of body fluid compartments. To obtain more accurate estimates of FFM and FM in reproductive women, we have chosen the Fuller 4-component model, which minimizes assumptions regarding the hydration of FFM and the bone mineral content of dry FFM, by incorporating measurements of total body water, body density, and BMC (55). This model also provides more accurate body composition estimates in non-pregnant individuals. A comparison of FM estimated by the standard and multicomponent body composition models is presented in Table 10. Systematic differences between methods in the estimation of fat mass are evident throughout pregnancy.

Table 10. Fat mass (FM) estimated by deuterium dilution, hydrodensitometry, total body potassium, DEXA, and skinfold thicknesses, and multi-component models

Low BMI Group		Pre-pregnancy n=18	8 wk pregnancy n=17	22 wk pregnancy n=18	36 wk pregnancy n=14	2 wk postpartum n=14	6 wk postpartum n=14	24 wk postpartum n=14
Deuterium dilution								
Total body water (TBW)	kg	28.76 ± 2.86*	28.91 ± 2.53	32.31 ± 3.54	34.76 ± 4.07	30.20 ± 2.88	29.89 ± 3.06	27.48 ± 3.43
FM (TBW)	kg	10.43 ± 3.26	10.99 ± 2.88	13.33 ± 3.74	17.05 ± 3.84	13.76 ± 3.06	12.05 ± 2.95	13.82 ± 5.30
Hydrodensitometry (HD)								
Body Volume	l	47.47 ± 3.70	49.51 ± 5.26	54.80 ± 4.69	60.98 ± 5.14	52.83 ± 5.15	52.29 ± 5.30	49.50 ± 4.70
FM (HD)	kg	10.64 ± 2.69	11.91 ± 3.01	13.95 ± 2.85	14.32 ± 3.89	13.07 ± 4.03	14.46 ± 3.86	12.67 ± 3.84
Total body potassium (TBK)	g	93.35 ± 11.37	91.70 ± 8.86	93.64 ± 9.67	100.75 ± 11.19	94.81 ± 12.52	93.21 ± 11.82	93.40 ± 11.05
FM (TBK)	kg	10.17 ± 3.54	13.05 ± 3.77	17.36 ± 5.26	19.84 ± 4.04	14.04 ± 4.23	14.16 ± 4.93	11.72 ± 4.27
DEXA - Dual energy x-ray absorptiometry								
FM (DEXA)	kg	11.01 ± 3.36	***	***	***	15.17 ± 4.94	14.80 ± 4.16	12.90 ± 3.99
Skinfold thickness (Dumin & Womersley)								
FM (SF)	kg	10.79 ± 1.96	12.71 ± 3.16	15.59 ± 3.03	17.75 ± 3.75	14.41 ± 3.29	13.87 ± 3.37	11.76 ± 2.42
Multi-component models								
FM-Siri (TBW, HD)	kg	10.59 ± 2.78	11.27 ± 2.71	13.70 ± 3.07	15.82 ± 3.85	13.27 ± 3.26	12.56 ± 2.82	13.49 ± 4.76
FM-Fuller (TBW, HD, DEXA)	kg	10.77 ± 2.88	11.45 ± 2.79	13.63 ± 3.07	15.63 ± 3.86	13.41 ± 3.34	12.67 ± 2.82	14.29 ± 4.55

Table 10. Fat mass (FM) ...(cont)

Normal BMI Group		Pre-pregnancy n=35	8 wk pregnancy n=31	22 wk pregnancy n=34	36 wk pregnancy n=32	2 wk postpartum n=34	6 wk postpartum n=35	24 wk postpartum n=34
Deuterium dilution								
Total body water (TBW)	kg	31.90 ± 3.84*	31.76 ± 3.81	34.29 ± 3.94	39.06 ± 3.49	33.18 ± 3.59	32.42 ± 3.71	32.27 ± 3.73
FM (TBW)	kg	16.54 ± 4.44	17.10 ± 3.59	19.28 ± 5.21	21.23 ± 6.61	20.91 ± 5.69	19.71 ± 6.00	18.78 ± 5.76
Hydrodensitometry (HD)								
Body Volume	l	58.18 ± 5.66	58.61 ± 6.68	63.74 ± 7.84	70.69 ± 8.89	63.55 ± 8.82	62.42 ± 9.16	60.24 ± 8.53
FM (HD)	kg	16.94 ± 3.75	16.80 ± 3.92	19.21 ± 4.80	20.15 ± 5.01	19.93 ± 5.65	20.42 ± 5.59	18.54 ± 5.17
Total body potassium (TBK)	g	102.52 ± 13.26	99.92 ± 13.79	102.24 ± 13.77	108.32 ± 14.87	103.09 ± 11.75	100.97 ± 12.97	102.37 ± 13.60
FM (TBK)	kg	16.43 ± 3.73	18.07 ± 3.95	22.13 ± 4.49	25.41 ± 6.58	21.21 ± 6.13	20.87 ± 5.93	18.57 ± 5.45
DEXA - Dual energy x-ray								
FM (DEXA)	kg	17.18 ± 3.69	***	***	***	21.52 ± 5.91	21.17 ± 5.54	20.19 ± 5.18
Skinfold thickness (Durnin & Womersley)								
FM (SF)	kg	17.21 ± 3.56	17.68 ± 3.96	20.83 ± 5.06	23.91 ± 5.78	20.64 ± 5.65	20.18 ± 5.75	18.81 ± 5.45
Multi-component models								
FM-Siri (TBW, HD)	kg	16.79 ± 4.00	17.07 ± 3.61	19.48 ± 4.84	20.86 ± 5.96	20.33 ± 5.54	20.09 ± 5.71	18.97 ± 5.45
FM-Fuller (TBW, HD, DEXA)	kg	17.08 ± 4.02	17.31 ± 3.72	19.66 ± 4.91	20.86 ± 5.88	20.61 ± 5.63	20.28 ± 5.80	19.55 ± 5.18
High BMI Group		Pre-pregnancy n=12	8 wk pregnancy n=10	22 wk pregnancy n=12	36 wk pregnancy n=11	2 wk postpartum n=11	6 wk postpartum n=11	24 wk postpartum n=8
Deuterium dilution								
Total body water (TBW)	kg	35.62 ± 5.29*	37.02 ± 4.40	38.64 ± 4.03	41.70 ± 3.54	37.81 ± 3.89	36.69 ± 3.76	35.94 ± 3.96
FM (TBW)	kg	29.93 ± 6.25	32.58 ± 9.59	33.58 ± 10.32	35.38 ± 8.59	32.45 ± 8.24	31.23 ± 7.96	27.67 ± 6.07
Hydrodensitometry (HD)								
Body Volume	l	78.37 ± 10.94	81.07 ± 11.63	85.27 ± 10.89	94.22 ± 11.32	85.33 ± 11.46	83.42 ± 12.67	77.02 ± 7.28
FM (HD)	kg	31.74 ± 6.46	32.91 ± 7.69	34.98 ± 7.17	37.33 ± 8.43	35.40 ± 8.24	36.03 ± 9.32	30.38 ± 7.42
Total body potassium (TBK)	g	106.77 ± 10.02	105.05 ± 12.17	109.89 ± 8.30	120.62 ± 9.41	108.63 ± 8.68	106.57 ± 11.11	107.89 ± 9.83
FM (TBK)	kg	31.48 ± 8.46	37.05 ± 8.76	39.24 ± 9.72	42.19 ± 10.23	38.71 ± 9.82	36.65 ± 10.18	31.99 ± 7.14
DEXA - Dual energy x-ray								
FM (DEXA)	kg	34.15 ± 7.10	***	***	***	37.10 ± 9.23	37.05 ± 10.59	33.10 ± 7.69
Skinfold thickness (Durnin & Womersley)								
FM (SF)	kg	29.79 ± 6.08	31.24 ± 6.25	33.40 ± 6.11	37.83 ± 7.33	32.47 ± 6.72	32.00 ± 6.57	28.98 ± 5.14
Multi-component models								
FM-Siri (TBW, HD)	kg	30.81 ± 6.37	33.08 ± 9.06	33.99 ± 9.16	36.48 ± 7.86	33.49 ± 8.21	32.55 ± 7.95	27.99 ± 5.30
FM-Fuller (TBW, HD, DEXA)	kg	31.71 ± 6.36	31.88 ± 4.54	35.03 ± 9.08	34.49 ± 7.35	33.71 ± 8.32	32.68 ± 8.13	28.26 ± 5.65

*Mean±SD

Mean pre-pregnancy body fat was 10.8 ± 2.9 kg, or $21.6 \pm 5.2\%$ for the low BMI group, 17.1 ± 4.0 kg, or $28.3 \pm 5.7\%$ for the normal BMI group, and 31.7 ± 6.4 kg, or $39.6 \pm 4.7\%$ for the high BMI group by the Fuller 4-component model (Table 10). FM and % body fat at the pre-pregnancy time point estimated by military equations and by standard and multi-component models are listed in Table 11. Fat mass predicted from military equations depends on different anthropometric sites. The Army equation uses the waist and hip circumferences and builds in BMI, ie. weight and height measurements; the Navy equation predicts density from abdomen, hip and height; the Air Force has adopted the Navy equations. Women in the low and normal BMI groups would be eligible for enrollment into each branch of the Armed Services, according to the currently used Army and Navy equations. Only a few women in the high BMI group would be eligible for enrollment into the Army, and none would qualify for the Navy and Air Force.

Pre-pregnancy %FM derived from the Fuller 4-component model and military equations were highly correlated ($r^2 = 0.94-0.96$). Percent FM derived from the Fuller 4-component model and military equations also were significantly correlated, but to a lesser extent ($r^2 = 0.68-0.78$) (Figures 1-3). The obsolete Air Force equation significantly overestimated both the mean FM and % FM, and the Army and Navy equations underestimated mean FM and % FM in all BMI groups. The Army and Navy prediction equations are conservative, in that direct measurement of FM would have rendered more women ineligible for service.

Bland-Altman plots illustrate the differences in pre-pregnancy %FM derived from the Fuller 4-component model and military equations (Figures 4-6). The negative slope indicates that the difference between methods is a function of %FM. The underestimation of %FM was greater in fatter women. The mean differences (or relative biases), $-2 \pm 4\%$ for the Army and Navy equations, and $2 \pm 5\%$ for the Air Force, were statistically significant ($p=0.001-0.01$). The standard deviations or estimations of error imply wide confidence intervals for the prediction of %FM for individuals.







We will also evaluate these predictive equations for use in postpartum women at 6 mo postpartum. %FM derived from the Fuller 4-component model and military equations were highly correlated ($r^2 = 0.83-0.89$). Percent FM derived from the Fuller 4-component model and military equations also were significantly correlated ($r^2 = 0.93-0.96$) (Figures 1-3). Bland-Altman plots illustrate the differences in pre-pregnancy %FM derived from the Fuller 4-component model and military equations (Figures 4-6). The mean differences (or relative biases), $-0.8 \pm 8.6\%$ for the Army and $0.1 \pm 7.8\%$ for the Navy equations, and $2.8 \pm 7.6\%$ for the Air Force, were statistically significant ($p=0.01$). The standard deviations were larger at 6 mo postpartum than prepregnancy values, indicating wide confidence intervals for the prediction of %FM for individuals.







FM derived from the Fuller-4 component model indicated an increase of 2.1 ± 4.1 kg FM at 6 mo postpartum, with means of 2.6 ± 3.7 , 1.7 ± 4.1 and 3.0 ± 5.0 kg in the low, normal and high BMI groups.

Table 11. Body fat mass (FM) predicted from military equations and body composition models at the pre-pregnancy and 24 wk postpartum

Low BMI Group		Pre-pregnancy n=18	24 wk postpartum n=14		Pre-pregnancy n=18	24 wk postpartum n=14
Army FM	kg	10.61 ± 1.28*	11.73 ± 2.29	% WT	21.28 ± 1.74	22.52 ± 3.07
Navy FM	kg	9.79 ± 1.28	11.41 ± 2.70	% WT	19.63 ± 2.02	22.02 ± 3.70
Air Force FM	kg	11.71 ± 1.92	13.34 ± 2.91	% WT	23.41 ± 2.73	25.82 ± 4.03
FM (TBW)	kg	10.43 ± 3.26	13.82 ± 5.30	% WT	20.87 ± 5.92	26.65 ± 9.06
FM (HD)	kg	10.64 ± 2.69	12.67 ± 3.84	% WT	21.34 ± 4.94	24.33 ± 6.12
FM (TBK)	kg	10.17 ± 3.54	11.72 ± 4.27	% WT	20.47 ± 7.08	22.64 ± 7.30
FM (DEXA)	kg	11.01 ± 3.36	12.90 ± 3.99	% WT	22.12 ± 6.45	24.90 ± 6.97
FM-Siri (TBW, HD)	kg	10.59 ± 2.78	13.49 ± 4.76	% WT	21.22 ± 4.98	26.02 ± 8.05
FM-Fuller (TBW, HD, DEXA)	kg	10.77 ± 2.88	14.29 ± 4.55	% WT	21.56 ± 5.17	27.45 ± 7.59
Normal BMI Group		Pre-pregnancy n=35	24 wk postpartum n=34		Pre-pregnancy n=35	24 wk postpartum n=34
Army FM	kg	16.10 ± 2.99	17.61 ± 4.26	% WT	26.61 ± 3.32	28.14 ± 3.98
Navy FM	kg	15.87 ± 3.09	17.88 ± 4.92	% WT	26.22 ± 3.32	28.45 ± 4.77
Air Force FM	kg	19.18 ± 3.64	21.11 ± 5.42	% WT	31.65 ± 3.71	33.60 ± 4.78
FM (TBW)	kg	16.54 ± 4.44	18.78 ± 5.76	% WT	27.38 ± 6.62	29.39 ± 6.44
FM (HD)	kg	16.94 ± 3.75	18.54 ± 5.17	% WT	28.07 ± 5.21	29.51 ± 5.69
FM (TBK)	kg	16.43 ± 3.73	18.57 ± 5.45	% WT	27.32 ± 5.62	29.56 ± 6.31
FM (DEXA)	kg	17.18 ± 3.69	20.19 ± 5.18	% WT	28.45 ± 4.91	31.98 ± 5.40
FM-Siri (TBW, HD)	kg	16.79 ± 4.00	18.97 ± 5.45	% WT	27.80 ± 5.74	29.63 ± 5.79
FM-Fuller (TBW, HD, DEXA)	kg	17.08 ± 4.02	19.55 ± 5.18	% WT	28.28 ± 5.69	30.46 ± 5.20
High BMI Group		Pre-pregnancy n=12	24 wk postpartum n=8		Pre-pregnancy n=12	24 wk postpartum n=8
Army FM	kg	28.43 ± 6.73	25.48 ± 2.53	% WT	35.82 ± 4.36	32.76 ± 2.87
Navy FM	kg	30.68 ± 8.11	29.56 ± 4.52	% WT	38.50 ± 5.21	37.80 ± 3.54
Air Force FM	kg	33.53 ± 7.63	32.33 ± 5.02	% WT	42.22 ± 4.42	41.30 ± 3.65
FM (TBW)	kg	29.93 ± 6.25	27.67 ± 6.07	% WT	38.00 ± 5.72	35.83 ± 6.54
FM (HD)	kg	31.74 ± 6.46	30.38 ± 7.42	% WT	39.88 ± 3.69	38.66 ± 7.04
FM (TBK)	kg	31.48 ± 8.46	31.99 ± 7.14	% WT	39.73 ± 6.77	40.71 ± 6.52
FM (DEXA)	kg	34.15 ± 7.10	33.10 ± 7.69	% WT	42.76 ± 3.64	42.17 ± 7.06
FM-Siri (TBW, HD)	kg	30.81 ± 6.37	27.99 ± 5.30	% WT	38.81 ± 4.70	36.27 ± 5.36
FM-Fuller (TBW, HD, DEXA)	kg	31.71 ± 6.36	28.26 ± 5.65	% WT	39.59 ± 4.66	36.73 ± 5.57

*Mean±SD





Fat-Free Mass Compartment

In terms of work capacity and physical fitness, the fat-free mass compartment is the metabolically active part of the body. We monitored changes in FFM during pregnancy and the postpartum period. We will test whether physical activity can maintain a higher level of physical fitness throughout pregnancy and prevent an undesirable loss of muscle mass associated with postpartum weight loss. We will also be able to relate changes in FFM to any changes in strength.

Based on the principle that the K/N ratio for muscle is higher than that of nonmuscle tissues (3.03 vs. 1.33 meq/g), TBK and TBN will be used to estimate the relative amounts of protein in muscle and nonmuscle components of the body and the mass of each component prior to conception and postpartum. Our results indicate no postpartum accretion of TBK or TBN due to pregnancy.

Prior to conception, total body bone mineral density of all subjects was within the normal range (1.00 to 1.36). Our results do not indicate significant pregnancy-induced changes in bone mineral content, adjusted for bone area.

Table 12. Fat-free mass (FFM) compartment determined by total body potassium, DEXA, and prompt-gamma activation								
Low BMI Group		Pre-pregnancy	8 wk pregnancy	22 wk pregnancy	36 wk pregnancy	2 wk postpartum	6 wk postpartum	24 wk postpartum
		n=18	n=17	n=18	n=14	n=14	n=13	n=12
Total body potassium (TBK)								
TBK	g	93.35 ± 11.37*	91.70 ± 8.86	93.64 ± 9.67	100.75 ± 11.19	94.81 ± 12.52	93.21 ± 11.82	93.40 ± 11.05
K40 LBM	kg	39.68 ± 4.83	38.97 ± 3.77	39.80 ± 4.11	42.82 ± 4.75	40.29 ± 5.32	39.61 ± 5.02	39.69 ± 4.69
Dual energy x-ray absorptiometry (DEXA)								
Bone mineral content (BMC)	kg	2.053 ± 0.209	***	***	***	2.031 ± 0.224	2.020 ± 0.226	1.947 ± 0.207
Bone mineral density (BMD)	g/cm ²	1.111 ± 0.053	***	***	***	1.094 ± 0.062	1.098 ± 0.077	1.067 ± 0.048
FFM (DEXA)	kg	38.63 ± 4.50	***	***	***	39.38 ± 4.72	38.66 ± 4.43	38.45 ± 5.28
Prompt-gamma activation analysis (PGA)								
Total body nitrogen	g	1279.1 ± 137.0				1289.0 ± 165.7		1242.7 ± 180.7
Muscle mass	kg	13.51 ± 4.50				13.76 ± 4.37		13.65 ± 2.72
Nonmuscle mass	kg	24.27 ± 5.38				24.33 ± 5.54		23.14 ± 4.86
Normal BMI Group		Pre-pregnancy	8 wk pregnancy	22 wk pregnancy	36 wk pregnancy	2 wk postpartum	6 wk postpartum	24 wk postpartum
		n=33	n=31	n=33	n=30	n=33	n=35	n=34
Total body potassium (TBK)								
TBK	g	102.52 ± 13.26	99.92 ± 13.79	102.24 ± 13.77	108.32 ± 14.87	103.09 ± 11.75	100.97 ± 12.97	102.37 ± 13.60
K40 LBM	kg	43.57 ± 5.64	42.46 ± 5.86	43.45 ± 5.85	46.03 ± 6.32	43.81 ± 4.99	42.91 ± 5.51	43.51 ± 5.78
Dual energy x-ray absorptiometry (DEXA)								
Bone mineral content (BMC)	kg	2.298 ± 0.328	***	***	***	2.305 ± 0.334	2.258 ± 0.298	2.275 ± 0.353
Bone mineral density (BMD)	g/cm ²	1.159 ± 0.091	***	***	***	1.150 ± 0.089	1.142 ± 0.089	1.140 ± 0.098
FFM (DEXA)	kg	42.56 ± 4.91	***	***	***	43.60 ± 4.70	42.18 ± 5.20	42.34 ± 5.32

Table 12 Fat free mass (FFM) ...(cont)

Prompt-gamma activation analysis (PGA)								
High BMI Group		Pre-pregnancy n=10	8 wk pregnancy n=10	22 wk pregnancy n=12	36 wk pregnancy n=11	2 wk postpartum n=11	6 wk postpartum n=11	24 wk postpartum n=8
Total body nitrogen	g	1499.2 ± 193.6				1501.4 ± 210.7		1472.8 ± 236.9
Muscle mass	kg	12.37 ± 3.35				12.64 ± 2.95		13.19 ± 3.43
Nonmuscle mass	kg	31.34 ± 5.47				31.17 ± 6.55		29.92 ± 7.47
Dual energy x-ray absorptiometry (DEXA)								
TBK	g	106.77 ± 10.02	105.05 ± 12.17	109.89 ± 8.30	120.62 ± 9.41	108.63 ± 8.68	106.57 ± 11.11	107.89 ± 9.83
K40 LBM	kg	45.38 ± 4.26	44.65 ± 5.17	46.70 ± 3.53	51.27 ± 4.00	46.17 ± 3.69	45.29 ± 4.72	45.85 ± 4.18
Bone mineral content (BMC)	kg	2.330 ± 0.260	***	***	***	2.336 ± 0.273	2.331 ± 0.266	2.256 ± 0.262
Bone mineral density (BMD)	g/cm ²	1.131 ± 0.078	***	***	***	1.134 ± 0.074	1.123 ± 0.063	1.117 ± 0.078
FFM (DEXA)	kg	44.97 ± 4.93	***	***	***	47.40 ± 5.44	46.54 ± 5.01	44.25 ± 4.58
Prompt-gamma activation analysis (PGA)								
Total body nitrogen	g	1591.3 ± 249.9				1714.6 ± 159.2		1626.4 ± 204.2
Muscle mass	kg	11.93 ± 4.99				9.44 ± 4.96		11.75 ± 1.81
Nonmuscle mass	kg	34.26 ± 10.02				39.75 ± 7.52		35.38 ± 6.40

*Mean ±SD

Energy Expenditure and Substrate Utilization by Room Respiration Calorimetry

Preconceptional baseline levels of energy expenditure and substrate utilization are summarized in Tables 13 and 14. The mean 24-h TEE and heart rates of the women confined to the room calorimeter are representative of sedentary conditions. TEE increases in absolute terms and relative to body weight throughout pregnancy. By 6 mo postpartum TEE has returned to pre-pregnancy baseline. These data will be used to evaluate changes in energy metabolism and their impact on energy requirements throughout the reproductive cycle in women with low to high BMIs.

Table 13. Room respiration calorimetry

Low BMI Group		Pre-pregnancy n=18	8 wk pregnancy n=14	22 wk pregnancy n=16	36 wk pregnancy n=12	24 wk postpartum n=14
Total energy expenditure (TEE)						
Heart rate	bpm	66.77 ± 9.92*	69.97 ± 8.58	75.31 ± 9.13	77.49 ± 9.56	66.01 ± 10.81
RQ		0.872 ± 0.028	0.894 ± 0.029	0.893 ± 0.026	0.875 ± 0.028	0.887 ± 0.028
TEE	kcal/d	1649 ± 169	1660 ± 164	1760 ± 158	2021 ± 192	1662 ± 146
TEE	kcal/kg/d	33.14 ± 3.49	32.89 ± 2.52	31.01 ± 2.60	31.81 ± 2.64	32.49 ± 2.89
Basal metabolic rate (BMR)						
Heart rate	bpm	62.01 ± 10.21	65.27 ± 8.67	71.09 ± 10.38	70.19 ± 7.71	60.05 ± 11.40
RQ		0.811 ± 0.047	0.805 ± 0.049	0.838 ± 0.068	0.802 ± 0.056	0.804 ± 0.033
BMR	kcal/d	1206 ± 135	1252 ± 130	1329 ± 117	1534 ± 163	1219 ± 117
BMR	kcal/kg/d	24.20 ± 2.30	24.82 ± 2.27	23.32 ± 2.04	24.13 ± 2.26	23.81 ± 2.24
TEE/BMR		1.37 ± 0.07	1.33 ± 0.07	1.33 ± 0.04	1.32 ± 0.05	1.37 ± 0.08

Table 13. Room respiration calorimetry ...(cont)

Normal BMI						
		Pre-pregnancy n=33	8 wk pregnancy n=31	22 wk pregnancy n=34	36 wk pregnancy n=30	24 wk postpartum n=32
Sleeping metabolic rate (SMR)						
Heart rate	bpm	58.49 ± 9.33	63.56 ± 8.89	69.19 ± 9.46	70.82 ± 9.29	58.06 ± 10.77
RQ		0.835 ± 0.026	0.857 ± 0.036	0.859 ± 0.031	0.859 ± 0.028	0.849 ± 0.028
SMR	kcal/kg/d	23.83 ± 2.31	24.65 ± 1.60	23.12 ± 1.92	24.37 ± 2.05	23.88 ± 1.82
Total energy expenditure (TEE)						
Heart rate	bpm	65.40 ± 8.77	69.86 ± 9.07	75.47 ± 8.66	80.09 ± 10.50	65.40 ± 8.69
RQ		0.876 ± 0.016	0.884 ± 0.024	0.885 ± 0.022	0.873 ± 0.026	0.876 ± 0.027
TEE	kcal/d	1777 ± 181	1794 ± 197	1872 ± 189	2184 ± 236	1803 ± 190
TEE	kcal/kg/d	29.19 ± 2.61	29.56 ± 2.27	28.73 ± 2.20	30.58 ± 2.97	28.94 ± 3.10
Basal metabolic rate (BMR)						
Heart rate	bpm	60.86 ± 8.57	66.05 ± 9.13	70.74 ± 9.34	74.31 ± 11.76	59.81 ± 9.06
RQ		0.805 ± 0.034	0.827 ± 0.051	0.811 ± 0.053	0.824 ± 0.049	0.815 ± 0.042
BMR	kcal/d	1322 ± 126	1353 ± 156	1418 ± 142	1673 ± 172	1325 ± 135
BMR	kcal/kg/d	21.73 ± 2.01	22.32 ± 2.24	21.77 ± 1.81	23.55 ± 2.26	21.29 ± 2.27
TEE/BMR		1.35 ± 0.07	1.33 ± 0.08	1.32 ± 0.08	1.30 ± 0.06	1.36 ± 0.07
Sleeping metabolic rate (SMR)						
Heart rate	bpm	56.02 ± 13.01	62.44 ± 9.03	69.10 ± 8.54	73.58 ± 11.17	57.84 ± 12.19
RQ		0.834 ± 0.022	0.844 ± 0.032	0.852 ± 0.031	0.845 ± 0.032	0.843 ± 0.031
SMR	kcal/kg/d	21.13 ± 1.94	21.67 ± 1.96	21.38 ± 1.94	23.45 ± 2.33	21.58 ± 2.07
High BMI Group						
		Pre-pregnancy n=10	8 wk pregnancy n=8	22 wk pregnancy n=11	36 wk pregnancy n=10	24 wk postpartum n=7
Total energy expenditure (TEE)						
Heart rate	bpm	73.41 ± 6.51	78.03 ± 5.96	85.79 ± 6.13	90.15 ± 5.23	74.74 ± 6.63
RQ		0.867 ± 0.018	0.880 ± 0.023	0.869 ± 0.019	0.863 ± 0.027	0.869 ± 0.034
TEE	kcal/d	2094 ± 221	2169 ± 261	2267 ± 276	2620 ± 343	1988 ± 190
TEE	kcal/kg/d	26.89 ± 1.79	26.10 ± 1.26	26.60 ± 1.82	27.66 ± 1.90	25.73 ± 1.96
Basal metabolic rate (BMR)						
Heart rate	bpm	65.77 ± 5.99	71.40 ± 6.27	79.33 ± 7.98	81.59 ± 10.25	66.61 ± 6.93
RQ		0.790 ± 0.052	0.822 ± 0.031	0.809 ± 0.052	0.796 ± 0.042	0.778 ± 0.089
BMR	kcal/d	1505 ± 153	1600 ± 213	1693 ± 210	2016 ± 254	1459 ± 122
BMR	kcal/kg/d	19.32 ± 1.12	19.22 ± 0.71	19.78 ± 1.67	21.54 ± 2.29	18.98 ± 2.36
TEE/BMR		1.39 ± 0.05	1.36 ± 0.06	1.36 ± 0.07	1.32 ± 0.05	1.37 ± 0.12
Sleeping metabolic rate (SMR)						
Heart rate	bpm	64.70 ± 4.74	68.18 ± 9.74	77.74 ± 5.23	83.24 ± 4.05	67.45 ± 6.81
RQ		0.823 ± 0.022	0.845 ± 0.030	0.833 ± 0.023	0.832 ± 0.031	0.826 ± 0.042
SMR	kcal/m	1.02 ± 0.12	1.10 ± 0.14	1.13 ± 0.15	1.40 ± 0.18	1.00 ± 0.09
SMR	kcal/kg/d	18.94 ± 1.40	19.11 ± 1.16	19.03 ± 1.58	21.25 ± 1.59	18.74 ± 1.75

*Mean ± SD

Table 14. Substrate utilization estimated from 24-h respiration calorimetry

Low BMI Group		Pre-pregnancy n=17	8 wk pregnancy n=14	22 wk pregnancy n=15	36 wk pregnancy n=8	24 wk postpartum n=8
Protein utilization	g/d	61.1 ± 12.7	71.1 ± 11.6	66.6 ± 11.1	66.1 ± 19.9	83.1 ± 24.9
Carbohydrate utilization	g/d	195.3 ± 37.8	219.2 ± 34.8	233.6 ± 29.5	240.3 ± 30.5	208.4 ± 17.9
Fat utilization	g/d	55.8 ± 20.6	41.4 ± 20.4	47.8 ± 20.4	77.7 ± 18.1	41.3 ± 10.1
Protein utilization	% TEE	17.7 ± 3.8	20.4 ± 3.1	18.0 ± 3.1	15.0 ± 3.6	23.4 ± 5.3
Carbohydrate utilization	% TEE	50.3 ± 9.6	56.2 ± 9.8	56.5 ± 9.1	49.4 ± 8.1	53.1 ± 6.4
Fat utilization	% TEE	31.9 ± 10.5	23.3 ± 10.4	25.3 ± 9.3	35.5 ± 6.5	23.4 ± 4.6
Carbohydrate utilization	%NPEE	61.4 ± 12.3	70.8 ± 12.7	69.1 ± 11.3	58.0 ± 7.9	69.3 ± 6.3
Fat utilization	%NPEE	38.6 ± 12.3	29.2 ± 12.7	30.9 ± 11.3	42.0 ± 7.9	30.7 ± 6.3
Normal BMI Group		Pre-pregnancy n=33	8 wk pregnancy n=32	22 wk pregnancy n=33	36 wk pregnancy n=28	24 wk postpartum n=28
Protein utilization	g/d	62.7 ± 10.7	73.5 ± 17.8	68.6 ± 17.0	72.7 ± 19.3	72.2 ± 14.0
Carbohydrate utilization	g/d	216.6 ± 29.6	225.7 ± 36.9	241.1 ± 36.4	264.8 ± 51.0	214.9 ± 46.9
Fat utilization	g/d	59.0 ± 14.6	51.0 ± 16.3	55.4 ± 18.1	77.8 ± 23.9	58.5 ± 19.7
Protein utilization	% TEE	16.8 ± 2.7	19.5 ± 3.7	17.3 ± 3.5	15.6 ± 3.6	19.0 ± 3.8
Carbohydrate utilization	% TEE	51.6 ± 5.8	53.3 ± 7.9	54.5 ± 7.7	50.8 ± 9.1	50.2 ± 9.4
Fat utilization	% TEE	31.5 ± 5.8	27.1 ± 8.4	28.0 ± 7.9	33.5 ± 9.2	30.6 ± 8.6
Carbohydrate utilization	%NPEE	62.1 ± 6.8	66.3 ± 10.1	66.1 ± 9.3	60.3 ± 10.6	62.0 ± 10.6
Fat utilization	%NPEE	37.9 ± 6.8	33.7 ± 10.1	33.9 ± 9.3	39.7 ± 10.6	38.0 ± 10.6
High BMI Group		Pre-pregnancy n=10	8 wk pregnancy n=7	22 wk pregnancy n=7	36 wk pregnancy n=6	24 wk postpartum n=3
Protein utilization	g/d	71.9 ± 12.6	90.3 ± 21.7	76.1 ± 11.6	75.8 ± 12.5	62.1 ± 19.6
Carbohydrate utilization	g/d	240.9 ± 34.4	256.2 ± 37.5	245.9 ± 41.9	296.6 ± 85.9	230.4 ± 47.1
Fat utilization	g/d	76.8 ± 19.2	71.5 ± 26.7	75.4 ± 8.8	89.2 ± 17.2	67.1 ± 39.7
Protein utilization	% TEE	16.5 ± 3.4	20.0 ± 6.2	17.1 ± 2.3	14.8 ± 3.0	15.3 ± 3.6
Carbohydrate utilization	% TEE	48.7 ± 6.0	49.4 ± 5.4	48.7 ± 4.7	50.1 ± 8.8	52.2 ± 14.9
Fat utilization	% TEE	34.6 ± 6.7	30.5 ± 7.8	34.1 ± 3.7	34.9 ± 7.5	32.4 ± 15.1
Carbohydrate utilization	%NPEE	58.5 ± 7.4	62.1 ± 7.4	58.8 ± 4.8	58.8 ± 9.4	61.7 ± 17.6
Fat utilization	%NPEE	41.5 ± 7.4	37.9 ± 7.4	41.2 ± 4.8	41.2 ± 9.4	38.3 ± 17.6

*Mean±SD

Abbreviations: EI=energy intake; RQ=respiratory quotient; TEE=total energy expenditure; NPEE=nonprotein energy expenditure.

Total Energy Expenditure by Doubly-Labeled Water Method

Free-living energy expenditure (TEE) averaged 2353 ± 269 kcal/d or 47 ± 6 kcal·kg⁻¹·d⁻¹ in the low BMI group, 2423 ± 369 kcal/d or 41 ± 6 kcal·kg⁻¹·d⁻¹ in the normal BMI group, and 2940 ± 421 kcal/d or 38 ± 4 kcal·kg⁻¹·d⁻¹ in the high BMI group. TEE, plus the estimated energy deposited in tissues, will be used to estimate dietary energy needs of pregnant women.

The mean physical activity levels (PAL=TEE/BMR) of 1.92 ± 0.25 , 1.84 ± 0.25 , and 1.96 ± 0.22 for the low, normal and high BMI groups, respectively, would be classified as heavy (1.82), according to FAO/WHO/UNU (4); however, there is considerable individual variation. The PAL values at 22 and 36 wk of pregnancy indicate a definite trend for physical activity to decrease progressively with advancing gestation in all three BMI groups.

Measurements of TEE, AEE, and PAL will allow us to assess changes in activity during pregnancy and the postpartum period. In conjunction with the fitness and strength tests, we can assess whether pregnancy-induced changes in activity influence the ability to return to military duty.

Low BMI Group	Pre-pregnancy n=18	22 wk pregnancy n=16	36 wk pregnancy n=10	24 wk postpartum n=8
^2H dilution space (kg)	29.91 ± 2.97*	33.60 ± 3.68	36.15 ± 4.23	29.30 ± 3.02
^{18}O dilution space (kg)	29.08 ± 3.00	32.67 ± 3.72	35.16 ± 3.92	28.31 ± 2.91
$^2\text{H}/^{18}\text{O}$	1.03 ± 0.02	1.03 ± 0.02	1.03 ± 0.01	1.03 ± 0.02
RCO_2 (mol/d)	18.46 ± 2.11	18.07 ± 3.01	19.59 ± 4.08	16.48 ± 2.34
RO_2 (mol/d)	21.47 ± 2.46	21.01 ± 3.50	22.78 ± 4.74	19.16 ± 2.73
EE (kcal/d)	2353 ± 269	2304 ± 384	2498 ± 520	2101 ± 299
EE (kcal/kg/d)	47.45 ± 6.23	40.60 ± 7.06	40.01 ± 9.21	41.67 ± 8.37
TEE/BMR	1.97 ± 0.25	1.74 ± 0.29	1.68 ± 0.37	1.72 ± 0.30
Normal BMI Group	Pre-pregnancy n=33	22 wk pregnancy n=31	36 wk pregnancy n=32	24 wk postpartum n=27
^2H dilution space (kg)	33.17 ± 3.99*	35.39 ± 3.95	40.56 ± 3.66	33.56 ± 3.88
^{18}O dilution space (kg)	31.92 ± 3.99	34.54 ± 3.88	39.62 ± 3.67	32.72 ± 3.83
$^2\text{H}/^{18}\text{O}$	1.04 ± 0.02	1.02 ± 0.02	1.02 ± 0.03	1.03 ± 0.02
RCO_2 (mol/d)	19.01 ± 2.89	19.77 ± 2.94	21.38 ± 2.76	19.67 ± 3.13
RO_2 (mol/d)	22.10 ± 3.36	22.99 ± 3.42	24.86 ± 3.21	22.88 ± 3.63
EE (kcal/d)	2423 ± 369	2521 ± 374	2726 ± 352	2508 ± 399
EE (kcal/kg/d)	40.37 ± 5.86	38.68 ± 5.86	37.80 ± 5.32	40.03 ± 5.65
TEE/BMR	1.84 ± 0.25	1.77 ± 0.28	1.63 ± 0.24	1.89 ± 0.25
High BMI Group	Pre-pregnancy n=10	22 wk pregnancy n=9	36 wk pregnancy n=7	24 wk postpartum n=6
^2H dilution space (kg)	36.92 ± 5.41*	40.18 ± 4.20	43.37 ± 3.69	37.32 ± 4.51
^{18}O dilution space (kg)	35.92 ± 5.07	38.88 ± 3.79	42.37 ± 3.57	36.58 ± 4.75
$^2\text{H}/^{18}\text{O}$	1.03 ± 0.02	1.03 ± 0.01	1.02 ± 0.02	1.02 ± 0.02
RCO_2 (mol/d)	23.07 ± 3.30	22.99 ± 3.80	22.86 ± 4.75	21.06 ± 3.24
RO_2 (mol/d)	26.82 ± 3.84	26.73 ± 4.41	26.58 ± 5.53	24.49 ± 3.76
EE (kcal/d)	2940 ± 421	2931 ± 484	2914 ± 606	2685 ± 413
EE (kcal/kg/d)	37.71 ± 4.40	34.70 ± 7.07	32.12 ± 4.68	35.34 ± 5.18
TEE/BMR	1.96 ± 0.22	1.73 ± 0.25	1.51 ± 0.26	1.77 ± 0.19

*Mean±SD

Fitness and Strength

The measurement of VO_2max allowed us to categorize these women in terms of pre-pregnancy fitness. The mean VO_2max of $40\text{ml}/\text{kg}^{-1}\text{min}^{-1}$ for the low BMI group, and $34\text{ml}/\text{kg}^{-1}\text{min}^{-1}$ for the normal BMI group would be in the "average" range of fitness, defined as $34\text{--}38\text{ml}/\text{kg}^{-1}\text{min}^{-1}$ for women. The mean VO_2max for the high BMI group of $26\text{ml}/\text{kg}^{-1}\text{min}^{-1}$ would be categorized in the "low" range of fitness. There was considerable variability in the level of fitness in all BMI groups. The maximal heart rates and the high RQ (1.2) also demonstrate that a true maximum was achieved. The maximal workload averaged 160, 170, and 148 watts, in the 3 respective groups, indicating a fairly high level of work reached at exhaustion.

For the one-repetition maximum strength testing, it appears that these women are fairly strong prior to pregnancy. The women can bench press about 72, 76, and 79 lbs of their body weight, in the 3 respective groups. For the lower body, the leg extension measurement indicates a high degree of strength (80, 90, 96 lbs). As with the fitness levels, there seems to be a great deal of variability in the group for strength. About half the women report that they strength train regularly.

Changes at 6 and 24 wk postpartum from pre-pregnancy measurements were examined. VO_2 max/kg was lower 6 wk postpartum and almost recovered to pre-pregnancy baseline at 24 wk postpartum (37, 29, 22 ml/kg⁻¹min⁻¹), in the 3 respective BMI groups. Strength measures also decreased for the leg press, leg extension and bench press at 6 wk postpartum, and showed improvement by 6 mo postpartum.

Table 16. Physical fitness and strength

Low BMI Group		Pre-pregnancy n=18	6 wk postpartum n=14	24 wk postpartum n=12
Steady state exercise on stationary cycle ergometer				
Workload	watts	50 ± 0*	50 ± 0	50 ± 0
Heart rate	bpm	122 ± 19	124 ± 23	122 ± 23
VO ₂	lpm	0.88 ± 0.08	0.86 ± 0.05	0.89 ± 0.07
VO ₂ /kg	ml/kg/m	17.80 ± 2.17	15.96 ± 1.74	17.40 ± 2.32
RQ		0.91 ± 0.06	1.04 ± 0.08	1.00 ± 0.09
Maximal exercise on stationary cycle ergometer				
Workload	watts	160 ± 41	128 ± 28	150 ± 48
Heart rate	bpm	180 ± 10	171 ± 13	174 ± 13
VO ₂	lpm	1.97 ± 0.42	1.58 ± 0.36	1.89 ± 0.49
VO ₂ /kg	ml/kg/m	39.68 ± 8.15	29.27 ± 6.67	36.82 ± 8.78
RQ		1.21 ± 0.06	1.26 ± 0.10	1.23 ± 0.09
Strength testing: 1-repetition maximum				
Leg press	lb	93 ± 29	75 ± 31	88 ± 40
Leg extension	lb	80 ± 19	77 ± 13	82 ± 19
Bench press	lb	72 ± 19	71 ± 20	69 ± 17
Latissimus pull-down	lb	62 ± 12	54 ± 9	57 ± 8
Submaximal exercise on treadmill @ 2.5 mph				
Heart rate	bpm	91 ± 14	104 ± 11	91 ± 16
VO ₂	lpm	0.58 ± 0.07	0.70 ± 0.06	0.58 ± 0.06
RQ		0.88 ± 0.04	0.87 ± 0.03	0.91 ± 0.04
Energy expenditure	kcal/m	2.88 ± 0.35	3.46 ± 0.32	2.89 ± 0.30

Table 16. Physical fitness and strength ...(cont)

Normal BMI Group		Pre-pregnancy n=33	6 wk postpartum n=34	24 wk postpartum n=34
Steady state exercise on stationary cycle				
Workload	watts	50 ± 0*	50 ± 0	50 ± 0
Heart rate	bpm	119 ± 17	121 ± 19	118 ± 21
VO ₂	lpm	0.91 ± 0.06	0.91 ± 0.09	0.89 ± 0.06
VO ₂ /kg	ml/kg/m	15.23 ± 1.44	14.46 ± 2.19	14.59 ± 1.80
RQ		0.93 ± 0.07	1.03 ± 0.08	1.01 ± 0.09
Maximal exercise on stationary cycle ergometer				
Workload	watts	170 ± 34	141 ± 32	150 ± 36
Heart rate	bpm	177 ± 9	172 ± 9	174 ± 12
VO ₂	lpm	2.08 ± 0.43	1.74 ± 0.34	1.82 ± 0.39
VO ₂ /kg	ml/kg/m	34.47 ± 6.20	27.43 ± 5.20	29.47 ± 5.60
RQ		1.22 ± 0.09	1.28 ± 0.09	1.28 ± 0.10
Strength testing: 1-repetition				
Leg press	lb	79 ± 34	67 ± 29	82 ± 29
Leg extension	lb	90 ± 22	86 ± 21	95 ± 20
Bench press	lb	76 ± 16	71 ± 16	74 ± 17
Latissimus pull-down	lb	63 ± 11	57 ± 13	63 ± 10
Submaximal exercise on treadmill @ 2.5 mph				
Heart rate	bpm	95 ± 13	105 ± 13	93 ± 15
VO ₂	lpm	0.68 ± 0.10	0.78 ± 0.14	0.66 ± 0.10
RQ		0.90 ± 0.03	0.88 ± 0.04	0.89 ± 0.03
Energy expenditure	kcal/m	3.37 ± 0.50	3.82 ± 0.66	3.27 ± 0.52
High BMI Group		Pre-pregnancy n=10	6 wk postpartum n=10	24 wk postpartum n=8
Steady state exercise on stationary cycle				
Workload	watts	50 ± 0*	50 ± 0	50 ± 0
Heart rate	bpm	123 ± 16	127 ± 16	140 ± 17
VO ₂	lpm	1.00 ± 0.07	1.01 ± 0.07	1.04 ± 0.14
VO ₂ /kg	ml/kg/m	12.86 ± 1.68	11.80 ± 0.72	13.36 ± 1.70
RQ		0.98 ± 0.05	1.05 ± 0.07	1.09 ± 0.08
Maximal exercise on stationary cycle ergometer				
Workload	watts	148 ± 27	130 ± 24	121 ± 25
Heart rate	bpm	179 ± 12	168 ± 16	175 ± 8
VO ₂	lpm	2.04 ± 0.30	1.76 ± 0.23	1.69 ± 0.27
VO ₂ /kg	ml/kg/m	26.14 ± 4.24	20.73 ± 3.14	21.74 ± 3.39
RQ		1.25 ± 0.08	1.26 ± 0.11	1.29 ± 0.09
Strength testing: 1-repetition				
Leg press	lb	60 ± 27	43 ± 34	70 ± 37
Leg extension	lb	96 ± 14	89 ± 21	105 ± 23
Bench press	lb	79 ± 14	79 ± 19	81 ± 24
Latissimus pull-down	lb	66 ± 7	62 ± 4	66 ± 7

Table 16. Physical fitness and strength ...(cont)

Submaximal exercise on treadmill @ 2.5 mph				
Heart rate	bpm	105 ± 12	125 ± 9	108 ± 13
VO ₂	lpm	0.88 ± 0.13	1.04 ± 0.14	0.84 ± 0.09
RQ		0.91 ± 0.03	0.91 ± 0.02	0.90 ± 0.04
Energy expenditure	kcal/m	4.35 ± 0.66	5.14 ± 0.70	4.15 ± 0.45

*Mean±SD

8. CONCLUSIONS

This study addresses the impact of pregnancy-induced changes in body composition and physical fitness on postpartum return to duty readiness. Prior to conception, the subjects represent healthy, moderately-active women. Based on military standards, nearly all subjects in the low and normal BMI groups would be eligible for the Armed Services. The weight, body fat, bone density and iron status of these women are within normal limits. Energy expenditure measurements indicate that these women are physically active. VO₂max and strength tests confirm that the women are fit and fairly strong. Our evaluation of military equations for the prediction of fat mass based on 117 women prior to pregnancy strongly indicates a need to revise the equations. The Army and Navy equations underestimate body fat. Although conservative, the equations do not perform equally across the range of body fat mass. The underestimation is exaggerated at higher levels of fat mass. Our data indicate a mean gestational weight gain of 15.5 kg. Postpartum weight retention at 6 mo averaged 2.7 ± 3.9 kg, and was greater in the high BMI group (4.0 ± 4.4 kg) than the low and normal BMI groups (2.4 ± 3.5 kg). BMI had increased by approximately one unit in all groups. FM derived from the Fuller-4 component model indicated an increase of 2.1 ± 4.1 kg FM at 6 mo postpartum, with means of 2.6 ± 3.7, 1.7 ± 4.1 and 3.0 ± 5.0 kg in the low, normal and high BMI groups, respectively. There was evidence of deconditioning at 6 wk postpartum with near recovery at 6 mo postpartum. The effect of gestational weight gain, fat accretion and deconditioning would affect a woman's ability to return to weight, body fat and fitness standards in the 6 mo following delivery.

9. REFERENCES

1. U.S. Department of Army. 1986. Army Regulation 600-9, "The Army Weight Control Program." September 1, Washington D.C.
2. Vogel JA, Kirkpatrick JW, Fitzgerald PI, Hodgdon JA, Harman EA. Derivation of anthropometry based body fat equations for the Army's weight control program. 1988. Technical Report No. T17-88, U.S. Army Research Institute of Environmental Medicine, Natick, MA.
3. Hodgdon JA, Beckett MB. Prediction of percent body fat for U.S. Navy women from body circumference and height. 1984. Report No. 84-29. Naval Health Research Center, San Diego, CA.
4. Wright HF, Wilmore JH. Estimation of relative body fat and lean body weight in a United States Marine Corps population. 1974. Aerospace Med 45: 301-6.

5. Brennan EH. Development of a binomial involving anthropometric measurements for predicting lean mass in young women. 1974. M.S. thesis. Incarnate Word College, San Antonio, TX.
6. U.S. Department of Army. 1983. Army Regulation 40-501, Change 34, "Standards of Medical Fitness." December 1, Washington, D.C.
7. Institute of Medicine, Food and Nutrition Board, Committee on Military Nutrition Research. 1992. In: Mariott BM, Grumstrup-Scott J. (ed), Body composition and physical performance. National Academy Press.
8. Laurence MT. Enlistment height/weight standards and attrition from the military. Defense Manpower Data Center, Arlington, VA.
9. Institute of Medicine, Food and Nutrition Board, Committee on Nutritional Status During Pregnancy and Lactation. Nutrition during pregnancy. Part I. Weight gain. Part II. Nutrient supplements. 1990. Washington, DC: National Academy Press.
10. Naeye RL. Weight gain and the outcome of pregnancy. 1979. *Am J Obstet Gynecol* 135:3-9.
11. Taffel SM. Maternal weight gain and the outcome of pregnancy: US, 1980. 1986. *Vital and Health Statistics, Series 21, No. 44.* DHHS Publ. No. (PHS) 86-1922.
12. Langhoff-Roos J, Lindmark G, Gebre-Medhin M. Maternal fat stores and fat accretion during pregnancy in relation to infant birthweight. 1987. *Brit J Obs Gyn* 94:1170-7.
13. Lawrence M, McKillop FM, Durnin JVGA. Women who gain more fat during pregnancy may not have bigger babies: implications for recommended weight gain during pregnancy. 1991. *Brit J Obs Gyn* 98:254-9.
14. Hytten FE. Weight gain in pregnancy. 1991. In: *Clinical physiology in obstetrics.* Hytten FE, Chamberlain G, eds. Blackwell Scientific Publications. Oxford.
15. Hytten FE, Thomson AM, Taggert N. Total body water in normal pregnancy. 1966. *J Obstet Gynaecol Br Commonw* 73:553-61.
16. Fitch WL, King JC. Protein turnover and 3-methylhistidine excretion in non-pregnant, pregnant and gestational diabetic women. 1987. *Hum Nutr Clin Nutr* 41C:327-39.
17. Naismith DJ. Maternal nutrition and the outcome of pregnancy - a critical appraisal. 1980. *Proc Nutr Soc* 39:1-11.
18. Sohlström A, Wahlund L-O, Forsum E. Total body fat and its distribution during human reproduction as assessed by magnetic resonance imaging. 1993. In: Ellis K (ed), *Human Body Composition* pp 181-184.

19. Dewey KG, Heinig MJ, Nommsen LA. Maternal weight-loss patterns during prolonged lactation. 1993. *Am J Clin Nutr* 58:162-166.
20. Brewer MM, Bates MR, Vannoy LP. Postpartum changes in maternal weight and body fat depots in lactating vs non-lactating women. 1989. *Am J Clin Nutr* 49:259-265.
21. Taggart NR, Holliday RM, Billewics WZ, Hytten FE, Thomson AM. Changes in skinfolds during pregnancy. 1967. *Br J Nutr* 21:439-51.
22. Goldberg GR, Prentice AM, Coward WA, Davies HL, Murgatroyd PR, Wensing C, Black AE, Harding M, Sawyer M. Longitudinal assessment of energy expenditure in pregnancy by the doubly labelled water method. 1993. *Am J Clin Nutr* 57:494-505.
23. van Raaij JMA, Peek MEM, Vermaat-Miedema SH, Schonk CM, Hautvast JGAJ. New equations for estimating body fat mass in pregnancy from body density or total body water. 1988. *Am J Clin Nutr* 48:24-9.
24. Pipe NGJ, Smith T, Halliday D, Edmonds CJ, Williams C, Coltart TM. Changes in fat, fat-free mass and body water in human normal pregnancy. 1979. *Br J Obstet Gynaecol* 86:929-40.
25. Forsum E, Sadurskis A, Wager J. Resting metabolic rate and body composition of healthy Swedish women during pregnancy. 1988. *Am J Clin Nutr* 47:942-47.
26. King JC, Butte NF, Bronstein MN, Kopp LE, Lindquist SA. Energy metabolism during pregnancy: influence of maternal energy status. 1994. *Am J Clin Nutr* 59:439S-45S.
27. Johnson EM. Weight changes during pregnancy and the postpartum period. 1991. *Progress in Food and Nutr Sci* 15:117-57.
28. Keppel K, Taffel S. Pregnancy-related weight gain and retention: implications of the 1990 Institute of Medicine Guidelines. 1993. *Am J Publ Health* 83:1100-3.
29. Butte NF, Garza C, Stuff JE, Smith EO, Nichols BL. Effect of maternal diet and body composition on lactational performance. 1984. *Am J Clin Nutr* 39:296-306.
30. Manning-Dalton C, Allen LH. The effects of lactation on energy and protein consumption, postpartum weight change and body composition of well nourished North American women. 1983. *Nutr Res* 3:293-308.
31. Strode MA, Dewey KG, Lonnerdal B. Effect of short-term maternal caloric restriction on lactation performance of well-nourished mothers. 1986. *Acta Paediatr Scand* 75:222-9.
32. Forbes GB. *Human Body Composition*. 1987. Springer-Verlag, New York.

33. Ballor DL, Poehlman. Exercise-training enhances fat-free mass preservation during diet-induced weight loss: a meta-analytical finding. 1994. *International J Obesity* 18:35-40.
34. McMurray RG, Mottola MF, Wolfe LA, Artal R, Millar L, Pivarnik JM. Recent advances in understanding maternal and fetal responses to exercise. 1993. *Med Sci Sports Exerc* 25:1305-21.
35. Sady SP, Carpenter MW, Sady MA, Haydon B, Hoegsberg B, Cullinane EM, Thompson PD, Coustan DR. Prediction of Vo_{2max} during cycle exercise in pregnant women. 1988. *J Appl Physiol* 65:657-61.
36. Lotgering FK, van Doorn MB, Struijk PC, Pool J, Wallenburg HCS. Maximal aerobic exercise in pregnant women: heart rate, O_2 consumption, CO_2 production, and ventilation. 1991. *J Appl Physiol* 70:1016-23.
37. South-Paul JE, Rajagopal KR, Tenholder MF. Exercise responses prior to pregnancy and in the postpartum state. 1992. *Med Sci Sports Exerc* 24:410-4.
38. American College of Obstetricians and Gynecologists. Exercise during pregnancy and the postnatal period. 1985. ACOG Home Exercise Programs. Washington, DC: ACOG.
39. Viteri FE. Influence of iron nutrition on work capacity and performance. 1989. In: *Dietary Iron: Birth to Two Years*, ed LF Filer, Jr. Raven Press, Ltd., New York, 141-60.
40. Lukaski HC, Ball CB, Siders WA. Altered metabolic response of iron-deficient women during graded, maximal exercise. 1991. *Eur J Appl Physiol* 63:140-5.
41. FAO/WHO/UNU Expert Consultation. Energy and protein requirements. 1985. Technical Report Series 724, WHO, Geneva.
42. National Research Council. Recommended dietary allowances. 10th ed. Washington, DC: National Academy Press, 1989.
43. Committee on Medical Aspects of Food Policy. Dietary reference values for food energy and nutrients for the United Kingdom. 1991. Dept Hlth Rep Hlth Soc Subj 41. London: HMSO.
44. Durnin JVGA, McKillop FM, Grant S, Fitzgerald G. 1987. Energy requirements of pregnancy in Scotland. *Lancet* ii:897-900.
45. van Raaij JMA, Schonk CM, Vermaat-Miedema SH, Peek MEM, Hautvast JGAJ. Body fat mass and basal metabolic rate in Dutch women before, during, and after pregnancy: a reappraisal of energy cost of pregnancy. 1989. *Am J Clin Nutr* 49:765-72.

46. Prentice AM, Goldberg GR, Davies HL, Murgatroyd PR, Scott W. Energy-sparing adaptations in human pregnancy assessed by whole-body calorimetry. 1989. *Br J Nutr* 62:5-22.
47. de Groot LCPGM, Boekholt HA, Spaaij CJK, van Raaij JMA, Drijvers JJMM, van der Heijden LJM, van der Heide D, Hautvast JGAJ. Energy balances of healthy Dutch women before and during pregnancy: limited scope for metabolic adaptations in pregnancy. 1994. *Am J Clin Nutr* 59:827-32.
48. Forsum E, Kabir N, Sadurskis A, Westerterp K. Total energy expenditure of healthy Swedish women during pregnancy and lactation. 1992. *Am J Clin Nutr* 56:334-42.
49. King N, Fridlund KE, Askew EW. Nutrition issues of military women. 1993. *J Amer Coll Nutr* 12:344-8.
50. Friedl KE, Marchitelli LJ, Sherman DE, Tulley R. Nutritional Assessment of Cadets at the US Military Academy: Part 1. Anthropometric and Biomedical Measures." Natick MA: US Army Research Institute of Environmental Medicine. 1990. USARIEM Technical Report T4-91.
51. U.S. Department of the Army, Navy and the Air Force. Army Regulation 40-25/Naval Command Medical Instruction 10110.1/Air Force Regulation 160-95. Nutrient Allowances, Standards, and Education. May 15, 1985, Washington, DC.
52. Akers R, Buskirk ER. An underwater weighing system utilizing "force cube" transducers. 1969. *J Appl Physiol* 26:649-52.
53. Wilmore JH, Vodak PA, Parr RB, Girandola RN, Billing JE. Further simplification of a method for determination of residual lung volume. 1980. *Med Sci Sports Exercise* 12:216-8.
54. Burkinshaw L, Hill GL, Morgan DB. Mass and composition of the fat-free tissues of patients with weight loss. 1979. *Clin Sci* 68:455-62.
55. Fuller NJ, Jebb SA, Laskey MA, Coward WA, Elia M. Four-compartment model for the assessment of body composition in humans: comparison with alternative methods, and evaluation of the density and hydration of fat-free mass. 1992. *Clinical Science* 82:687-93.
56. International Dietary Energy Consulting Group. The doubly-labeled water method for measuring energy expenditure: Technical Recommendations for use in humans. 1990. (Prentice AM, ed) International Atomic Energy Agency, Vienna, Austria.

57. Black AE, Prentice AM, Coward WA. Use of food quotients to predict respiratory quotients for the doubly labeled water method for measuring energy expenditure. 1986. *Hum Nutr Clin Nutr* 40C:381-91.
58. de V Weir JB. New methods for calculating metabolic rate with special reference to protein metabolism. 1949. *J Physiol* 109:1-9.
59. Moon JK, Vohra FA, Valerio-Jimenez OS, Butte NF. Closed-loop control of carbon dioxide concentration and pressure improves response of room respiration calorimetry. 1995. *J Nutr* 125:220-8.
60. Livesey G, Elia M. Estimation of energy expenditure, net carbohydrate utilization, and net fat oxidation and synthesis by indirect calorimetry: evaluation of errors with special reference to the detailed composition of fuels. 1988. *Amer J Clin Nutr* 47:608-25.
61. Taylor HL, Jacobs DR, Schucker B, Knudsen J, Leon AS, Bebacker G. A questionnaire for the assessment of leisure time physical activities. 1978. *J Chron Dis* 31:741-55.
62. Shapiro S, Weinblatt E, Frank CW, Sagen V. The H.I.P. study of incidence and prognosis of coronary heart disease. 1965. *J Chron Dis* 18:527-58.
63. Dixon WJ (ed) 1983 BMDP, Biomedical computer programs. University of California Press, Berkeley, CA.
64. Minitab version 6.1 1988 Minitab, Inc., State College, PA.
65. Katch FI, McArdle WD (ed) 1983 Nutrition, weight control and exercise. Philadelphia, Lea & Febiger, p291.

10. BIBLIOGRAPHY OF PUBLICATIONS

Because of the longitudinal nature of this study, data have not been submitted for publication. Preliminary data were publically presented at a March of Dimes Scientific Conference on Nutrition and Optimal Human Development on June 6-9, 1999.

11. LIST OF PERSONNEL

Nancy Butte
Judy Hopkinson
Carolyn Heinz
Nitesh Mehta
Roman Shypailo



DEPARTMENT OF THE ARMY

US ARMY MEDICAL RESEARCH AND MATERIEL COMMAND
504 SCOTT STREET
FORT DETRICK, MARYLAND 21702-5012

REPLY TO
ATTENTION OF:

MCMR-RMI-S (70-1y)

23 Aug 01

MEMORANDUM FOR Administrator, Defense Technical Information
Center (DTIC-OCA), 8725 John J. Kingman Road, Fort Belvoir,
VA 22060-6218


SUBJECT: Request Change in Distribution Statement

1. The U.S. Army Medical Research and Materiel Command has reexamined the need for the limitation assigned to the technical reports listed at enclosure. Request the limited distribution statement for these reports be changed to "Approved for public release; distribution unlimited." These reports should be released to the National Technical Information Service.

2. Point of contact for this request is Ms. Judy Pawlus at DSN 343-7322 or by e-mail at judy.pawlus@det.amedd.army.mil.

FOR THE COMMANDER:

Encl


PHYLIS M. RINEHART
Deputy Chief of Staff for
Information Management

Reports to be Downgraded to Unlimited Distribution

ADB241560	ADB253628	ADB249654	ADB263448
ADB251657	ADB257757	ADB264967	ADB245021
ADB263525	ADB264736	ADB247697	ADB264544
ADB222448	ADB255427	ADB263453	ADB254454
ADB234468	ADB264757	ADB243646	
ADB249596	ADB232924	ADB263428	
ADB263270	ADB232927	ADB240500	
ADB231841	ADB245382	ADB253090	
ADB239007	ADB258158	ADB265236	
ADB263737	ADB264506	ADB264610	
ADB239263	ADB243027	ADB251613	
ADB251995	ADB233334	ADB237451	
ADB233106	ADB242926	ADB249671	
ADB262619	ADB262637	ADB262475	
ADB233111	ADB251649	ADB264579	
ADB240497	ADB264549	ADB244768	
ADB257618	ADB248354	ADB258553	
ADB240496	ADB258768	ADB244278	
ADB233747	ADB247842	ADB257305	
ADB240160	ADB264611	ADB245442	
ADB258646	ADB244931	ADB256780	
ADB264626	ADB263444	ADB264797	