

UNCLASSIFIED

AD NUMBER
ADB274630
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies only; Proprietary Info.; Apr 2001. Other requests shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott St., Ft. Detrick, MD 21702-5012.
AUTHORITY
USAMRMC ltr, 26 Nov 2002

THIS PAGE IS UNCLASSIFIED

AD_____

Award Number: DAMD17-98-1-8559

TITLE: Estrogen Metabolism and Prostate Cancer Risk

PRINCIPAL INVESTIGATOR: Paola Muti, M.D.

CONTRACTING ORGANIZATION: State University of New York
Amherst, New York 14228-2567

REPORT DATE: April 2001

TYPE OF REPORT: Final

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Distribution authorized to U.S. Government agencies only (proprietary information, April 2001). Other requests for this document shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott Street, Fort Detrick, Maryland 21702-5012.

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

20020131 161

NOTICE

USING GOVERNMENT DRAWINGS, SPECIFICATIONS, OR OTHER DATA INCLUDED IN THIS DOCUMENT FOR ANY PURPOSE OTHER THAN GOVERNMENT PROCUREMENT DOES NOT IN ANY WAY OBLIGATE THE U.S. GOVERNMENT. THE FACT THAT THE GOVERNMENT FORMULATED OR SUPPLIED THE DRAWINGS, SPECIFICATIONS, OR OTHER DATA DOES NOT LICENSE THE HOLDER OR ANY OTHER PERSON OR CORPORATION; OR CONVEY ANY RIGHTS OR PERMISSION TO MANUFACTURE, USE, OR SELL ANY PATENTED INVENTION THAT MAY RELATE TO THEM.

LIMITED RIGHTS LEGEND

Award Number: DAMD17-98-1-8559

Organization: State University of New York

Those portions of the technical data contained in this report marked as limited rights data shall not, without the written permission of the above contractor, be (a) released or disclosed outside the government, (b) used by the Government for manufacture or, in the case of computer software documentation, for preparing the same or similar computer software, or (c) used by a party other than the Government, except that the Government may release or disclose technical data to persons outside the Government, or permit the use of technical data by such persons, if (i) such release, disclosure, or use is necessary for emergency repair or overhaul or (ii) is a release or disclosure of technical data (other than detailed manufacturing or process data) to, or use of such data by, a foreign government that is in the interest of the Government and is required for evaluational or informational purposes, provided in either case that such release, disclosure or use is made subject to a prohibition that the person to whom the data is released or disclosed may not further use, release or disclose such data, and the contractor or subcontractor or subcontractor asserting the restriction is notified of such release, disclosure or use. This legend, together with the indications of the portions of this data which are subject to such limitations, shall be included on any reproduction hereof which includes any part of the portions subject to such limitations.

THIS TECHNICAL REPORT HAS BEEN REVIEWED AND IS APPROVED FOR PUBLICATION.

Nimisha Chandra Mirm
01/03/2002

REPORT DOCUMENTATION PAGEForm Approved
OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE April 2001	3. REPORT TYPE AND DATES COVERED Final (1 Oct 98 - 30 Mar 01)	
4. TITLE AND SUBTITLE Estrogen Metabolism and Prostate Cancer Risk			5. FUNDING NUMBERS DAMD17-98-1-8559	
6. AUTHOR(S) Paola Muti, M.D.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) State University of New York Amherst, New York 14228-2567 Email - muti@buffalo.edu			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012			10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Distribution authorized to U.S. Government agencies only (proprietary information, April 2001). Other requests for this document shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott Street, Fort Detrick, Maryland 21702-5012.				12b. DISTRIBUTION CODE
13. ABSTRACT (Maximum 200 Words) Experimental evidence shows that estrogens are related to prostate cancer risk – In addition recent data suggest that 16 α hydroxylated estrogen metabolites, biologically strong estrogens, are associated with cancer risk, while Z-hydroxyisted metabolites, with lower estrogenic activity, are weakly related to this disease. This study analyzes the association of prostate cancer with estrogen metabolism in a population-based study conducted in Erie and Niagara Counties – (the Protein Study) – Data in 81 prostate cancer cases and 318 control subjects showed that a higher ratio of 2 hydroxyestrone to 16 α hydroxyestrone uses associated with a reduced risk of prostate cancer.				
14. SUBJECT TERMS Prostate Cancer, Estrogen Metabolism, Epidemiology				15. NUMBER OF PAGES 47
				16. PRICE CODE
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

Table of Contents

Introduction.....	3
Body.....	5
Key Research Accomplishments.....	15
Reportable Outcomes.....	16
Conclusions.....	17
References.....	24
Appendices.....	35

Introduction

Prostate cancer is the most common cancer among men in the United States (IARC, 1995). And the second most common in the European Community (IARC, 1995). The causes of prostate cancer, however, remain largely unknown, with age, race, and family history being the only established risk factors (Nomura et al., 1997). The prostate gland has historically been considered the prototype of an androgen-dependent organ. However, there is evidence that estrogens may induce mitosis of prostatic epithelial cells in many species, including humans (Leav et al., 1978; Schulze et al., 1987). In humans, 16α hydroxyestrone and estriol are biologically significant estrogens, and their biologies can contribute to the overall expression of estrogenic action. Hydroxylation at the 16α position, one of the two major and mutually exclusive biotransformation pathways of estradiol, leads to estriol, a substance which is recognized as a biologically potent estrogen equivalent to estradiol (Clark et al., 1977). The other prominent pathway for metabolism is hydroxylation at C-2, producing 2-hydroxyestrone (Fishman J, 1963), a substance which has virtually no estrogenic activity (Gordon S., 1964; Martucci et al., 1977), except for some of the central activities of estradiol, such as the regulation of pituitary hormone release (Naftolin et al., 1975)

In 1979, Thomas Dao postulated that the way in which estrogens are metabolized might be important for cancer development (Dao T., 1979). The theory, known as "the unconventional estrogen hypothesis", proposed that the products of estrogen metabolism may be of etiological significance for hormone

related cancer. Several experimental, clinical and epidemiologic studies support this hypothesis (Schneider J et al., 1982; Clarck JK, et al., 1977; Fishman J. et al., 1980; Martucci et al., 1977). In *in-vitro* studies, 16 α -hydroxylation has been shown to have strong biological estrogenic activity (Telang NT, et al., 1992; Suto A, et al. 1993) and genotoxic characteristics (Telang NT et al., 1992), while 2-hydroxylation metabolites had virtually no peripheral estrogenic effects (Suto A, et al., 1993). In mice, 16 α -hydroxylation of estrone was associated with increased spontaneous incidence of tumors (Bradlow HL, et al., 1985).

When estrogen metabolism has been studied in relation to breast cancer risk in population-based studies, results supported the proposed hypothesis. In case-control studies an increase in estrone 16 α -hydroxylation in breast cancer cases was observed compared with healthy controls, in particular in postmenopausal women (Osborn MP, et al., 1993; Ursin G, et al., 1997; Kabat GC, et al., 1997; Zheng W, et al., 1977). Two prospective studies conducted to investigate the role of estrogen metabolism as predictor of breast cancer concordantly found that study participants characterized by elevated 2-hydroxyestrone /16 α -hydroxyestrone ratio had a 40% reduction in breast cancer risk compared with those in the lowest tertile (Meilahn EN et al., 1998; Muti P. et al., 2000).

The present report analyzes the association between prostate cancer and estrogen metabolism investigated in a case-control study. In particular, we tested the hypothesis that the pathway favoring 2-hydroxylation over 16 α -hydroxylation may be associated with decrease in prostate cancer risk.

Body

Ninty-one prostate cancer cases and 318 control subjects participated in the study between 1999 and 2001. However, the active recruitment of participants started only after one year since it took a long time to get the approval of the study protocols from Institutional Review Boards of the five area hospitals. In the Statement of Work (Task 1), we estimated that we would only require 3 months to complete the approval process. This task, however, took almost a full year. We encountered some difficulties in obtaining approval, in particular, from one of the area hospitals. These difficulties however, have been resolved.

During the entire study period, we collaborated with a few physicians and have finalized the questions on risk factors for prostate cancer as well as preparing letters for both cases and controls and developing strategies for recruitment. In addition, we tried to expand the base of prostate cancer case recruitment by including all private practices in Erie and Niagara Counties. After six months of contacts and calls and visits, we had several offices that were collaborating with us.

We devoted massive attention, time and energy to recruit prostate cancer cases:

a) we developed posters and brochures to be distributed to doctors' offices to support our recruitment efforts;

b) we also developed a staging form for the urologists to use in their private offices, memo pads and pens were distributed often along with a letter reminding them of the study and the need for their cancer case referral;

c) periodically, recent articles on prostate cancer etiology and treatment were sent to the area urologists again to remind them that the study was still on-going;

d) we met the HMO organizations and contacted them weekly to get support to our recruitment effort at the urologists's private offices;

e) in response to this, the Promedicus and the IPA sent letter to their participating urologists to help us in prostate cancer case enrollment. We also contacted a social organizations of prostate cancer cases such as US TOO and A FAMILY AFFAIR to inform their public about our research.

Although our efforts were equally spread all over the Erie and Niagara counties, most of our prostate cancer cases originated from three community hospitals in the Buffalo area: Buffalo General Hospital (BGH), Millard Fillmore Hospital (MFH), and the Department of Veterans Affairs Medical Center of Western New York Health Care System (VAMC). In particular, VAMC was remarkably efficient and open to our collaboration with a total of 64 prostate cancer cases recruited in the study there.

Prostate cancer patient recruitment: All patients, 65 to 85 years old, with newly diagnosed, cytologically and/or histologically confirmed prostate cancer have been considered for the study. All cases have been recruited in the study before starting any cancer treatment. After obtaining names of cases with newly

diagnosed prostate cancer, we mailed a request to the physician for permission to interview the patient. Once the approval was received, we sent a letter to the patient inviting them to participate. A written consent was obtained from the patient for both the blood draw and for the interview before the actual process took place

Eligibility criteria Since the major focus of the study was the relation of estrogen metabolism with prostate cancer risk, patients on hormonal treatment (current or in the six months prior the diagnosis), or affected with metabolic diseases affecting endocrine profile (i.e., hypogonadism, hyperadrenalism) were excluded. Those affected with chronic or acute liver diseases also were excluded because of their potential influence on the endocrine pattern. Patients with a previous history of cancer (except of skin cancer) have been excluded as well.

During the active phase of study recruitment (December 1999- April 2001), 146 prostate cancer cases have been identified. Of these 146, 108 met eligibility criteria, and were approved by the physicians at VAMC and invited to join the PROMEN study. Of these 108, 17 refused to participate. Thus, 74% (108/146) of the subjects were potential participants in the study, while the actual participation rate was 84% (91/108).

Prostate control recruitment: Control have been men, 65 to 85 years of age, selected from the rolls of the Health Care Finance Administration and matched

with prostate cancer cases on area of residence (first three digits of the zip code).

Eligibility criteria Eligibility criteria for controls was the same as for cases. We excluded men on hormonal treatment (current or in the six months prior to the contact), or affected with metabolic or endocrine diseases. Participants with a previous history of cancer (except of skin cancer) have been excluded as well.

During the same fifteen months of the study, 489 eligible controls were contacted. 84 of these potential candidates were deceased and 52 were too ill to participate (136/489, 27% of the sample). 318 of the remaining 353 subjects (90%) were enrolled and interviewed.

Since there is high prevalence of latent prostate carcinoma in men over age 50 (Breslow et al., 1977; Bostwick et al., 1992), we determined Prostate-Specific Antigen (PSA) in all blood samples obtained from controls. Controls with a PSA value higher than 4ng/ml were excluded from the control group according to the criterion proposed by the American Cancer Society Prostate Cancer Detection Project (Babaian et al., 1992) until the completion of diagnostic procedures to determine their true case-control status.

We have been able to identify 8 prostate cancer cases as a result of the PSA determinations that were done in controls. The PSA results were sent to the

participant's primary physician along with a letter to draw attention to the fact that it was outside of the normal range. The primary physician then referred the participant to an urologist where a biopsy was done to confirm prostate cancer. In our records, we modified the case-control status after the confirmation of prostate cancer diagnosis.

During the study period, procedures have been finalized for the ongoing maintenance of the biological specimen bank, processing of samples for immediate determinations and for storage, tracking of samples and mapping of the freezer as outlined in the Statement of Work (Task 2). For standardization purposes, morning spot urine was collected and fasting samples (57ml) of blood was drawn during the same time interval of the day (7:00AM – 9:00AM) from all participants. The time at specimens collection was recorded.

In accordance with the Statement of Work (Task 2), preparations for the data entry of the interview, maintenance of files from the computer-assisted interview, and entry of data from the sections of the interview completed by hand by the participant were developed during the recruitment phase of the study. The database for the prostate cancer study is written in Microsoft Access and contains information on all participants. Each form that a study participant filled out has a computer equivalent. Each of these databases is housed in the same location for security, backup, and data analysis purposes. These programs were written in Microsoft Access as well. Data validation, analysis, and compatibility

(for integration with SPSS) have been written in Visual Basic. Training has also been provided for the identification and staging of prostate cancer cases in area hospitals as well as the private physician offices.

Hormonal derminations in prostate cancer cases and control subjects (Tasks 4 to 5)

Stored urinary samples from prostate cancer cases and related controls were handled identically and randomly located across the laboratory runs. All laboratory personnel were blinded with regard to case-control status.

Analyses of 2-hydroxyestrone (2OHE1) and 16 α -hydroxyestrone (16 α -OHE1) were performed using a competitive solid-phase enzyme immunoassay (IMMUNA CARE Corporation, Bethlehem, PA). The urinary forms of these estrogen metabolites are found as glucuronide conjugates and require the removal of the sugar moiety before recognition by the monoclonal antibodies. A mixture of β -glucuronidase and arylsulphatase (glusulase from *H. Pomatia*, Sigma Chemical Co., St. Louis, MO) was used for this purpose. The enzyme digest was then neutralized. Assay incubation time was 3 hours at room temperature. The assay was read kinetically using a Ceres 900 HDI plate reader (Biotek Instruments, Winooski, VT) and the data were reduced using Kineticalc EIA Application software (Bio-Tek[®] Instruments). Both assays have been shown to demonstrate 100% recovery of metabolites with serial dilution and "spiking" of exogenous estrogens into urine samples. The EIA kits have been evaluated for validity and reproducibility and the values for each metabolite were compared

with values obtained by gas chromatography-mass spectrometry (Bradlow HL. Et al., 1998; Klug TL, et al., 1994; Sepkovic DW et al., 1994). As a measure of reproducibility, control samples were included and their values had to fall within two standard deviations from the mean of a continuous Levy-Jennings control plot. In addition, 10% duplicates were included with each batch of samples to determine reproducibility. Intra-assay coefficient of variation for 2-OHE1 and 16- α OHE1 were 3.6% and 3.8%, respectively. Interassay coefficients of variation were 5.9% and 10.2%, respectively.

Statistical Analysis (Task 6)

2-hydroxyestrone and 16 α -hydroxyestrone urinary levels were standardized by the total creatinine. We used conditional logistic regression to obtain the odds ratios of prostate cancer in relation to estrogen metabolites and their ratio (Breslow NE, Day NE, 1980). The independent variables of interest were 2-OHE1, 16 α -OHE1, and the ratio of 2-OHE1 and 16 α -OHE1 by tertiles of urine concentration. We based the cutoff points for each tertile on the distribution of the estrogen metabolites in controls. In the present analysis on estrogen metabolism, we included 70 prostate cancer cases since 21 participants refused to donate biological specimens and accepted only to answer to the questionnaires. 318 controls were available for the present analysis.

We identified age, weight, waist-to-hip ratio, as potential covariates according to their potential biologic relevance and logistic regression was used to control for these covariates. Since the information on all covariates was missing for part of the prostate cancer cases and control subjects, a total number of 61

prostate cancer cases and 235 controls were then considered in the analysis considering more complex models. In the initial regression model, we examined all variables. We evaluated each covariate for confounding by removing each from the fully adjusted model. Age, weight, and waist-to-hip ratio, did not substantially modify the results. None of the potential covariates was a confounder of the association between prostate cancer and estrogen metabolites and their ratio. Nevertheless, we included them in further analysis to provide fully adjusted estimates for comparison with those reported in the published literature, in particular with the previous studies on hormones and prostate cancer risk.

Study results (task 7).

Characteristics of the study population are reported in Table 1.

Prostate cancer cases were more likely to be slightly younger and to have higher weight, and a higher waist-to-hip ratio and a lower education. In a descriptive analysis reported in Table 2 conducted on control subjects, urinary levels of estrogen metabolites did not significantly differ by age strata although both 2-OHE1 and 16- α OHE1 were higher in older age in both prostate cancer cases and controls while the 2-OHE1/ 16- α OHE1 ratio was higher in younger subjects. African Americans showed to have a higher levels of estrogen metabolites and a higher ratio, although the difference did not reach the statistical significance due to the low number of African American study participants. Estrogen metabolites differ by waist-to-hip ratio tertiles with the lowest levels of estrogen metabolites and their ratio being observed in the

highest tertile of abdominal fat. Lighter subjects showed to have significant higher levels of 2-OHE1 and significant higher level of 2-OHE1/ 16- α OHE1 ratio.

Current smokers showed to have a higher levels of estrogen metabolites, and the lowest ratio, in comparison with never and former smokers. Finally, education, as an index of social and economic status and thus of life-style pattern, showed to have an effect on the estrogen metabolism. Participants classified in the lowest education level had the highest levels of both 2OHE1 and 16- α OHE1 and the lowest ratio .

Table 3 reports data on prostate cancer risk in relation to tertile of distribution of the estrogen metabolites, their ratio, and their 95% coefficients of variation. There was a light protective effect for the highest tertile of the 2-OHE1, however the confidence intervals included the unity. Conversely, there was a light risk effect in the highest tertile of 16 α -OHE1 and again the confidence intervals included the unity .

The increase in 2-OHE1/16 α -OHE1 ratio was associated with a reduction in odds ratios for prostate cancer across tertiles: the highest tertiles of 2-OHE1/16 α -OHE1 ratio had the lowest risk estimate, even after adjustment for covariates.

Key Research accomplishments.

The present research has found that:

- Estrogen metabolites can be found and measured in men;
- Estrogen metabolism is affected by age, abdominal adiposity, weight, smoking status, and education as an index of social and economic status;
- Estrogen metabolism may be implicated in prostate cancer development;
- Estrogen metabolism may become a potential tool for preventive strategies against the development of prostate cancer and for its therapeutic management.

Reportable Outcomes

The present data have to be considered preliminary results since the laboratory determinations have been completed on May 15th. However, we included in the present report data from a corollary preliminary study conducted in the context of the PROMEN study in collaboration with the Johns Hopkins University, Baltimore, MD. The study focused on the relation between human papilloma virus infection and risk of prostate cancer. Preliminary results of that study are included in the appendix in the form of a poster for a conference communication. The data will be presented by Dr. Sharita Womak, a NCI postdoctoral fellow mentored by Dr. Muti at Professional Development Workshop, sponsored by the Comprehensive Minority Biomedical Branch (CMBB), Office of Centers, Training, Resources (OCTR), Office of Deputy Director for Extramural Science (ODDES), and National Cancer Institute (NCI); June 4-6, 2001.

The final aim of the HPV determination is to evaluate potential biological interaction between viral infection and estrogen metabolism in the development of prostate cancer.

Conclusions.

The main finding of the present study was that estrogen metabolism pathway favoring 2-hydroxylation over 16 α -hydroxylation is associated with a reduced risk of prostate cancer risk. This is the first time that estrogen metabolism has been studied in relation to prostate cancer and the first observation supporting the potential protective role of estrogen metabolites characterized by low biological activity in prostate cancer development.

Analytical epidemiological studies on effects of estrogens in relation to prostate cancer risk, in particular serum estrone and estradiol have provided conflicting results. Serum estrone levels in patients with prostatic cancer have been reported to be higher than (Jackson et al., in African American, 1980; Ahluwalia et al., 1981; Hill et al., 1982; Zumoff et al., 1982), lower than (Drafta et al., 1982) and similar to (Bartsch et al., 1977a; Bartsch et al., 1977b; Jackson et al., in Nigerians, 1980; Nomura et al., 1988; Hsing et al., 1989; Barrett-Connor et al., 1990; Signorello et al., 1997) those in healthy controls. Serum estradiol showed similar conflicting results (Harper et al., 1976; Bartsch et al., 1977a; Bartsch et al., 1977b; Hammond et al., 1978; Ahluwalia et al., 1981; Hill et al., 1982; Hoisaeter et al., 1982; Meikle et al., 1982; Ranikko et al., 1983; Hulka et al., 1987; Hsing et al., 1989; Meikle et al., 1989; Andersson et al., 1993; Gann et al., 1996; Signorello et al., 1997 Drafta et al., 1982).

The inconsistency of results may be due to chance, low statistical power (low number of cases and controls), different strategy of control selection and different methodology in specimen collection (i.e., different control of some

sources of hormone variability as circadian rhythm or different storage conditions). It may also be that the relevant measure was not the serum level of estrone and estradiol but the estrogen metabolism.

Plasma concentrations of unconjugated estriol and 16α hydroxyestrone are low relative to estrone and estradiol (Fishman et al., 1976; Longcope et al., 1977) but their biological impact may be significant because of their lack of affinity for SHBG (sex hormone binding globulin) (Fishman et al., 1980). Considering that 16α hydroxyestrone is virtually unsequestered by SHBG, it may be that its biological effectiveness could equal or even exceed that of estradiol, even though the concentration of the latter in plasma is several-fold greater. In circumstances of comparable hormone secretion, therefore, estradiol metabolism shifted in the 16α hydroxylation direction could produce a hyperestrogenic milieu, while a predominance of 2-hydroxylation could produce hypoestrogenic conditions. 16α hydroxyestrone has been found to be elevated in strains of mice susceptible to breast cancer (Bradlow, 1985). In humans, estrogen metabolism has been primarily studied in women in relation to breast cancer risk. There is some evidence that 16α hydroxyestrone is elevated in women with breast cancer (Bradlow, 86; Fishman et al., 1984, Kabat et al., 1997, Muti. 2000).

There are a number of potential explanations for these findings, including potential effects of the neoplastic tissue on estrogen metabolism. Another potential source of bias in our study may be related to the preferential participation of prostate cancer cases from the Department of Veterans Affairs Medical Center of Western New York Health Care System (VAMC) while control

subjects were selected from the general population. Thus, differences in social and economic class and lifestyle may explain the present results. In order to limit the potential effects of social and economic and lifestyle bias, the control subjects were frequency-matched on area of residence (neighborhood controls) and randomly selected every two weeks from the general population of Erie and Niagara Counties. In the present study urinary estrogen metabolite levels were determined controlling for several sources of hormone variability by both inclusion criteria and highly standardized conditions at blood drawing. Cases were drawn before cancer treatment was begun, and control subjects were evaluated for potential presence of latent prostate cancer by serum analysis for prostate-specific antigen. Cases and controls on current or recent (six months prior the participation in the study) local or systemic hormonal therapy, and those affected with endocrine diseases, acute or chronic liver diseases were excluded to reduce, at least in part, some of the sources of hormone variability potentially influencing study results. Extreme care was taken in order to control for hormone circadian variation. All hormone determinations for all cases and controls were performed at the end of the recruitment phase of the study, during the last month of the study period to reduce interassay variability. The laboratory assaying estrogen metabolites was blinded to case-control status.

In conclusion, this study supports the hypothesis that there is difference in the way estrogens are metabolized between patients affected with prostate cancer and control subjects. Further studies are needed to corroborate these findings and to offer a new perspective on hormone involvement in prostate

cancer development. This, in turn, will allow to improve prevention and/or treatment.

Table 1. Characteristics of the Study Participants

	Cases		Controls	
	n	%	n	%
Age (years)				
45-64	12	19.7	21	8.9
65-74	36	59.0	142	60.4
75-85	13	21.3	72	30.6
total	61	100.0	235	100.0
Race				
Whites	44	72.1	213	90.6
African Americans	17	27.9	22	9.4
Waist-to-hip ratio (tertiles)				
1 st	19	31.1	79	33.6
2 nd	16	26.2	83	35.3
3 rd	26	42.6	73	31.1
Weight (tertiles)				
1 st	17	27.9	82	34.9
2 nd	21	34.4	76	32.3
3 rd	23	37.7	77	32.8
Smoking status				
Never	20	32.8	79	33.6
Current	6	9.8	15	6.4
Former	35	57.4	141	60.0
Education				
Less than high school	20	32.8	37	15.7
High school	18	29.5	81	34.5
College	23	37.7	117	49.8

Table 2. Levels of Estrogen Metabolites among Participants without Prostate Cancer

	n	2-hydroxyestrone (2-OHE1)		16 α -hydroxyestrone (16 α -OHE1)		2-OHE1/16 α -OHE1 ratio	
		Mean	SD	Mean	SD	Mean	SD
Age (yrs)							
45-64	21	5.40	2.24	4.42	2.22	1.35	0.60
65-74	142	5.67	2.77	4.33	1.74	1.36	0.53
75-85	72	5.54	2.34	4.52	1.83	1.29	0.47
<i>P</i>		0.88		0.77		0.64	
Race							
White	213	5.56	2.48	4.38	1.70	1.33	0.52
Black	22	6.11	3.53	4.53	2.71	1.46	0.58
<i>P</i>		0.34		0.72		0.27	
W/H ratio							
1 st	79	6.00	2.52	4.33	1.63	1.46	0.58
2 nd	83	6.00	2.64	4.61	2.12	1.38	0.45
3 rd	73	4.72	2.42	4.22	1.59	1.67	0.49
<i>P</i>		0.002		0.39		0.002	
Weight							
1 st	82	6.19	2.43	4.53	1.68	1.44	0.54
2 nd	76	5.55	2.75	4.43	2.05	1.32	0.48
3 rd	77	5.04	2.49	4.21	1.68	1.25	0.54
<i>P</i>		0.02		0.53		0.07	
Never	79	5.45	2.40	4.23	1.53	1.35	0.53
Current	15	5.85	2.23	5.63	2.53	1.15	0.51
Former	141	5.67	2.74	4.36	1.83	1.36	0.52
<i>P</i>		0.77		0.02		0.36	
Education							
<High sc	37	5.80	3.12	4.64	2.27	1.32	0.48
High sc	81	5.64	2.20	4.59	1.82	1.30	0.50
College	117	5.52	2.68	4.18	1.61	1.37	0.55
<i>P</i>		0.85		0.20		0.61	

N, sample size. SD, standard deviation. P, p value for test in mean differences.

W/H ratio, waist-to-hip ratio (tertile). Sc, School.

Table 3. Prostate Cancer Risk by Tertiles of Estrogen Metabolite Levels and Their Ratio

	n (Cases/Controls)	Crude OR (95% CI)	Adjusted OR * (95% CI)
2-hydroxyestrone (2-OHE1)			
1 st	22/76	1.00 (reference)	1.00 (reference)
2 nd	24/75	1.11 (0.57-2.14)	1.11 (0.54-2.26)
3 rd	15/84	0.62 (0.30-1.28)	0.61 (0.28-1.33)
16 α - hydroxyestrone (16 α -OHE1)			
1 st	19/79	1.00 (reference)	1.00 (reference)
2 nd	15/84	0.74 (0.35-1.56)	0.90 (0.41-1.96)
3 rd	27/72	1.56 (0.80-1.04)	1.73 (0.84-3.55)
2-OHE1/16 α -OHE1 ratio			
1 st	25/73	1.00 (reference)	1.00 (reference)
2 nd	22/77	0.83 (0.43-1.61)	0.80 (0.40-1.59)
3 rd	14/85	0.48 (0.23-0.99)	0.46 (0.21-0.99)

OR, odds ratio. 95% CI, 95% confidence interval.

* Adjusted for age, race, education, smoking status, and waist-hip-ratio using logistic regression.

References

- Abraham GE Ed. Radioassay Systems in Clinical Endocrinology. New York, Marcel Dekker, Inc. 1981
- Andersson S.O., Adami H.O., Bergstrom R., Wide L. Serum pituitary and sex steroid hormone levels in the etiology of prostatic cancer - a population based case-control study. *Br. J. Cancer* 1993; 68:97-102
- Babaian R.J., Mettlin C., Kane R. The relationship of prostate-specific antigen to digital examination and transrectal ultrasonography. Findings of the America Cancer Society National Prostate Cancer Detection Project. *Cancer* 1992; 69:1195-1200
- Baron JA, La Vecchia C, Levi F The antiestrogenic effect of cigarette smoking in women *Am J Obstet Gynecol* 1990; 162:502-14
- Block G., Hartman A.M., Dresser C.M., Carroll M.D., Gannon J., Gardner L. A data based approach to diet questionnaire design and testing. *Am. J. Epidemiol.* 1986; 124:453-469
- Block G., Woods M., Potosky A., Clifford C. Validation of a self-administered diet history questionnaire using multiple diet records. *J. Clin. Epidemiol.* 1990; 43:1327-1335
- Bolelli G.F., Muti P., Micheli A., Sciajno R., Franceschetti F., Krogh V., Pisani P., Berrino F. Validity for epidemiological studies of long-term cryoconservation of steroid and protein hormones in serum and plasma. *Cancer Epidemiology, Biomarkers & Prevention.* 1995; 4:509-513
- Bonen A., Long WY, MacIntyre KP Effects of exercise on the serum concentrations of FSH, LH, progesterone and estradiol *Eur J Appl Physiol* 1979; 42:15-23
- Bradlow HL, Hershcopf RJ, Martucci CP, Fishman J. Estradiol 16 α -hydroxylation in the mouse correlates with mammary tumor incidence and presence of murine mammary tumor virus: a possible model for the hormonal etiology of breast cancer in humans. *Proc. Natl. Acad. Sci. USA* 1985; 82:6295-6299
- Bradlow HL, Sepkovic DW, Klug T, Osborne MP. Application of an improved ELISA assay to the analysis of urinary estrogen metabolites. *Steroids* 63:406-413, 1998

Breslow N., Chan C.W., Dhorn G., Latent carcinoma of prostate at autopsy in seven areas. *Intl.J. Cancer*. 1977; 20:680-688

Brinckmann P., Biggemann M., Hilweg D. Prediction of the compressive strength of human lumbar vertebrae. *Spine*. 1989; 14:606-610

Browner WS, Pressman AR, Nevitt MC, Cauley JA, Cummings SR Association between low bone density and stroke in elderly women:the Study of Osteoporotic Fractures Stroke 1993; 24:940-946

Browner WS, Seely DG, Vogt TM, Cummings SR Non-trauma mortality in elderly women with low bone mineral density:Study of Osteoporotic Research Fractures Group *Lancet* 1991; 338:355-8

Burger H., van Daele P.L.A., Algra D., van den Ouweland F.A., Grobbee D.E., Hofman A., van Kuijk C., Schutte H.E., Birkenhager J.C., Pols H.A. P. The association between age and bone mineral density in men and women aged 55 years and over: The Rotterdam Study. *Bone Miner*. 1994; 25:1-13

Canadian Task force on the periodic health examination 1991 Periodic health examination, 1991 update: 3. Secondary prevention of prostate cancer. *Can med Assoc J*. 1991; 145:413-428

Carr B. Disorders of the ovaries and female reproductive tract in: *Endocrinology* (Wilson JD, Foster DW, Kronenberg HM, Larsen PR, eds) Saunders Company Philadelphia, USA, 1998, pp.751-817

Catalona W.J. Diagnosis, staging, and surgical treatment of prostatic carcinoma. *Arch. Intern. Med*. 1987; 147:361-80

Catalona W.J., Smith D.S., Rartliff T.L., et al. Measurement of prostate-specific antigen in serum as a screening test for prostate cancer. *New Engl. J. Med*. 1991; 324:1156-1161

Cauley JA, Lucas FL, Kuller LH, Vogt MT, Browner WS, Cummings SR Bone mineral density and risk of breast cancer in older women:the Study of Osteoporotic Fractures *JAMA* 1996; 276: 1404-1408

Clark JK, Paszko Z, Peck EJ. Nuclear binding and retention of the receptor estrogen complex: relation to the agonist and antagonist properties of estriol. *Endocrinology* 1977; 100:91-96

Cummings S.R., Block G., McHenry K., Baron R.B. Evaluation of two food-frequency methods measuring dietary calcium intake. *Am. J. Epidemiol*. 1987; 126:796-802

Dagogo-Jack S., Al-Ali N., Qurtom M Aumentation of bone mineral density in hirsute women J Clin Endocrinol Metab 1997; 82:2821-2825

Dai WS, Kuller LH, LaPorte RE, Gutai J, Falvo-Gerard L, Caggiula A The epidemiology of plasma testosterone levels in middle-aged men Am J Epidemiol 1981; 114:804-815

Dao T. Metabolism of estrogens in breast cancer Biochimica et Biophysica Acta 1979; 560:397-426

Deslypere J.P., Vermeulen A. Leydig cell function in normal men: effect of age, life-style, residence, diet and activity. J. Clin. Endocrinol. Metab. 1984; 59:955-962

Finkelstein J.S., Klibanski A., Neer R.M. Osteoporosis in men with idiopathic hypogonadotropic hypogonadism. Ann. Intern. Med. 1987; 106:354-361

Finkelstein J.S., Neer R.M., Biller B.M.K. Osteopenia in men with a history of delayed puberty. N. Engl. J. Med. 1992; 326:600-604

Fishman J, Martucci C. Biological properties of 16 alpha-hydroestrone: implications in estrogen physiology and pathophysiology. J Clin Endocrinol Metab 1980; 51:611-615

Foresta C., Ruzza G., Mioni R., Guarneri G., Gribaldo R., Meneghello A., Mastrogiacomo I. Osteoporosis and decline of gonadal function in the elderly male. Hormone Research. 1984; 19(1):18-22.

Garn S.M. The earlier gain and the later loss of cortical bone in nutritional perspective (Garn S.M., eds), Charles C., Thomas, Springfield, IL, 1970

Gilsanz V, Roe TF, Mora S, Costin, Goodman WG Changes in vertebral bone density in black girls and white girls during childhood and puberty N Engl J Med 1991; 325:1597-1600

Girasole G., Passeri G. Upregulation of osteoclastogenic potential for the marrow is induced by orchidectomy and is reversed by testosterone replacement in the mouse. J. Bone Miner. Res. 1992; 7:s96

Griffin J.E., Wilson J.D. Disorders of the tests and the male reproductive tract. In: Endocrinology (eds. Wilson, Foster, Kronenberg, Larsen). W.B. Saunders Company, Philadelphia, PA. 1998;pg836-837

Hammond GL, Ruokonen A, Konturi M, et al. The simultaneous radio-immunoassay of seven steroids in human spermatic and peripheral venous blood J Clin Endocrinol Metab 1977; 45:16-24

Hankinson S, Manson J, Spiegelman D, Willett WC, Longcope C, Speizer FE
Reproducibility of plasma hormone levels in postmenopausal women over a
2-3 year period *Cancer Epidemiol Biomarkers & Prev* 1995; 4:649-654

Harman S.M., Tsitouras P.D. Reproductive hormones in aging men.
I: Measurement of sex steroids, basal luteinizing hormone, and Leydig cell
response to human chronic gonadotropin. *J Clin Endocrinol Metab.* 1980;
51:35-40

Hsieh F.Y., Bloch D.A., Larsen M.D. A simple method of sample size
calculation for linear and logistic regression. *Statistics in Medicine.* 1998;
17:1623-1634

Jackson J.A., Riggs M.W., Spiekerman A.M. Testosterone deficiency as a
risk factor for hip fracture in men: A case-control study. *Am. J. Med. Sci.*
1992; 304:4-8

Johansson A.G., Lindh E, Blum WF, Kollerup G, Sorensen OH, Ljunhall S
Effects of GH and IGF-I in men with idiopathic osteoporosis *J Clin Endocrinol
Metab* 1996; 81:44-48

Johansson A.G., Forslund A., Hambræus L. Growth-hormone-dependent
insulin-like growth factor binding protein is a major determinant of bone
mineral density in healthy men. *J. Bone Miner. Res.* 1994; 9:915-921

Jones JJ, Clemmons DR. Insulin-like growth factors and their binding proteins:
biological actions. *Endocr Rev* 1995; 16:3-34

Kaaks R, Nutrition, hormones, and breast cancer: Is insulin the missing link?
Cancer Causes and Control 1996; 7:605-25

Kabat GC, Chang CJ, Sparano JA, Sepkovic DW, Hu X-P, Khalil A,
Rosenblatt R, Bradlow HL. Urinary estrogen metabolites and breast cancer: a
case-control study. *Cancer Epidemiol Biomarkers Prev* 1997; 6:505-509

Keinbaum D.G., Kupper L., Muller KE, Nizam A *Regression Diagnostics in:
Applied Regression Analysis and Multivariable Methods*, Duxbury Press,
Pacific Grove, 1998

Keinbaum D.G., Kupper L., Morgenstern H. Information bias in Epidemiologic
Research- Principles and quantitative methods (Keinbaum D.G., Kupper L.,
Morgenstern H., eds.) pp. 220-241, Von Nostrand Reinhold Company, New
York, 1982

Kelly P.J., Pocock N.A., Sambrook P.N., Eisman J.A. Dietary calcium, sex
hormones, and bone mineral density in men. *Br. Med. J.* 1990; 300:1361-
1364

Klug TL, Bradlow HL, Sepkovic, DW. Monoclonal antibody-based enzyme immunoassay for simultaneous quantitation of 2- and 16 α -hydroxyesterone in urine. *Steroids* 59:648-655, 1994.

Kosary C.L., Ries L.A.G., Miller B.A., Hankey B.F., Harras A., Edwards B.K., eds SEER Cancer Statistics Review 1973-1992 (tables and Graphs). DHHS Pub. No. (NIH) 96-2789 Bethesda, MD: National Cancer Institute, 1995

Liel Y, Edwards J, Shary J, Spicer TJ, Gordon L, Bell NH The effect of race and body habitus on bone mineral density of the radius, hip, and spine in premenopausal women *J Clin Endocrinol Metab* 1988; 66:1247-1250

Lindsay R., Hart D.M., Forest C., Baird C. Prevention of spinal osteoporosis in oophorectomized women. *Lancet*. 1980; 2:1151-1154

Looker AC, Harris TB, Madans JH, Sempos CS Dietary calcium and hip fracture risk: the NHANES I epidemiologic follow-up study *Osteoporosis Int* 1993; 3: 177-184

MacDonald PC, Madden JD, Brenner PF, et al., Origin of estrogen in normal men and in women with testicular feminization *J Clin Endocrinol Metab* 1979; 49:905-916

Machwate M, Zerath E, Holy X, IGF-I increases trabecular bone formation and osteoblastic cell proliferation in unloaded rats *Endocrinology*, 1994; 134:1031-1038

Martucci C, Fishman J. Direction of estradiol metabolism as a control of its hormonal action-uterotrophic activity of estradiol metabolism. *Endocrinology* 1977; 101:1709-1715

Meier D.E., Orwoll E.S., Keenan E.J., Fagerstrom R.M. Marked decline in trabecular bone mineral content in healthy men with age. Lack of association with sex steroid levels. *J. Am. Geriatr. Soc.* 1987; 35:189-197

Meier D.E., Orwoll E.S., Jones JM Marked disparity between trabecular and cortical bone loss with age in healthy men: measurement by vertebral computed tomography and radioal photon absorptometry *Ann Inter Med* 1984; 01:605-612

Mohan S., Baylink D Autocrine and paracrine aspects of bone metabolism *Growth Genet Horm* 1990; 6:1-9

Muti P, Trevisan M, Modlich F, Krogh V Why and how to use a biological specimen bank in epidemiological and clinical research: *Nutr Metab Cardiovasc Dis* 1998; 8:200-204

Muti P, Trevisan M, Micheli A, Krogh V, Bolelli G.F, Sciajno R, Berrino F. Alcohol consumption and total estradiol in replicated measures over a year in premenopausal women. *Cancer Epidemiology, Biomarkers & Prevention*, 7:189-193, 1998

Muti P., Trevisan M., Micheli A., Krogh V., Bolelli G.F., Sciajno R., Berrino F. Reliability of serum hormones in premenopausal and postmenopausal women over a one-year period. *Cancer Epidemiol. Biomarker Prev.* 1996; 5:917-922

Muti P, Micheli A, Krogh V, Freudenheim JL, Yang J, Imamura D, Schünemann HJ, Trevisan M, Berrino F. *Estrogen metabolism and breast cancer risk* *Epidemiology* 11:635-640, 2000

Nankin H.R., Calkins J.H. Decrease bioavailable testosterone in aging normal and impotent men. *J Clin Endocrinol Metab.* 1986; 63:1418-1420

Nomura AMY, Kolonel, LN Prostate cancer: a current prospective *Am J Epidemiol* 1991; 13:200-227

Olsson H, Hagglung G, Reduced cancer morbidity and mortality in a prospective cohort of women with distal forearm fractures *Am J Epidemiol* 1992; 136:422-427

Orgebin-Crist M.C., Eller B.C., Danzo B.J. The effect of estradiol, tamoxifen, and testosterone on the weights and the histology of the epididymis and accessory sex organs of sexually immature rabbits. *Endocrinology.* 1983; 113:1703-1709

Orwoll ES, Kelin RF Osteoporosis in men in: *Osteoporosis* (Marcus R, Feldman D, Kelsey J, eds), S. Diego Academic Press, 1996, pp.745-784

Osborne MP, Bradlow HL, Wong GYC, Telang NT. Upregulation of estradiol C16 α -hydroxylation in human breast tissue: a potential biomarker of breast cancer risk. *J Natl Cancer Inst* 1993; 85:1917-1920

Oursler M.J., Cortese C., Keeting P.E., Anderson M.A., Bonde S.K., Riggs B.L., Spelsberg T.C. Modulation of transforming growth factor- β production in normal human osteoblast-like cells by 17 β -estradiol and parathyroid hormone. *Endocrinology.* 1991; 129:3313-3320

Panico S., Pisani P., Muti ID ec hione C., Totis A. Diurnal variation of testosterone and estradiol: a source of bias in comparative studies on breast cancer. *J Endocrinol Invest.* 1990 – 13:423-427

- Persson I, Adami HO, McLaughlin JK, Naessen T, Fraumeni JF Jr Reduced risk of breast and endometrial cancer among women with hip fracture (Sweden) *Cancer Causes and Control* 1994; 5:523-528
- Potosky A.L., Block G., Hartman A.M. The apparent validity of questionnaires is influenced by number of diet-record days used for comparison. *J. Am. Dietetic Assoc.* 1990; 90:810-813
- Prentice A, Laskey MA, Shaw J, Cole TJ, and Frazer DR Bone mineral density content of Gambian and British children aged 0-36 months *Bone Miner.* 1990; 10:211-224
- Ravaglia G, Forti P, Maioli F, Nesi B, Pratelli L, Cucinotta D, Bastagli L, Cavalli G Body composition, sex steroids, IGF-I, and bone mineral density in aging men *Journal of Gerontology* 2000; 55: 516-521
- Reid IR, Plank LD, Evans C Fat mass is an important determinant of whole body bone density in premenopausal women but not in men *J Clin Endocrinol Metab* 1992; 75, 779-782
- Reiter EO, Rosenfeld RG Normal and aberrant growth In: Wilson JD, Foster DW, Kronenberg HM, Larsen PR, eds. *Endocrinology*. Sect.16, WB Saunders Com, Philadelphia, 1998, pg. 1447-1457
- Riggs BL, Melton LJ Medical Progress: Involutional osteoporosis *N Engl J Med* 1986; 1676-1686
- Rosen CJ, Donahue LR, Hunter SJ Insulin-like growth factors and bone: the osteoporosis connection *Proc Soc Exp Biol Med* 1994; 206:83-102
- Rothman K. *Modern Epidemiology*. Little Brown, Boston. 1986
- Rothman K.J., Greenland S. *Modern Epidemiology*. Lippincott-Raven Publishers, Philadelphia, PA. 1998
- Seeman E., Melton L.J., O'Fallon W.F. Risk factors for osteoporosis in men. *Am. J. Med.* 1983; 75:977-983
- Sepkovic DW, Bradlow HL, Ho G, Hankinson S, Gong L, Osborne MP, Fishman J. Estrogen metabolite ratios and risk assessment of hormone related cancers: Assay validation and prediction of cervical cancer risk. *Ann New York Acad Sci* 768:312-316, 1995.
- Sepkovic DW, Bradlow HL, Michnovicz J, Murtezani S, Levy I, Osborne MP. Catechol estrogen production in rat microsomes after treatment with indole-3-carbinol, ascorbigen, or B-naphthaflavone: A comparison of stable isotope dilution gas chromatography-mass spectrometry and radiometric methods. *Steroids* 59:318-323, 1994.

Siiteri P.K., MacDonald P.C. Role of extraglandular estrogen in human endocrinology. In: Greep RO, Astwood EB, eds Handbook of Physiology Sect 7: Endocrinology. Vol II. Female Reproductive System. Part I. Washington, DC: American Physiological Society. 1973;615-629

Slemenda C., Hui S.L., Longcope C., Johnston C.C. Sex steroids and bone mass: A study of changes about the time of menopause J. Clin. Invest. 1987; 80:1261-1269

Slemenda C.W., Longcope C., Zhou L., Hui S.L., Peacock M., Johnston C.C. Sex steroid and bone mass in older men. J. Clin. Invest. 1997; 7:1755-1759

Smals AGH, Kloppenborg PWC, Njo KT Alcohol-induced Cushingoid syndrome BMJ 1976; 2:1298-1299

Rees LH, Besser GM, Jeffcoate WJ Alcoholic pseudo-Cushing's Lancet 1977; 1:726-728

Siiteri PK, McDonald PC, Role of extraglandular estrogen in human endocrinology In: Greep RO, Astwood EB, eds. Handbook of physiology, Sec. 7: Endocrinology Vol II. Washington, DC: American Physiological Society 1973; 615-629

Silverman SL, Madison RE. Decreased incidence of hip fractures in Hispanics, Asians, and blacks: California Hospital Discharge Data Am J Public Health 1988; 78:1482-1483

Smith D.A.S., Walker M.S. Changes in plasma steroids and bone density in Klinefelter's syndrome. Calcif. Tissue Int. 1997; 22:225-228

Smith E.P., Boyd G.R., Frank H., Takahashi R.M., Cohen R.M., Specker B., Williams T.C., Lubahn D.B., Korach K.S. Estrogen resistance caused by a mutation in the estrogen-receptor gene in a man. N. Engl. J. Med. 1994; 331:1056-1061

Schneider J, Kinne D, Fracchia A, Pierce V, Anderson KE, Bradlow HL, Fishman J. Abnormal oxidative metabolism in women with breast cancer Proc. Natl. Acad. Sci. USA 1982; 79:3047-3051

Snow R.C., Barbeiri R.L., Frish R.E. Estrogen 2-hydroxylase oxydation and menstrual function among elite oarswomen. J. Clin. Endocrinol. Metab. 1989; 69:369-376

Snow-Harter C, Whalen K, Myburg K, Arnaud S, Marcus R Bone mineral density, muscle strength, and recreational exercise in men J Bone Miner Res 1992; 7:1291-1296

Sobell J., Block G., Koslowe P., Tobin J., Andres R. Validation of a retrospective questionnaire assessing diet 10-15 years ago. *Am. J. Epidemiol.* 1989; 130:173-187

Solomon L. Bone density in ageing Caucasian and African populations *Lancet* 1979; 2:1326-1330

Stanley H.L., Schmitt B.P., Poses R.M. Does hypogonadism contribute to the occurrence of a minimal trauma hip fracture in elderly men? *J. Am. Geriatr. Soc.* 1991; 39:766-771

Staszewski J., Haenzel W. Cancer mortality among the Polish-born in the United States. *J. Natl. Cancer Inst.* 1965; 35:291-297

Suto A, Bradlow HL, Wong GY, Osborne MP, Telang NT. Experimental down-regulation intermediate biomarkers of carcinogenesis in mouse mammary epithelial cells *Breast Cancer Res Treat* 1993; 27:193-202

Sugimoto T, Nishiyama K, Kuribayashi F, Serum levels of insulin-like growth factor (IGF)-I, IGF Binding Protein (IGFBP)-2 and IGFBP-3 in Osteoporitic Patients with and without spinal fractures. *J of Bone and Mineral Research* 1997; 12:1272-1279

Telang NT, Suto A, Wong GY, Osborn MP, Bradlow HL. Induction by estrogen metabolite 16 α -hydroxyestrone of genotoxic damage and aberrant proliferation in mouse mammary epithelial cells *J Natl Cancer Inst* 1992; 84:634-638

Tietz NW, Ed.: *Fundamentals of Clinical Chemistry*, Saunders Company, Philadelphia 1987

Thorner MO, Vance ML, Laws ER, Horvath E, Kovacs K The anterior pituitary in "Williams textbook of Endocrinology" (Wilson JD, Foster DW, Kronenberg HM, Larsen PR, eds), 1998, pp.249-340

Ursin G, London S, Stanczyk FZ, Gentzschein E, Paganini-Hill A, Ross RK, Pike MC. A pilot study of urinary estrogen metabolism (16 α -OHE1 and 2-OHE1) in postmenopausal women with and without breast cancer. *Environ Health Perspect* 1997; 105 (Suppl.3):601-605

Vermeulen A., Deslypere J.P. Intratesticular unconjugated steroids in elderly men. *J. Steroid Biochem.* 1986; 24:1079-1083

- Vermeulen A., Verdonck G. Representativeness of a single point plasma testosterone level for the long term hormonal milieu in men. *J. Clin. Endocrinol. and Metabolism*. 1992; 74:939-942
- Vesterby A., Mosekilde L., Gundersen H.J.G. Biologically meaningful determinants of the in vitro strength of lumbar vertebrae. *Bone*. 1991; 12:219-224
- Wahner H. Use of Densitometry in management of osteoporosis in: *Osteoporosis* (Marcus R, Feldman D, Kelsey J, eds), S. Diego Academic Press, 1996, pp.1055-1074
- Walther P.J. Prostate cancer screening. Why the controversy? *Surg. Oncology Clin. North America*. 1995; 4:315-334
- Wanjus S. Dai, Kuller L.H., LaPorte R.E., Gutai J.P., Falvo-Gerard L., Caggiula A. The epidemiology of plasma testosterone levels in middle-aged men. *Am. J. Epidemiol.* 1981; 114:804-817
- Westerling K. The role of estrogen metabolism in aging-related diseases: Breast cancer, osteoporosis and coronary heart disease. Communication at the AACR- Special Conference of Basic and Clinical Aspects of breast Cancer, Keystone, Colorado. March 7-12, 1997
- Wilson J.D. Recent studies on the mechanism of action of testosterone. *N. Engl. J. Med.* 1972; 287:1284-1291.
- Wilson JD, Metabolism of testicular androgens In: Wilson JD, Foster DW, Kronenberg HM, Larsen PR, eds. *Endocrinology*. Sect.16, WB Saunders Com, Philadelphia, 1998: 822-829
- Wink C.S. and Felts W.J.L. Effects of castration on the bone structure of male rats: a model of osteoporosis. *Calcif. Tissue Int.* 1980; 32:77-82
- Young H., Kent J. Plasma testosterone levels in patients with prostatic carcinoma before and after treatment. *J Urol.* 1968; 99:788-792
- Yu H, Rohan T. Role of the insulin-like growth factor family in cancer development and progression. *J Natl Cancer Inst* 2000; 92:1472-89.
- Zeyneloglu HB, Arici A, Olive DL. Environmental toxins and endometriosis. [Review] [118 refs] [Journal Article. Review. Review, Academic] *Obstetrics & Gynecology Clinics of North America*. 24(2):307-29, 1997 Jun.
- Zheng W, Dunning L, Jin F, Holtzman J. Correspondence re: Kabat GC, Chang CJ, Sparano JA, Sepkovic DW, Hu X-P, Khalil A, Rosenblatt R, Bradlow HL Urinary estrogen metabolites and breast cancer: a case-control study. *Cancer Epidemiol Biomarkers Prev* 1977; 6:500-504

Ziegler RG, Rossi SC, Fears TR, Bradlow HL, Adlercreutz H, Sepkovic DW, Kiuru P, Wahala K, Vaught JB, Donaldson JL, Falk RT, Fillmore C-M, Siiteri PK, Hoover RN, Gail MH. Quantifying estrogen metabolism: An evaluation of the reproducibility and validity of enzyme immunoassays for 2-hydroxyesterone and 16 α -hydroxyesterone in urine. *Environ Health Perspect* 105(3): 607-614, 1997.

Zhang Y, Kiel DP, Kreger BE, Cupples LA, Ellison RC, Dorgan JF, Schatzkin A, Levy D, Felson D Bone Mass and the risk of breast cancer among postmenopausal women *N Engl J Med* 1997; 336:611-617

Zhang Y, Felson DT, Ellison RC, Kreger BE, Schatzkin A, Dorgan JF, Cupples LA, Levy D, Kiel DP Bone mass and risk of colon cancer among postmenopausal women *Am J Epidemiol* 2001; 153:31-37

Ziegler R. G., Rossi S.C., Fears T. R., Brad low H. L., Adlercreutz, Sepkovic D., Kiuru P., Wahala K., Vaught J.B., Donaldson J.L., Falk R.T., Fillmore C.M., Siiteri P.K., Hoover R.N., Gail M.H. Quantifying estrogen metabolism: an evaluation of the reproducibility and validity of enzyme immunoassays for 2-hydroxyestrone and 16 α -hydroxyestrone in urine. *Environ. Health Perspect.* 1997; 105:607-614

Zumoff B., Levin J., Strain G.W., Rosenfeld R.S., O'Connor J., Freed S.Z., Kream J., Whitmore W., Fukushima D., Heilman L. Abnormal levels of plasma hormones in men with prostate cancer: evidence toward a 'two-disease' theory. *Prostate.* 1982; 3:579-588

Appendices

Poster on HPV and prostate cancer risk.

Human papillomavirus seropositivity in PROMEN case and control study participants

Preliminary Results


Background

- Prostate Cancer is leading cancer diagnosed among men (*SEER, 1988-92*)
- 2nd leading cause of male cancer deaths in the country (*Parkin et al, 1997*)
- In 2000, 180,400 men would be diagnosed and 37,000 deaths (*ACS facts and figures, 1999*)
- Mortality rates are more than twice as high as rates in white men (*SEER 1988-92*)
- Factors suggested to play a role: age, weight, high fat diet, vasectomy, physical activity,
 - and infectious agents (*Ross RK and Schottenfield D, 1996*)
- HPV has been reported from prostate tissues in several recent investigations (*Cuzick, 1995*)

Common Risk Factors and Preventive Factors for Prostate Cancer

Risk factors

Hormonal factors
 Advancing age
 Family history/genetics
 Dietary fat



BPH
 Vasectomy

Preventive measures

Vitamin E
 Soy
 Selenium
 Physical Activity

Gallagher et al, CMAJ 159(7) 1998; updated

Role of Sexual Behavior in Prostate Cancer Etiology

- Key et al., 1995 (systematic review)
 - sexual behavior emerged as one of the stronger and more consistently identified risk factors
 - increased RR for early intercourse and for multiple sexual partners
 - highest RR was for a history of sexually transmitted disease (RR=1.86; 95% CI, 1.43-2.42)
 - recall and selection bias
- Two cohort studies
 - (Nilsen TL et al, 2000 and Hayes RB et al, 2000)
 - found that men who were divorced or separated ↑
 - syphilis, gonorrhea, sex with prostitutes and unprotected sexual intercourse were indicators of contact with a STI ↑

HPV infection of the Prostate and Prostate Cancer Risk

- HPV has been detected in prostate tissues in several recent investigations
- Polymerase chain reaction (PCR) and other methods
- Case-control studies have provided conflicting results

Table 1
Summary of Studies Investigating HPV in the Prostate Specimen

Reference	HPV detection	No.	Subjects		HPV %
			Type		
McNicol 1990	Southern blot for HPV-16 and -18 SPP	4	PCa cases	75%	
		12	BPH control	33%	
McNicol 1990	E6 PCR HPV-16, -18	4	PCa cases	100%	
		5	BPH controls	93%	
			normal autops.	20%	
Masood 1991	ISH for HPV-6,11, 16,18,31,33,35	20	PCa cases	none	
		20	BPH controls	none	
MCNicol 1991	E6 PCR HPV-16, -18	27	PCa cases	52%	
		56	BPH controls	63%	
Anwar 1992	E6 PCR HPV-16, -18,33	68	PCa cases	41%	
		10	BPH controls	none	
		10	normal autops.	None	
Effort 1992	Differential E6 PCR 30 HPV-16 and 18	30	Pac cases	none	
		8	Cervical care,	38%	
Dodd 1993	Reverse transcription; PCR E6/E7 MRNA of HPV-16	7	PCa cases	43%	
		10	BPH controls	50%	
Tu 1994	L1 consensus primer-43 PCR	17	PCa cases	2%	
		1	metastases	6%	
		1	normal	none	
Mayer-Lalle 1995	E6 PCR HPV-16; and 18	17	PCa cases	53%	
		22	BPH controls	32%	
Wideroff 1996	L1 consensus primer 56 PCR, E6 PCR for SPP HPV-6,11,16,18,31, 33, and 45	42	PCa cases	L1 13%	E6 0%
			BPH controls	L1 10%	E6 0%
Suzuki 1996	L1 consensus primer 51	51	PCa cases	16%	
Strickler 1998	L1 consensus primer 63 And E6 PCR	61	PCa cases	none	
			BPH controls	none	
	Serum antibodies HPV -16 and -11	63	PCa cases	no difference	
		144	BPH controls		

HPV infection of the Prostate and Prostate Cancer Risk

- 2 Prospective Studies (Dillner et al, 1998 and Hisada et al, 2000)
increased prostate cancer risk among subjects
with HPV 16/18 antibody compared with those w/o the antibody
 - » 2.4 to 2.6 fold risk ($p < 0.005$)
 - » OR= 2.7; 95% (CI, 0.9-7.9)

Specific Aims

1. To determine the HPV prevalence of serum antibodies to HPV 16, 18, and 31 in stored serum specimens collected from the first 79 prostate cancer cases and 280 control subjects in the ongoing PROMEN study.
2. To determine the strength of the association of HPV seropositivity in PROMEN case and control study participants.

Elisa Assay

- HPV 16, 18, and 31 Virus-like particles (VLP) prepared from insect cells (expressing the L1 and L2 viral capsid protein)
- VLP's coat wells of 96-well microtiter plate
- control wells coated with PBS
- plates washed 5 times
- add test serum to each antigen
- incubated 2 hrs at 37°C
- plates washed 5 times
- VLP reaction detected with horseradish peroxidase conjugated Protein G
- incubate 30min
- hydrogen peroxide solution (incubate 30 min)

PROMEN Study

PROstate cancer and Metabolism of EstrogeNs

- An ongoing study case-control investigation of the of prostate cancer and two estrogen hormones.
- This study is being conducted in the greater Buffalo, New York area.
- Funded by the U.S Army Prostate Cancer Research Program
- Includes men, 49-85 with incident, primary, pathologically confirmed prostate cancer
 - diagnosed at the Department of Veterans Affairs Medical Center of Western New York (VAMC),
 - ECMC, Buffalo General, Millard Fillmore and RPCI

PROMEN Study

continue....

- Data collected:

- physical measurements*: weight, height, waist to hip, etc

- questionnaire*-- sexual behavior across different age periods, expressed as number of partners; age at first intercourse;

- lifetime alcohol intake history, reproductive history, diet history, medical history, residential history, physical activity and vasectomy

- clinical specimens*-- blood and urine specimens

- questionnaires and measurements repeated in 51 controls over a one year period (test reproducibility)

PROMEN Study

continue....

- Cases:

- men attending annual physical exam at VAMC and other clinics in Buffalo 49-85 years old, with newly diagnosed histologically confirmed prostate cancer prior to cancer treatment

- to assure more accurate case definition, only clinical apparent prostate cancer cases (stage B and over) were included

- participation rate was 84%

PROMEN Study

continue....

- Controls:

- men 49-85 years old
 - selected from driver's licenses list of Erie and Niagara county residence;
 - controls age 65 and over are randomly selected from the rolls
 - of the Health Care Finance Administration
 - frequency matched to cases on age, race, and area of current residence

participation rate 72%

PROMEN Study

continue....


- Exclusion criteria:

- cases: on hormonal treatment or with metabolic/endocrine diseases
 - affected with chronic or acute liver disease
 - with previous history of cancer

- controls: on hormonal treatment or with metabolic disease or endocrine diseases
 - PSA value higher than 4ng/ml (to exclude potential latent carcinoma)

RESULTS: Figure 1

Percentage of PROMEN Study Participants by Selected Variables



RESULTS: Table 2

Reliability Coefficient Selected Variables

Waist to Hip	0.82
Height (cm)	0.96
Weight (kg)	0.90

Results: Table 3
Sexual Behavior and Reproductive Variables of PROMEN

Selected Variables	Cases Mean (SD)	Controls Mean (SD)	Reliability coefficient
Age first intercourse	17.9 (4.20)	19.5 (4.02)*	0.97
# sex partners at 20-30	3.8 (4.4)	3.6 (4.7)	.70
# sex partners at 31-40	3.0 (4.6)	1.8 (4.6)*	.87
# sex partners at 41-50	2.8 (4.3)	1.5 (3.8)*	.49
# sex partners at 51-60	2.1 (3.3)	1.3 (3.1)	.91
# sex partners at 61-70	1.8 (3.7)	1.1 (2.6)	.87
# subjects with vasectomy	7 (7%) †	19 (6.6%) †	1.00**

*t-test for unpaired data ($P < 0.05$); † prevalence in percentage; ** percent agreement

Results: Table 4
Adjusted Odds Ratio (OR) of PROMEN Study Participants for HPV 16, 18, and 31 seropositivity

Infectious agent	Percent Positive		Crude OR (95% CI)	OR adjusted for age, race and BMI	OR adjusted for age, race, WHR
	Cases	Controls			
HPV 16	3	.4	7.3 (.65-81.0)	2.7 (.15-45.1)	2.8 (.23-34.4)
HPV 18	11	11	1.01 (.49-2.4)	1.1 (.48-2.6)	0.9 (.37-2.1)
HPV 31	4	.7	5.5 (.90-33.4)	4.5 (.55-37.2)	6.9 (1.1-44.6)
HPV 16, 18, 31	20	10	1.5 (.74-3.1)	1.4 (.63-3.1)	1.3 (.62-2.8)

Conclusion

- 1) ↑ WHR and weight are related to increased risk
» *Cancer Epid., Biomark. & Prev. (Hsing et al, 2000)*
- 2) Increased number of sexual partners over decades and earlier age at first intercourse were shown for cases (..recall bias may exist)
- 3) Serology data indicate a potential association between HPV infections and prostate cancer risk, however estimates always included unity
» small sample size or
» large variability of the serological method

Future Studies

- 1) Strength of association of prostate cancer using two more sensitive HPV detection assay/method performed in urine and compare results with current serology data for HPV 16, 18, 31
» *L1 consensus primer-based polymerase chain reaction (PCR)*
» *Hybrid Capture II probe B)*
- 2) To evaluate potential variants in HPV 16, 18, and 31 in relation to prostate cancer risk
- 3) Immunologic characteristics in relation to HPV and cancer risk

Limitations

- 1) Serological measurements of HPV is consistently reported as an insensitive method compared to other assays/method***
 - » low titer
 - » not detected in a proportion of subjects previously or currently HPV positive
- 2) Cause-Effect relationship can not be derived from case-control study**

Acknowledgements

Dr. P Muti
Dr. R Viscidi
Dr. KV Shah



DEPARTMENT OF THE ARMY
US ARMY MEDICAL RESEARCH AND MATERIEL COMMAND
504 SCOTT STREET
FORT DETRICK, MARYLAND 21702-5017

REPLY TO
ATTENTION OF:

MCMR-RMI-S (70-1y)

26 Nov 02

MEMORANDUM FOR Administrator, Defense Technical Information
Center (DTIC-OCA), 8725 John J. Kingman Road, Fort Belvoir,
VA 22060-6218


SUBJECT: Request Change in Distribution Statement

1. The U.S. Army Medical Research and Materiel Command has reexamined the need for the limitation assigned to technical reports written for this Command. Request the limited distribution statement for the enclosed accession numbers be changed to "Approved for public release; distribution unlimited." These reports should be released to the National Technical Information Service.

2. Point of contact for this request is Ms. Kristin Morrow at DSN 343-7327 or by e-mail at Kristin.Morrow@det.amedd.army.mil.

FOR THE COMMANDER:

Encl



PHYLLIS M. RINEHART
Deputy Chief of Staff for
Information Management

ADB263708
ADB257291
ADB262612
ADB266082
ADB282187
ADB263424
ADB267958
ADB282194
ADB261109
ADB274630
ADB244697
ADB282244
ADB265964
ADB248605
ADB278762
ADB264450
ADB279621
ADB261475
ADB279568
ADB262568
ADB266387
ADB279633
ADB266646
ADB258871
ADB266038
ADB258945
ADB278624