

UNCLASSIFIED

Defense Technical Information Center
Compilation Part Notice

ADP011066

TITLE: Alternobaric Vertigo: Incidence in Portuguese Air Force Pilots

DISTRIBUTION: Approved for public release, distribution unlimited

This paper is part of the following report:

TITLE: Operational Medical Issues in Hypo-and Hyperbaric Conditions
[les Questions medicales a caractere operationel liees aux conditions
hypobares ou hyperbares]

To order the complete compilation report, use: ADA395680

The component part is provided here to allow users access to individually authored sections of proceedings, annals, symposia, etc. However, the component should be considered within the context of the overall compilation report and not as a stand-alone technical report.

The following component part numbers comprise the compilation report:

ADP011059 thru ADP011100

UNCLASSIFIED

Alternobaric Vertigo: Incidence in Portuguese Air Force Pilots

Ten/med. João Subtil; Maj/med. Jorge Varandas; LtCol/med. Alves dos Santos

Aeronautical Medical Center, ENT department

Portuguese Air Force

Azinhaga da Torre do Fato

1600 Lisbon, Portugal

Abstract

Alternobaric Vertigo is a sudden and transient vertigo caused by asymmetrical changes in middle ear pressure (Lündergren – 1965). These changes are directly related to the daily activity of aircraft pilots (and divers). This vertigo strikes often in critical maneuvers (attacks or evasions) causing sudden disorientation. We used an anonymous enquiry, multiple-choice type, preceded by a briefing about alternobaric vertigo. We found 29% of the pilots with 1 or more episodes of alternobaric vertigo, all being caused directly by rapid ascents and/or brisk Valsalva maneuvers.

Introduction

Alternobaric Vertigo is a sudden and transient vertigo caused by asymmetrical changes in middle ear pressure (Lündergren – 1965). These changes are directly related to the daily activity of aircraft pilots (and divers), which implies variation of altitude (or depth). This vertigo typically lasts few seconds, with prompt *restitutio ad integrum*, and no organ damage is observed. The precise physiopathologic mechanism is not known, even though several studies have been made searching for circulatory, thermal or pressure changes in internal ear, as well as anatomical variations.

Its importance derives from the fact that it strikes often in critical maneuvers (attacks or evasions) causing sudden disorientation.

Known in divers since 1896, and observed in pilots since 1937, was first coined “Alternobaric Vertigo” by Lündergren in 1965. Probably (predictably) fairly common in high-performance aircrafts, as well as in divers:

RAF pilots 10% - Jones, 1956⁽³⁾

Swedish Air Force Pilots 17% - Lündergren 1966⁽⁴⁾

Divers 26% Lündergren 1965⁽⁴⁾

It is assumable that pilots are reluctant to report such symptoms and so these data are most likely underestimated...

The importance of this situation drove our Aeronautical Medical Center to enquire our high performance pilots, to get *au pair* of our numbers, as well as to learn more about how to prevent and deal with it.

Materials and Methods

It was used an anonymous enquiry, multiple-choice type, preceded by a *briefing* about alternobaric vertigo. We included questions about specific details of the episode, as well as risk and precipitating factors.

This quiz was filled by 12 Alphajet and 12 F16 pilots (about 2/3 of our pilots flying each of these aircrafts)

Results

More than predicted by previous studies, we found 7 pilots with 1 or more episodes of alternobaric vertigo (29%): 3 in F16, 2 in Alphajet, 1 in Epsilon and 1 on the floor prior to flying. In 5 cases there were more than 1 episode. We were unable to determine a relation between the number of episodes and the total flown hours; even though, 5 pilots have more than 1000h. No incidents were officially reported.

Characteristics:

In most cases, the vertigo was rotary horizontal, with 2 episodes of sagittal vertigo. All but 1 episode were not accompanied by other symptoms, the exception being tinnitus.

Precipitating factors

All the episodes were precipitated either by rapid ascent or by Valsalva maneuver.

Risk factors

About 71% (5) of the episodes were connected with recent or contemporary upper respiratory tract infection. 1 pilot has history allergy with predominant terrain in the upper respiratory tract. 59% (4) of the referred pilots are smokers (14% light smoker, 28% moderate and 14% heavy). 28% (2) mentioned chronic recurrent upper respiratory tract afflictions (sinusitis, rhinitis...). Only 1 pilot had no identified risk factors. None of our study pilots had had adenoidectomy.

Discussion

Even though we had few pilots to enquiry (24), the numbers were impressive in two ways: the problem seems more common than expected, probably because of the increased performance of the aircrafts since the studies presented, in spite of the fact the selection methods are more careful nowadays. The second striking aspect was the role clearly present for the precipitating factors, having all the episodes being caused directly by rapid ascents and/or brisk Valsalva maneuvers, demonstrating the critical role of pressure changes in this type of vertigo. Confirming this datum are the risk factors that were almost omnipresent.

As this vertigo is typically brief and self-limited, one can only prevent it and limit its consequences:

- Learn to promptly recognize it and deal with it properly – leveling the flight, limiting the pressure changes.
- Avoid flying with active risk factors – UTI, active allergy...
- Avoid smoking and other respiratory irritants.
- Avoid precipitating factors – make frequent and progressive Valsalva maneuvers, especially when flying low level or 100% O₂.
- Pilots should pay frequent visits to the flight surgeon, to rule out any Eustachian tube dysfunction
- Flight surgeons should also keep pilots on continuous formation, including this subject in physiology refreshment courses.

Finally we remind that Alternobaric Vertigo is (as we've seen) an important cause of sudden spatial disorientation, leading to sudden incapacity in flight that cannot be omitted when investigating causes for aircraft accidents.

It is also obvious that it is also important to divers, for the same mechanism(s) inflicts the same pressure changes in the middle ear.

The fact of being brief and self-limited should not minimize it's importance in our minds. It can occur at any critical time!

References

1. Wicks, RE. Alternobaric Vertigo: An Aeromedical Review. ASEM. 1989; 60:67-72
2. Armstrong HG, Heim JW. The effect of flying on the middle ear. JAMA. 1937; 109:417-21
3. Jones MG. Review of current problems associated with disorientation in man controlled flight. FPRC, RAF, GB, Oct 1957
4. Lundgren CEG, Malm LU. Alternobaric vertigo among pilots. Aerospace Med. 1966
5. Enders LJ Lopez ER. Aeromedical consultation case report: alternobaric vertigo. Aerospace Med. 41(2):200-202. 1970
6. Lundgren CEG Alternobaric vertigo: a diving hazard Brit. Med. J., 1965
7. Benson AJ. Textbook of aviation Physiology. Pergamon Press. 1965
8. Boies et al. Fundamentals of Otolaryngology. Saunders. 1964.
9. Fatori B. et al. Alternobaric and hyperbaric oxygen therapy in the immediate and long term treatment of Meniere's disease. Audiology 1996 Nov-Dec;35(6):322-34
10. Suzuki M. et al. The influence of vestibular and cochlear aqueducts on vestibular response to middle ear pressure changes in guinea pigs. Acta Otolaryngologica 1994;Sup510:16-9
11. Suzuki M. et al. Involvement of round and oval windows in the vestibular response to pressure changes in the middle ear of guinea pigs. Acta Otolaryngologica 1998;118:712-716

12. Nagai H. et al. Effect of increased middle ear pressure on blood flow to the middle ear, inner ear and facial nerve in guinea pigs. *Acta Otolaryngologica* 1996;116:439-442
13. Tjernstrom O Further studies on alternobaric vertigo. *Acta Otolaryngologica* 1974;78:221-231
14. Tjernstrom O Middle ear mechanics and alternobaric vertigo. *Acta Otolaryngologica* 1974;78:376-384
15. Suzuki M. et al. The influence of rates of pressure change on pressure-induced vestibular response in guinea pigs. *Nippon Jibiinkoka Gakkai Kaiho* 1995 May; 98 (5):820-4
16. Molvaer OI et al. Alternobaric vertigo in professional divers. *Undersea Biomed Res* 1998 Jul;15(4):271-82
17. Molvaer OI et al. Facial baroparesis: a review. *Undersea Biomed Res* 1987 May;14(3):277-95
18. Densert O et al. Pressure-regulating mechanisms in the inner ear. *ORL J Otorhinolaryngol Relat Spec* 1979;40(6):319-24
19. Carlborg BI et al. Transmission of cerebrospinal fluid pressure via the cochlear aqueduct and endolymphatic sac. *Am J Otolaryngol* 1983 Jul-Aug;4(4):273-82
20. Ivarson A. et al. Volume-pressure properties of round and oval windows. A quantitative study on human temporal bone. *Acta Otolaryngol (Stockh)* 1977 Jul-Aug;84(1-2):38-43
21. Suzuki M. et al. The influence of vestibular and cochlear aqueducts on vestibular response to middle ear pressure changes in guinea pigs. *Acta Otolaryngologica Suppl (Stockh)* 1994;510:16-9
22. Suzuki M. et al. The influence of middle ear pressure changes on vestibular neurons in guinea pigs. *Acta Otolaryngologica Suppl (Stockh)* 1994;510:9-15
23. Konradson KS et al. Perilymph pressure during hypobaric conditions -cochlear aqueduct obstructed. *Acta Otolaryngologica (Stockh)* 1994 Jan;114(1):24-9
24. Kossowski M, Coulet O, Florentin JL, Bonete D, Gauvin Y, Bonne L, Cohat JP. Incidence of Vertigo in Diving. *Rev Laryngol Otol Rhinol (Bord)* 1997; 118(5):301-5

Address:

Ten. / Med. João Subtil
Portuguese Ministry of Defense - Portuguese Air Force
Aeronautical Medicine Center
Azinhaga da Torre do Fato, 1600
Lisboa - Portugal

E-mail:

jsubtil@netcabo.pt

This page has been deliberately left blank



Page intentionnellement blanche