

UNCLASSIFIED

Defense Technical Information Center  
Compilation Part Notice

ADP011665

TITLE: Effects of Anisotropy in Light Scattering by Anisotropic Layer  
Around a Spherical Particle in Uniaxial Medium

DISTRIBUTION: Approved for public release, distribution unlimited

This paper is part of the following report:

TITLE: International Conference on Electromagnetics of Complex Media  
[8th], Held in Lisbon, Portugal on 27-29 September 2000. Bianisotropics  
2000

To order the complete compilation report, use: ADA398724

The component part is provided here to allow users access to individually authored sections  
of proceedings, annals, symposia, etc. However, the component should be considered within  
the context of the overall compilation report and not as a stand-alone technical report.

The following component part numbers comprise the compilation report:

ADP011588 thru ADP011680

UNCLASSIFIED

# Effects of Anisotropy in Light Scattering by Anisotropic Layer around a Spherical Particle in Uniaxial Medium

A. D. Kiselev<sup>1</sup>, V. Yu. Reshetnyak<sup>2</sup>, and T. J. Sluckin<sup>3</sup>

<sup>1</sup> Department of Pure & Applied Mathematics,  
Chernigov State Technological University,  
Shevchenko Street 95, 14027 Chernigov, Ukraine  
Fax: +380-46-22 34244; Email: kisel@elit.chernigov.ua

<sup>2</sup> Department of Theoretical Physics, Kiev State University,  
Prospect Glushkova 6, 03680 Kiev, Ukraine

<sup>3</sup> Faculty of Mathematical Studies, University of Southampton,  
Southampton SO17 1BJ, United Kingdom

## Abstract

We have used a Mie-type theory to study the light scattering from an annular anisotropic layer around a spherical colloidal particle. We have derived an exact solution of the scattering problem in the case when the distribution of the optical axes around the particles possesses some special transformation properties under rotation and, outside of the layer, the ambient medium is isotropic. We have then calculated the dependence of the scattering cross-section on particle size, anisotropy parameter, and layer thickness for different optical axis distributions. We find that the scattering cross-section is strongly affected by the type of anisotropy. The presence of disclinations enhances scattering efficiency. We determine the region of validity of Rayleigh-Gans approximation by comparing approximate values of the scattering cross-section with the results computed from the exact solution. As an additional effect specific to anisotropic scatterer, it is found that for structures with broken central symmetry there is the phase shift proportional to the logarithm of layer thickness that enters the scattering amplitudes.

In order to study the case of anisotropic ambient medium, approximate theory has been developed. The phase shift is found to affect the scattering amplitudes even if the central symmetry is unbroken.

## 1. Introduction

There are a large number of physical contexts in which it is useful to understand light scattering by impurities [1]. A particular example of recent interest concerns liquid crystal devices. There are now a number of systems in which liquid crystal droplets are suspended in a polymer matrix – the so-called PDLC systems – or the inverse system, involving colloids now with a nematic liquid crystal solvent. The latter systems are commonly known as filled nematics [2, 3].

In such systems it is required to calculate the scattering of light by composite anisotropic particles embedded in an isotropic matrix. In this paper we discuss some model cases of light scattering by such particles, which may be supposed to represent local liquid crystalline director structures, using both the Mie and the Rayleigh-Gans (R-G) approaches.

The large majority of exact solutions of the single-scattering problem have been derived for isotropic scatterers [4]. However, there are a few cases for which Mie theory has been extended to the case of anisotropic scatterers [5, 6, 7, 8].

In order that the problem have an analytic solution, we find it necessary to restrict consideration to cases in which the optical axis distributions within the anisotropic layer around the core possess some special symmetry properties under rotation. The analysis is then based on a systematic expansion of the electromagnetic field over vector spherical harmonics [9, 10]. The specific form of the expansions is known as the  $T$ -matrix ansatz. This has been widely used in the related problem of light scattering by nonspherical particles [11].

In this study we investigate the dependence of light scattering from the layer on the internal structure of the optical tensor, and implicitly on the liquid crystal director texture. We also compare our exact results with those obtained by the simpler but less accurate R-G method. For brevity, in what follows we shall leave aside the details on this comparison and do not discuss extensions of the theory.

## 2. Scatterers

We consider scattering by a spherical particle of radius  $R_1$  embedded in a uniform isotropic dielectric medium with dielectric constant  $\epsilon_{ij} = \epsilon \delta_{ij}$  and magnetic permeability  $\mu_{ij} = \mu \delta_{ij}$ . The scattering particle consists of an inner isotropic core of radius  $R_2$ , surrounded by an anisotropic annular layer of thickness  $d = R_1 - R_2$ .

Within the inner core of the scatterer the dielectric tensor  $\epsilon$ , and the magnetic permittivity  $\mu$  take the values  $\epsilon_{ij} = \epsilon_2 \delta_{ij}$ ,  $\mu_{ij} = \mu_2 \delta_{ij}$ . The dielectric tensor within the annular layer is locally uniaxial. The optical axis distribution is defined by the vector field  $\hat{\mathbf{n}}$ . (Hats will denote unit vectors.) Then within the annular layer  $\epsilon_{ij}(\mathbf{r}) = \epsilon_1 \delta_{ij} + \Delta \epsilon_1 (\hat{\mathbf{n}}(\mathbf{r}) \otimes \hat{\mathbf{n}}(\mathbf{r}))_{ij}$  and  $\mu_{ij} = \mu_1 \delta_{ij}$ . The unit vector  $\hat{\mathbf{n}}$  corresponds to a liquid crystal director for material within the annular region.

We shall suppose that the director field can be written in the following form

$$\hat{\mathbf{n}} = n_r \hat{\mathbf{r}} + n_\varphi \hat{\boldsymbol{\varphi}} + n_\vartheta \hat{\boldsymbol{\vartheta}}, \quad (1)$$

where  $\hat{\mathbf{r}}$  is the unit radial vector and  $\hat{\boldsymbol{\varphi}}$ ,  $\hat{\boldsymbol{\vartheta}}$  are the vectors tangential to the unit sphere. The components  $n_r, n_\varphi, n_\vartheta$  are constants.

## 3. Generalized Mie Theory

The electromagnetic field can be expanded using the vector spherical harmonic basis,  $\mathbf{Y}_{j+\delta jm}(\phi, \theta) \equiv \mathbf{Y}_{j+\delta jm}(\hat{\mathbf{r}})$  ( $\delta = 0, \pm 1$ ) [9] as follows:

$$\mathbf{E} = \sum_{jm} \mathbf{E}_{jm} = \sum_{jm} \left[ p_{jm}^{(0)}(r) \mathbf{Y}_{jm}^{(0)}(\hat{\mathbf{r}}) + p_{jm}^{(e)}(r) \mathbf{Y}_{jm}^{(e)}(\hat{\mathbf{r}}) + p_{jm}^{(m)}(r) \mathbf{Y}_{jm}^{(m)}(\hat{\mathbf{r}}) \right], \quad (2a)$$

$$\mathbf{H} = \sum_{jm} \mathbf{H}_{jm} = \sum_{jm} \left[ q_{jm}^{(0)}(r) \mathbf{Y}_{jm}^{(0)}(\hat{\mathbf{r}}) + q_{jm}^{(e)}(r) \mathbf{Y}_{jm}^{(e)}(\hat{\mathbf{r}}) + q_{jm}^{(m)}(r) \mathbf{Y}_{jm}^{(m)}(\hat{\mathbf{r}}) \right], \quad (2b)$$

where  $\mathbf{Y}_{jm}^{(m)} = \mathbf{Y}_{jjm}$  and  $\mathbf{Y}_{jm}^{(e)} = [j/(2j+1)]^{1/2} \mathbf{Y}_{j+1jm} + [(j+1)/(2j+1)]^{1/2} \mathbf{Y}_{j-1jm}$  are electrical and magnetic harmonics respectively, and  $\mathbf{Y}_{jm}^{(0)} = [j/(2j+1)]^{1/2} \mathbf{Y}_{j-1jm} - [(j+1)/(2j+1)]^{1/2} \mathbf{Y}_{j+1jm}$  are longitudinal harmonics.

Outside the scatterer  $p_{jm}^{(\alpha)}(r)$  and  $q_{jm}^{(\alpha)}(r)$  can be expressed in terms of linear combinations of spherical Bessel functions and spherical Hankel functions. For the incident and scattered waves

in the far field region we have:

$$\mathbf{E}_{inc} = \sum_{jm} \left( \alpha_{jm}^{(inc)} j_j(\rho) \mathbf{Y}_{jm}^{(m)} + \tilde{\alpha}_{jm}^{(inc)} D j_j(\rho) \mathbf{Y}_{jm}^{(e)} \right), \quad (3)$$

$$\mathbf{E}_{sca} = \sum_{jm} \left( \beta_{jm}^{(sca)} h_j^{(1)}(\rho) \mathbf{Y}_{jm}^{(m)} + \tilde{\beta}_{jm}^{(sca)} D h_j^{(1)}(\rho) \mathbf{Y}_{jm}^{(e)} \right), \quad (4)$$

where  $\rho = kr$  and  $Df(x) \equiv x^{-1} \frac{d}{dx}(xf(x))$ . Since the scattering problem is linear, the coefficients  $\{\beta_{jm}^{(sca)}, \tilde{\beta}_{jm}^{(sca)}\}$  are linearly related to  $\{\alpha_{jm}^{(inc)}, \tilde{\alpha}_{jm}^{(inc)}\}$  through the elements of the  $T$ -matrix.

In order to derive the  $T$ -matrix, we need to find the general expressions for the electromagnetic field inside the anisotropic layer and the isotropic core. The ingoing and outgoing waves then can be related by using continuity of the tangential components of the electric and magnetic fields at  $r = R_2$  and  $r = R_1$  as boundary conditions.

Substituting the expansions (2) into the Maxwell equations gives a system of equations for the components  $p_{jm}^{(\alpha)}$  and  $q_{jm}^{(\alpha)}$ . For the distributions (1) we have found that the system can be reduced to a system of coupled Bessel equations for the magnetic components:  $\{p_{jm}^{(m)}, q_{jm}^{(m)}\}$ . For the structures  $\hat{\mathbf{n}} = \hat{\boldsymbol{\vartheta}}$  and  $\hat{\mathbf{n}} = \cos \gamma \hat{\mathbf{r}} + \sin \gamma \hat{\boldsymbol{\varphi}}$  solutions of this system can be obtained in the closed form. The result for transverse components is

$$p_{jm}^{(m)} = \alpha_{jm} j_j(\rho_e) + \beta_{jm} h_j^{(1)}(\rho_e), \quad q_{jm}^{(e)} = n_e / \mu_1 D p_{jm}^{(m)}(\rho_e), \quad (5a)$$

$$q_{jm}^{(m)} \equiv \tilde{\rho}_e^{i\delta_j} \tilde{q}_{jm}^{(m)} = \tilde{\rho}_e^{i\delta_j} \left[ \tilde{\alpha}_{jm} j_{\tilde{j}}(\tilde{\rho}_e) + \tilde{\beta}_{jm} h_{\tilde{j}}^{(1)}(\tilde{\rho}_e) \right], \quad p_{jm}^{(e)} = -\mu_1 / \tilde{n}_e \tilde{\rho}_e^{i\delta_j} D \tilde{q}_{jm}^{(m)}(\tilde{\rho}_e), \quad (5b)$$

$$\tilde{j}(\tilde{j} + 1) = \frac{1 + u_1(1 - n_{\tilde{\theta}}^2)}{(1 + u_1 n_r^2)^2} j(j + 1), \quad \delta_j = \tilde{u}_1 \tilde{m}_e^2 \sqrt{j(j + 1)} n_{\varphi} n_r, \quad (5c)$$

$$u_1 \equiv \Delta \epsilon_1 / \epsilon_1, \quad m_e^2 = \frac{1 + u_1}{1 + u_1(1 - n_{\tilde{\theta}}^2)}, \quad \tilde{m}_e^2 = \frac{1 + u_1(1 - n_{\tilde{\theta}}^2)}{1 + u_1 n_r^2}, \quad (5d)$$

where  $\alpha_{jm}$  and  $\beta_{jm}$  are integration constants;  $\rho_e = m_e \rho_1 = m_e k_1 r$  and  $\tilde{\rho}_e = \tilde{m}_e \rho_1$ . For brevity, the corresponding rather cumbersome expressions for the  $T$ -matrix have been omitted. It is seen, that anisotropy affects the analytical expressions in the following manner: (a) it renormalizes the order of corresponding Bessel functions; (b) it changes the arguments of the functions by replacing  $k_1$  with  $k_e$ ; (c) it leads to the appearance of the geometric factor  $\propto \rho^{i\delta_j}$  for the tilted configuration.

#### 4. Scattering Efficiency

In this section we discuss briefly the results concerning the scattering efficiency,  $Q$ , that is the ratio of the total scattering cross section  $C_{sca}$  and area of the composite particle,  $S = \pi R_1^2$ .

Dependence of the scattering efficiency  $Q$  on the dimensionless size parameter  $kR_2$ , is depicted in Figs. 1. It is assumed that refractive indexes of the surrounding medium and the isotropic core are equal,  $n = n_2$ . Clearly, the scattering cross section is strongly affected by the type of anisotropy characterizing by the unit vector  $\hat{\mathbf{n}}$ . Fig. 1 shows that the structure  $\hat{\mathbf{n}} = \hat{\boldsymbol{\varphi}}$  has the largest value of  $Q$  at small  $kR_2$ , whereas the least scattering efficiency corresponds to the configuration with  $\hat{\mathbf{n}} = \hat{\boldsymbol{\vartheta}}$ . We found that the helical structure  $\hat{\mathbf{n}} = \hat{\boldsymbol{\varphi}}$  remains the most efficient scatterer with increase in the anisotropy parameter  $u_1$  for sufficiently small particle size. On the other hand, Fig. 1 reveals that the configuration  $\hat{\mathbf{n}} = \hat{\boldsymbol{\vartheta}}$  becomes the most efficient scatterer as size of the scatterer increases.

#### Acknowledgement

This work is performed under INTAS grant 97-1315.




FIGURE 1: Dependence of  $Q$  on the size parameter  $kR_2$  at  $u_1 = 0.5$  and  $d/R_2 = 0.2$  for different types of anisotropy. It is shown that the scattering efficiency reveals crossover behaviour in this range of size parameters.

## References

- [1] A. Ishimaru, *Wave Propagation and Scattering in Random Media*. London: Academic Press, 1978.
- [2] M. Kreuzer and R. Eidenschink, in *Liquid Crystals in Complex Geometries*, G. P. Crawford and S. Žumer (Eds.), Chap. 15. London: Taylor & Francis, 1996.
- [3] T. Bellini, N. A. Clark, V. Degiorgio, F. Mantegazza, and G. Natale, "Light scattering measurement of the nematic correlation length in a liquid crystal with quenched disorder," *Phys. Rev. E.*, vol. 57, pp. 2996-3006, 1998.
- [4] C. F. Boren and D. R. Huffman, *Absorption and Scattering of Light by Small Particles*. New York: Wiley, 1983.
- [5] J. Roth and M. J. Digman, "Scattering and extinction cross sections for a spherical particle coated with an oriented molecular layer," *JOSA*, vol. 63, pp. 308-311, 1973.
- [6] B. Lange and S. R. Aragon, "Mie scattering from anisotropic thin spherical shells," *J. Chem. Phys.*, vol. 92, pp. 4643-4650, 1990.
- [7] D. K. Hahn and S. R. Aragon, "Mie scattering from anisotropic thick spherical shells," *J. Chem. Phys.*, vol. 101, pp. 8409-8417, 1994.
- [8] H. Karacali, S. M. Risseler, and K. F. Ferris, "Scattering of light from small nematic spheres with radial dielectric anisotropy," *Phys. Rev. E.*, vol. 56, pp. 4286-4293, 1997.
- [9] L. C. Biedenharn and J. D. Louck, *Angular Momentum in Quantum Physics*. Massachusetts: Addison-Wesley, 1981.
- [10] R. G. Newton, *Scattering Theory of Waves and Particles*. London: McGraw Hill, 1966.
- [11] M. I. Mishchenko, L. D. Travis, and D. W. Mackowski, "T-matrix computations of light scattering by nonspherical particles: A review," *J. of Quant. Spectr. & Radiat. Transf.*, vol. 55, pp. 535-575, 1996.