

THIS REPORT HAS BEEN DELIMITED  
AND CLEARED FOR PUBLIC RELEASE  
UNDER DOD DIRECTIVE 5200.20 AND  
NO RESTRICTIONS ARE IMPOSED UPON  
ITS USE AND DISCLOSURE.

DISTRIBUTION STATEMENT A

APPROVED FOR PUBLIC RELEASE;  
DISTRIBUTION UNLIMITED.

# Services Technical Information Agency

ur limited supply, you are requested to return this copy WHEN IT HAS SERVED  
OSE so that it may be made available to other requesters. Your cooperation  
ciated.

# 1238

OR OTHER DRAWINGS, SPECIFICATIONS OR OTHER DATA  
HER THAN IN CONNECTION WITH A DEFINITELY RELATED  
OPERATION, THE U. S. GOVERNMENT THEREBY INCURS  
OBLIGATION WHATSOEVER; AND THE FACT THAT THE  
MULATED, FURNISHED, OR IN ANY WAY SUPPLIED THE  
NS, OR OTHER DATA IS NOT TO BE REGARDED BY  
AS IN ANY MANNER LICENSING THE HOLDER OR ANY OTHER  
INVENTION THAT MAY IN ANY WAY BE RELATED THERETO.

Reproduced by  
ENT SERVICE CENTER  
T BUILDING, DAYTON, 2, OHIO

# CLASSIFIED

AD No. 40238

ASTIA FILE COPY

U. S. NAVY  
OFFICE OF NAVAL RESEARCH  
Research Contract N7onr-32912  
Project NR 092 162

STUDIES ON THE PHOTOCHEMICAL SYNTHESIS OF HYDRAZINE  
AND OTHER ENDOTHERMIC COMPOUNDS

Status Report  
for the period 15 April to 15 June 1954

by  
Harry E. Gunning  
with  
C. Luner and C. C. McDonald

Illinois Institute of Technology  
Department of Chemistry  
Technology Center  
3300 Federal Street  
Chicago 16, Illinois

STUDIES ON THE PHOTOCHEMICAL SYNTHESIS OF HYDRAZINE  
AND OTHER ENDOTHERMIC COMPOUNDS

Status Report

for the period 15 April to 15 June 1954

A. THE PHOTOCHEMICAL SYNTHESIS OF HYDRAZINE FROM AMMONIA

The results of our investigations of the photolysis of ammonia at 1849 Å under both static and dynamic conditions were submitted to the Journal of Chemical Physics some time ago. The paper appeared in the May 1954 issue of the journal.

A manuscript covering the salient features of our studies of the mechanism of hydrazine formation in the reaction of ammonia with Hg  $6(^3P_1)$  atoms has been prepared. It will be submitted shortly to the Journal of Chemical Physics.

B. THE MERCURY- $6(^3P_1)$ -PHOTOSENSITIZED DECOMPOSITION OF ETHYLENEIMINE

This reaction has been studied at 25°C under circulating conditions. The principal products of the reaction have been shown to be ethylene, nitrogen and hydrogen. Small amounts of butanes, methane and a polymeric material have also been isolated. The analyses were largely carried out on the mass spectrometer. The rates of formation, and the quantum yields for the principal products of the reaction have been determined as functions of both substrate pressure and duration of exposure. At the present time, the data are in process of evaluation toward the development of a mechanism for the reaction. As soon as a reasonable mechanism can be formulated, the results of the study will be submitted to ONR as a technical report.

C. THE PHOTOOXIDATION OF CHLORATE TO PERCHLORATE

A number of experiments have been performed on the irradiation of suspensions of zinc oxide in dilute potassium chlorate solutions in the

presence of oxygen. The reactions were carried out in Pyrex vessels perfused with the radiation from a quartz mercury arc. Since the Pyrex absorbs all wavelengths shorter than 3200 Å, and the zinc oxide does not absorb above 4000 Å, the actinic energy can be considered as essentially arising from the strong 3660 Å group from the arc. A major problem encountered early in the investigation was the analysis for perchlorate in the presence of chlorate. At the present time, our efforts are being devoted exclusively to the analytical aspects of the problem. It is expected that the technique used will shortly be refined sufficiently to enable quantitative determination of the small amounts of perchlorate ion present after irradiation. Thereafter the kinetic studies will be resumed.

#### D. THE PHOTODECOMPOSITION OF TITANIUM TETRACHLORIDE

A flexible high-vacuum reaction assembly has now been constructed for the study of the photodecomposition of titanium tetrachloride, in the gas phase, under flow conditions. Mercury vapor will be used as a getter for chlorine atoms in the reaction. Analytical methods for quantitative determination of the lower halides of titanium are also being explored.

<u>DISTRIBUTION</u>	Director, Office of Naval Research Chicago Branch Office 86 East Randolph Street Chicago 1, Illinois	2 copies
	Chief of Naval Research Navy Department Washington 25, D. C. Attention: Code 429	5 copies
	Illinois Institute of Technology	8 copies

# Armed Services Technical Information Agency

Because of our limited supply, you are requested to return this copy WHEN IT HAS SERVED YOUR PURPOSE so that it may be made available to other requesters. Your cooperation will be appreciated.

# AD

# 40238

NOTICE: WHEN GOVERNMENT OR OTHER DRAWINGS, SPECIFICATIONS OR OTHER DATA ARE USED FOR ANY PURPOSE OTHER THAN IN CONNECTION WITH A DEFINITELY RELATED GOVERNMENT PROCUREMENT OPERATION, THE U. S. GOVERNMENT THEREBY INCURS NO RESPONSIBILITY, NOR ANY OBLIGATION WHATSOEVER; AND THE FACT THAT THE GOVERNMENT MAY HAVE FORMULATED, FURNISHED, OR IN ANY WAY SUPPLIED THE SAID DRAWINGS, SPECIFICATIONS, OR OTHER DATA IS NOT TO BE REGARDED BY IMPLICATION OR OTHERWISE AS IN ANY MANNER LICENSING THE HOLDER OR ANY OTHER PERSON OR CORPORATION, OR CONVEYING ANY RIGHTS OR PERMISSION TO MANUFACTURE, USE OR SELL ANY PATENTED INVENTION THAT MAY IN ANY WAY BE RELATED THERETO.

Reproduced by  
**DOCUMENT SERVICE CENTER**  
KNOTT BUILDING, DAYTON, 2, OHIO

# UNCLASSIFIED