

THIS REPORT HAS BEEN DELIMITED
AND CLEARED FOR PUBLIC RELEASE
UNDER DOD DIRECTIVE 5200.20 AND
NO RESTRICTIONS ARE IMPOSED UPON
ITS USE AND DISCLOSURE.

DISTRIBUTION STATEMENT A

APPROVED FOR PUBLIC RELEASE,
DISTRIBUTION UNLIMITED.

Armed Services Technical Information Agency

Because of our limited supply, you are requested to return this copy WHEN IT HAS SERVED YOUR PURPOSE so that it may be made available to other requesters. Your cooperation will be appreciated.

AD

42475

NOTICE: WHEN GOVERNMENT OR OTHER DRAWINGS, SPECIFICATIONS OR OTHER DATA ARE USED FOR ANY PURPOSE OTHER THAN IN CONNECTION WITH A DEFINITELY RELATED GOVERNMENT PROCUREMENT OPERATION, THE U. S. GOVERNMENT THEREBY INCURS NO RESPONSIBILITY, NOR ANY OBLIGATION WHATSOEVER; AND THE FACT THAT THE GOVERNMENT MAY HAVE FORMULATED, FURNISHED, OR IN ANY WAY SUPPLIED THE SAID DRAWINGS, SPECIFICATIONS, OR OTHER DATA IS NOT TO BE REGARDED BY IMPLICATION OR OTHERWISE AS IN ANY MANNER LICENSING THE HOLDER OR ANY OTHER PERSON OR CORPORATION, OR CONVEYING ANY RIGHTS OR PERMISSION TO MANUFACTURE, USE OR SELL ANY PATENTED INVENTION THAT MAY IN ANY WAY BE RELATED THERETO.

Reproduced by
DOCUMENT SERVICE CENTER
KNOTT BUILDING, DAYTON, 2, OHIO

UNCLASSIFIED

AD No. 42475
ASTIA FILE COPY

OFFICE OF NAVAL RESEARCH

Task Contract N5ori-07822

Project Designation No. NR 356-096

Technical Report No. 21

CYCLIC POLYOLEFINS. XXXIII.
COMPOUNDS DERIVED FROM CYCLO⁸OCTATETRAENYLMETHYL ALCOHOL.
ISOMERIZATION OF CYCLOOCTATETRAENYLACETONITRILE

by

Arthur C. Cope, Ronald M. Pike and Donald F. Rugen

(Prepared for publication in the
Journal of the American Chemical Society)

Massachusetts Institute of Technology
Department of Chemistry
Cambridge 39, Mass.

July 27, 1954

[Contribution from the Department of Chemistry,
Massachusetts Institute of Technology]

CYCLIC POLYOLEFINS. XXXIII. COMPOUNDS DERIVED FROM
CYCLOOCTATETRAENYLMETHYL ALCOHOL. ISOMERIZATION OF
CYCLOOCTATETRAENYLACETONITRILE¹

(1) Supported in part by the Office of Naval Research under contract
N5ori-07822, Project Designation NR356-096.

By Arthur C. Cope, Ronald M. Pike and Donald F. Rugen

Received _____

N,N-Dimethyl cyclooctatetraenylmethylaniline (II) and
cyclooctatetraenylacetone (III) have been prepared by displacement reactions
from cyclooctatetraenylmethyl bromide (I). The nitrile III was converted into
an isomer (VIII or IX) by migration of one of the nuclear double bonds into
conjugation with the nitrile group on heating with triethylamine, or on treat-
ment with sodium hydroxide at room temperature. Both of the nitriles were
converted to amides by treatment with alkaline hydrogen peroxide.

This paper reports an investigation of several cyclooctatetraene
derivatives prepared from cyclooctatetraenylmethyl alcohol, an allylic-type
alcohol obtained by copolymerization of acetylene with propargyl alcohol.²

(2) A. C. Cope and D. F. Rugen, This Journal, 75, 3215 (1953).

Attempts to prepare the p-toluenesulfonate of cyclooctatetraenyl-


methyl alcohol under conditions used for other alcohols in the cyclooctatetraene series^{2,3} and to use the ester as an intermediate in displacement reactions

(3) A. C. Cope and R. M. Pike, *ibid.*, 75, 3220 (1953).

gave erratic results, probably because of side reactions between the reactive allylic-type tosylate and the base (usually pyridine) employed in its preparation. It was possible to prepare cyclooctatetraenylmethyl bromide from the alcohol and phosphorus tribromide in hexane in the presence of pyridine, however, and the pure bromide I was isolated by distillation in 65% yield. Reaction of the bromide with dimethylamine in benzene yielded N,N-dimethylcyclooctatetraenylmethylamine (II, 43%), which was characterized as the crystalline picrate and hydrochloride. Cyclooctatetraenylacetonitrile (III) also was prepared from the bromide by reaction with potassium cyanide in aqueous tetrahydrofuran in the presence of potassium iodide. The nitrile III was purified and isolated in 44% yield by conversion into a non-crystalline silver nitrate complex, followed by regeneration with ammonium hydroxide and chromatography on silica gel. Infrared spectra of the bromide, tertiary amine and nitrile (Fig. 1) contain the bands previously noted as characteristic of monosubstituted cyclooctatetraenes,² and accordingly provide evidence supporting the structures of these compounds. In addition, the spectrum of the nitrile III contains a sharp band at 4.4μ characteristic of the (unconjugated) nitrile group.

Reaction of cyclooctatetraenylacetonitrile with aqueous sodium hydroxide at room temperature did not yield the carboxylic acid, but resulted in partial isomerization forming a nitrile with an infrared absorption band at 4.48μ , indicating that the nitrile group was conjugated. Isomerization of the

nitrile III was most readily accomplished by heating under reflux with triethylamine in a nitrogen atmosphere for seventy hours, at which time the nitrile band of III at 4.4μ had disappeared completely and the pure isomer with a nitrile band at 4.48μ was isolated in 37% yield. In the isomerization, migration of an α -hydrogen occurs and a double bond shifts from the cyclooctatetraene nucleus of III into conjugation with the nitrile group. Formula VIII represents the structure that would be formed by a three-carbon tautomeric shift, while the cross-conjugated structure IX could be formed by a second three-carbon shift. In addition to differences between the infrared spectra of the isomeric nitriles (Fig. 1), the conjugated isomer has two strong maxima at 314 and $242.5\text{ m}\mu$ in the ultraviolet spectrum (Fig. 2), while cyclooctatetraenylacetonitrile, like other cyclooctatetraene derivatives,² shows weak ultraviolet absorption becoming strong near $230\text{ m}\mu$. The existence of two maxima in the ultraviolet spectrum provides evidence favoring the cross-conjugated structure IX for the nitrile.



I, R = CH_2Br

II, R = $\text{CH}_2\text{N}(\text{CH}_3)_2$


III, R = CH_2CN

IV, R = CH_2CONH_2


V, R = $\text{CH}_2\text{CH}_2\text{CN}$

VI, R = $\text{CH}_2\text{CH}_2\text{CONH}_2$

VII, R = $\text{CH}_2\text{CH}_2\text{COOH}$




VIII, X = CN
X, X = CONH_2



IX, X = CN
XI, X = CONH_2

Since bases isomerized the nitrile III to VIII or IX, an attempt was made using β -cyclooctatetraenylethyl cyanide² (V) as a model compound to find milder conditions of hydrolysis. Reaction of V with dilute hydrogen peroxide in acetone containing a small amount of sodium carbonate yielded the crystalline amide, β -cyclooctatetraenylpropionamide (VI, 36%), which reacted with nitrous acid to form the known β -cyclooctatetraenylpropionic acid (VII). Under these conditions cyclooctatetraenylacetonitrile was converted to cyclooctatetraenylacetamide (IV, 74%), and the isomeric nitrile (VIII or IX) also yielded the corresponding amide (X or XI), which was rather unstable but could be isolated as a crystalline solid. The ultraviolet spectrum of cyclooctatetraenylacetamide is very similar to the spectrum of cyclooctatetraenylacetonitrile and indicates that the cyclooctatetraene nucleus is present in the amide (formula IV). The spectrum of the isomeric amide (X or XI), like the corresponding nitrile, contains two maxima (at 243 and 314 m μ). The ultraviolet spectra do not rule out bridged structures (XII and XIII) for the conjugated nitrile and amide, since 1,3,5-cyclooctatriene and the bridged isomer into which it is partially converted on heating, bicyclo [4.2.0] octa-2,4-diene, have very similar ultraviolet absorption spectra.⁴

(4) A. C. Cope, A. C. Haven, Jr., F. L. Ramp and E. R. Trumbull, ibid., 74, 4867 (1952).



Preliminary attempts to convert cyclooctatetraenylacetamide to cyclooctatetraenylacetic acid with nitrous acid under the conditions used for conversion of VI to VII were unsuccessful, for the product failed to crystallize and was converted to neutral, polymeric material on attempted short path distillation under reduced pressure.

Pharmacological

We are indebted to Dr. Karl H. Beyer and Mr. C. A. Ross of the Sharp and Dohme Division, Merck and Co., Inc., for preliminary pharmacological study of the hydrochloride of the amine II and of N,N-dimethyl- β -cyclooctatetraenylethylamine (V, ref. 2) and γ -cyclooctatetraenyl-n-butylamine (VI, ref. 3). Each compound was injected in the femoral vein of a dog under barbiturate anesthesia with both vagus nerves severed at the cervical level, and carotid blood pressure, respiration and heart rate were recorded, and electrocardiograms were taken. All three compounds produced weak vasopressor responses of moderate duration, but had less than one-thousandth of the vasopressor activity of epinephrine. The phenyl analogs of two of the amines (N,N-dimethylbenzylamine and N,N-dimethyl- β -phenylethylamine) also were tested as the hydrochlorides, and each produced a larger vasopressor response than the corresponding cyclooctatetraene derivative.

Experimental ⁵

-
- (5) Melting points are corrected and boiling points are uncorrected. We are indebted to Dr. S. M. Nagy and his associates for analyses, for the infrared spectra, which were determined with a Baird Double Beam Infrared Recording Spectrometer, Model B, fitted with a sodium chloride prism, and for the ultraviolet spectra, which were determined with a Cary Ultraviolet Recording Spectrophotometer, Model 11 MS.
-

Cyclooctatetraenylmethyl Bromide (I). - Cyclooctatetraenylmethyl alcohol was prepared by copolymerization of propargyl alcohol and acetylene and purified by chromatography on silica gel by procedures described previously. ²

A solution of 0.95 g. of phosphorus tribromide in 15 ml. of dry hexane was added dropwise with stirring to a mixture of 1.34 g. of cyclooctatetraenylmethyl alcohol, 0.27 g. of pyridine and 10 ml. of dry hexane in an atmosphere of nitrogen at 0°. After the addition was complete, the ice bath was removed, and the mixture was stirred for 6 hours at room temperature. The hexane solution was removed by decantation, washed twice with 25-ml. portions of water, and dried over sodium sulfate. After filtration and concentration, distillation of the residue through a semimicro column yielded 1.28 g. (65%) of I, b.p. 75-77.5° (1.6 mm.), n_D^{25} 1.5849-1.5874. A redistilled analytical sample had b.p. 78° (1.6 mm.), n_D^{25} 1.5893, d_4^{25} 1.3634.

Anal. Calcd. for C_8H_3Br : C, 54.88; H, 4.60; Br, 40.57.

Found: C, 55.03; H, 4.73; Br, 40.31.

N,N-Dimethyl Cyclooctatetraenylmethylamine (II). - A solution of 2.43 g. of the bromide I in 30 ml. of benzene was saturated with gaseous dimethylamine and allowed to stand overnight at room temperature. Water (20 ml.) was added, and the benzene was washed with a second 20-ml. portion of water and concentrated under reduced pressure at room temperature. Dry hydrogen chloride was passed through the resulting dry solution of II in a small volume of benzene, and the precipitate of the hydrochloride was separated by filtration. Recrystallization from a mixture of methanol and ether yielded 1.45 g. of the hydrochloride of II, m.p. 209-210.2°. The hydrochloride was added to 15 ml. of 10% sodium hydroxide and the amine II was extracted with two 20-ml. portions of ether. The extracts were diluted with benzene, concentrated under reduced pressure, and the residue was fractionated through a semimicro column. The yield of II was 0.85 g. (43%), b.p. 34-35° (0.1 mm.), n_D^{25} 1.5216-1.5222. An analytical sample had b.p. 35° (0.1 mm.), n_D^{25} 1.5222, d_4^{25} 0.9169.

Anal. Calcd. for $C_{11}H_{15}N$: C, 81.94; H, 9.38; N, 8.68.

Found: C, 82.24; H, 9.34; N, 8.77.

N,N-Dimethyl Cyclooctatetraenylmethanamine Picrate was prepared by adding 0.05 g. of the amine II to a saturated solution of picric acid in ether, and was washed with ether; the yield was 0.06 g., m.p. 145.7-146.2° (capillary inserted at 130°).

Anal. Calcd. for $C_{17}H_{19}N_4O_7$: C, 52.31; H, 4.65; N, 14.35.

Found: C, 52.02; H, 4.87; N, 14.52.

N,N-Dimethyl Cyclooctatetraenylmethanamine Hydrochloride was prepared by passing dry hydrogen chloride through a solution of 0.60 g. of II in dry ether. The precipitate was treated with activated charcoal in a mixture of methanol and ether and crystallized from that solvent; yield 0.41 g., m.p. 209-209.5° (dec., capillary inserted at 195°).

Anal. Calcd. for $C_{11}H_{18}ClN$: C, 66.82; H, 8.16; N, 7.09.

Found: C, 67.12; H, 8.21; N, 7.09.

Cyclooctatetraenylacetonitrile (III). - A solution of 5.42 g. of potassium cyanide and 1.5 g. of potassium iodide in 20 ml. of water was added to a solution of 6.76 g. of the bromide I in 50 ml. of tetrahydrofuran, and the mixture was heated under reflux on a steam bath for 6 hours. The cooled solution was poured into 40 ml. of water, and then extracted with two 30-ml. portions of ether. The extracts were washed with two 25-ml. portions of water, dried over sodium sulfate, and concentrated under reduced pressure. Fractionation through a semimicro column yielded 5.34 g. of the crude nitrile III, b.p. 53-90° (0.06 mm.). A solution of the crude nitrile in 150 ml. of absolute ethanol was heated to boiling, 20.4 g. of powdered silver nitrate was added, and the mixture was heated under reflux until a clear solution was obtained. The

excess silver nitrate was separated by filtering the hot solution, and the filtrate was cooled in an ice bath. The solvent was decanted from the gummy, yellowish green solid that separated, and the solid residue was dissolved in 20 ml. of water. The solution was added to 100 ml. of ice-cold concentrated ammonium hydroxide, which was extracted immediately with two 50-ml. portions of ether. The combined extracts were washed with water, dried over sodium sulfate, and concentrated under reduced pressure, yielding 2.8 g. of a light-yellow liquid residue. This residue was chromatographed on a 13 x 2-cm. column of silica gel,² and the nitrile III was eluted with 200 ml. of a 10% solution of ether in pentane. Concentration and fractionation of the residue through a semimicro column yielded 2.16 g. (44%) of III, b.p. 78-79° (0.10 mm.), n_D^{25} 1.5434-1.5442. An analytical sample had b.p. 79° (0.10 mm.), n_D^{25} 1.5442, d_4^{25} 1.0167.

Anal. Calcd. for $C_{10}H_9N$: C, 83.88; H, 6.34; N, 9.78.

Found: C, 83.84; H, 6.16; N, 9.84.

Cyclooctatetraenylacetamide (IV). - The nitrile III (0.30 g.) was added to 3 ml. of 10% hydrogen peroxide, and acetone was added until a homogeneous solution was obtained, followed by 0.5 ml. of 10% sodium carbonate solution.⁶ The solution was allowed to stand at room temperature for 3

(6) Conditions adapted from a procedure described by J. V. Murray and J. B. Cloke, This Journal, 56, 2749 (1934).

days, after which the acetone was removed with a stream of air and the aqueous suspension was cooled in an ice bath. The pale yellow crystals of IV that separated were recrystallized from a mixture of benzene and 30-60° petroleum

ether, yielding 0.25 g. (7½%) of IV, m.p. 138.6-139.2°. An analytical sample that was recrystallized and sublimed at 0.03 mm. with a heating block temperature of 120-140° had m.p. 138.4-139°.

Anal. Calcd. for $C_{10}H_{11}NO$: C, 74.50; H, 6.83; N, 8.69.

Found: C, 74.79; H, 7.14; N, 8.74.

Attempts to convert the amide IV to cyclooctatetraenylacetic acid by treatment with sodium nitrite and hydrochloric acid under conditions described below for conversion of the next higher homolog VI into the acid VII were unsuccessful. The crude product was a viscous liquid which failed to crystallize and formed a dark, neutral, non-volatile solid on attempted short path distillation, apparently by decarboxylation and polymerization.

β-Cyclooctatetraenylpropionamide (VI). - β-Cyclooctatetraenylethyl cyanide ² (0.30 g.) was treated with alkaline hydrogen peroxide in aqueous acetone for 4 days under the conditions described above for conversion of III to IV. The product was isolated in the same manner as IV, and purified by sublimation at 1.5 mm. with a heating block temperature of 110°. The yield of VI was 0.12 g. (36%), m.p. 61.8-63°. An analytical sample with the same melting point was recrystallized from a mixture of benzene and pentane.

Anal. Calcd. for $C_{11}H_{13}NO$: C, 75.40; H, 7.48; N, 8.00.

Found: C, 75.32; H, 7.49; N, 7.77.

β-Cyclooctatetraenylpropionic Acid (VII) was prepared by adding 1 drop of concentrated hydrochloric acid to a solution of 56.8 mg. of the amide VI and 22.4 mg. of sodium nitrite in 4 ml. of water and 8 ml. of dioxane, and stirring the solution at room temperature for 10 hours. The solution was cooled with ice, made basic by addition of sodium hydroxide, and extracted with benzene. The alkaline aqueous solution was acidified with hydrochloric

acid, and extracted with two 20-ml. portions of ether. The extracts were dried over sodium sulfate, concentrated, and the residue was crystallized from hexane, yielding 20 mg. (35%) of the acid VII, m.p. and mixed m.p. with an authentic sample,² 58.9-59.8°.

2,4,6- (or 2,4,7-)Cyclooctatrienylideneacetonitrile (VIII or IX). - A solution of 0.68 g. of the nitrile III in 10 ml. of freshly distilled triethylamine was heated under reflux in a nitrogen atmosphere for 70 hours. The solution was cooled, diluted with 20 ml. of ether, and extracted with four 25-ml. portions of 3% hydrochloric acid at 0°. The ether solution was dried over sodium sulfate, concentrated under reduced pressure, and the residue was fractionated through a semimicro column. The yield of the isomeric nitrile VIII or IX was 0.59 g. (87%), b.p. 79° (0.3 mm.), n_D^{25} 1.6235-1.6255. An analytical sample had b.p. 79° (0.3 mm.), n_D^{25} 1.6255, d_4^{25} 1.0226, $\lambda_{\text{max.}}$ 242.5 m μ (log ϵ 4.31), $\lambda_{\text{max.}}$ 314 m μ (log ϵ 4.00) in 95% ethanol.

Anal. Calcd. for C₁₀H₉N: C, 83.83; H, 6.34; N, 9.79.

Found: C, 83.95; H, 6.38; N, 9.83.

The isomeric nitrile VIII or IX with an infrared spectrum essentially identical with the spectrum of the sample described above also was obtained by shaking the nitrile III with 20% sodium hydroxide at room temperature for 55 hours.

2,4,6-(or 2,4,7-)Cyclooctatrienylideneacetamide (X or XI). -

The nitrile VIII or IX was treated with hydrogen peroxide in alkaline aqueous acetone under the conditions described for conversion of III to IV. The extract of the amide was concentrated at a low temperature in a nitrogen atmosphere to avoid decomposition and the product was purified by sublimation

at 0.1 mm. with a heating block temperature of 120-130°. The rather unstable amide X or XI was isolated as yellow crystals, m.p. 96-98° (dec.), in a yield of 0.11 g. (32%). An analytical sample prepared by resublimation and crystallization from a mixture of benzene and 30-60° petroleum ether had m.p. 99-100° (dec.); $\lambda_{\text{max.}_1}$ 243 m μ (log ϵ 4.32), $\lambda_{\text{max.}_2}$ 314 m μ (log ϵ 3.92) in 95% ethanol.

Anal. Calcd. for $\text{C}_{10}\text{H}_{11}\text{NO}$: C, 74.50; H, 6.88; N, 8.69.

Found: C, 74.45; H, 7.05; N, 8.55.

Cambridge, Massachusetts




Fig. 1. - Infrared absorption spectra: curve 1, cyclooctatetraenylnitrile (III); curve 2, 2,4,6-(or 2,4,7-)cyclooctatrienyldeneacetonitrile (VIII or IX), both as pure liquids in a 0.025-cm. cell.

DISTRIBUTION LIST

<u>No. of Copies</u>	<u>Addressee</u>
2	Commanding Officer Office of Naval Research Branch Office 150 Causeway Street Boston, Massachusetts
1	Commanding Officer Office of Naval Research Branch Office 844 North Rush Street Chicago 11, Illinois
1	Commanding Officer Office of Naval Research Branch Office 346 Broadway New York 13, New York
1	Commanding Officer Office of Naval Research Branch Office 1000 Geary Street San Francisco 9, California
1	Commanding Officer Office of Naval Research Branch Office 1030 N. Green Street Pasadena 1, California
2	Officer-in-Charge Office of Naval Research Branch Office Navy Number 100 Fleet Post Office New York, New York
6	Director, Naval Research Laboratory Washington 25, D. C. Attention: Technical Information Officer
4	Chief of Naval Research Office of Naval Research Washington 25, D. C. Attn: Chemistry Branch
1	Dr. Ralph G. H. Siu, Research Director General Laboratories QM Depot 2800 S. 20th Street Philadelphia 45, Pennsylvania
1	Dr. Warren Stubblebine, Research Director Chemical and Plastics Section, RDB-MPD Quartermaster General's Office Washington 25, D. C.
1	Dr. A. Stuart Hunter, Tech. Director Research and Development Branch MFD Quartermaster General's Office Washington 25, D. C.

<u>No. of Copies</u>	<u>Addressee</u>
1	Dr. A. Weissler Department of the Army Office of the Chief of Ordnance Washington 25, D. C. Attention: ORDTB-PS
1	Research and Development Group Logistics Division, General Staff Department of the Army Washington 25, D. C. Attn: Dr. W. T. Read, Scientific Adviser
2	Director, Naval Research Laboratory Washington 25, D. C. Attention: Chemistry Division
2	Chief of the Bureau of Ships Navy Department Washington 25, D. C. Attention: Code 340
2	Chief of the Bureau of Aeronautics Navy Department Washington 25, D. C. Attention: Code TD-4
2	Chief of the Bureau of Ordnance Navy Department Washington 25, D. C. Attention: Code Rexd
1	Dr. H. A. Zahl, Tech. Director Signal Corps Engineering Laboratories Fort Monmouth, New Jersey
1	U.S. Naval Radiological Defense Lab. San Francisco 24, California Attention: Technical Library
1	Naval Ordnance Test Station (Inyokern) China Lake, California Attention: Head, Chemistry Division
1	Office of Ordnance Research 2127 Myrtle Drive Durham, North Carolina
1	Technical Command Chemical Corps Chemical Center, Maryland
1	U.S. Atomic Energy Commission Research Division Washington 25, D. C.

<u>No. of Copies</u>	<u>Addressee</u>
1	U.S. Atomic Energy Commission Chemistry Division Brookhaven National Laboratory Upton, New York
1	U.S. Atomic Energy Commission Library Branch, Tech. Information, ORE P.O. Box E Oak Ridge, Tennessee
1	National Bureau of Standards Washington 25, D. C. Attn: Dr. L. A. Wall
5	ASTIA Document Service Center Knott Building Dayton 2, Ohio
1	Office of Technical Services Department of Commerce Washington 25, D. C.
1	Office of Secretary of Defense Pentagon, Room 3D1041 Attn: Library Branch Washington 25, D. C. (R and D)
1	Dr. A. G. Horney Office of Scientific Research R and D Command - USAF Box 1395 Baltimore, Maryland

Armed Services Technical Information Agency

Because of our limited supply, you are requested to return this copy WHEN IT HAS SERVED YOUR PURPOSE so that it may be made available to other requesters. Your cooperation will be appreciated.

AD

42475

NOTICE: WHEN GOVERNMENT OR OTHER DRAWINGS, SPECIFICATIONS OR OTHER DATA ARE USED FOR ANY PURPOSE OTHER THAN IN CONNECTION WITH A DEFINITELY RELATED GOVERNMENT PROCUREMENT OPERATION, THE U. S. GOVERNMENT THEREBY INCURS NO RESPONSIBILITY, NOR ANY OBLIGATION WHATSOEVER; AND THE FACT THAT THE GOVERNMENT MAY HAVE FORMULATED, FURNISHED, OR IN ANY WAY SUPPLIED THE SAID DRAWINGS, SPECIFICATIONS, OR OTHER DATA IS NOT TO BE REGARDED BY IMPLICATION OR OTHERWISE AS IN ANY MANNER LICENSING THE HOLDER OR ANY OTHER PERSON OR CORPORATION, OR CONVEYING ANY RIGHTS OR PERMISSION TO MANUFACTURE, USE OR SELL ANY PATENTED INVENTION THAT MAY IN ANY WAY BE RELATED THERETO.

Reproduced by
DOCUMENT SERVICE CENTER
KNOTT BUILDING, DAYTON, 2, OHIO

UNCLASSIFIED