

THIS REPORT HAS BEEN DELIMITED
AND CLEARED FOR PUBLIC RELEASE
UNDER DOD DIRECTIVE 5200.20 AND
NO RESTRICTIONS ARE IMPOSED UPON
ITS USE AND DISCLOSURE.

DISTRIBUTION STATEMENT A

APPROVED FOR PUBLIC RELEASE;
DISTRIBUTION UNLIMITED.

Armed Services Technical Information Agency

Because of our limited supply, you are requested to return this copy WHEN IT HAS SERVED YOUR PURPOSE so that it may be made available to other requesters. Your cooperation will be appreciated.

AD

43632

NOTICE: WHEN GOVERNMENT OR OTHER DRAWINGS, SPECIFICATIONS OR OTHER DATA ARE USED FOR ANY PURPOSE OTHER THAN IN CONNECTION WITH A DEFINITELY RELATED GOVERNMENT PROCUREMENT OPERATION, THE U. S. GOVERNMENT THEREBY INCURS NO RESPONSIBILITY, NOR ANY OBLIGATION WHATSOEVER; AND THE FACT THAT THE GOVERNMENT MAY HAVE FORMULATED, FURNISHED, OR IN ANY WAY SUPPLIED THE SAID DRAWINGS, SPECIFICATIONS, OR OTHER DATA IS NOT TO BE REGARDED BY IMPLICATION OR OTHERWISE AS IN ANY MANNER LICENSING THE HOLDER OR ANY OTHER PERSON OR CORPORATION, OR CONVEYING ANY RIGHTS OR PERMISSION TO MANUFACTURE, USE OR SELL ANY PATENTED INVENTION THAT MAY IN ANY WAY BE RELATED THERETO.

Reproduced by
DOCUMENT SERVICE CENTER
KNOTT BUILDING, DAYTON, 2, OHIO

UNCLASSIFIED

AD No. ~~43632~~

ASTIA FILE COPY

Final Report

on


Contract No. Nonr-495(OL)

NR 055 298

Shalom Sarel and Melvin S. Newman

5 August 1954

**THE OHIO STATE UNIVERSITY
RESEARCH FOUNDATION**



RF Project 497
Report No. Final

FINAL
R E P O R T

By

THE OHIO STATE UNIVERSITY
RESEARCH FOUNDATION

Columbus 10, Ohio

To: OFFICE OF NAVAL RESEARCH
Contract No. Nonr-495(04)
NR 055 298

On: STERIC FACTORS IN ORGANIC CHEMISTRY

For the period: January 1, 1953 - July 31, 1954

Submitted by: Shalom Sarel and Melvin S. Newman
Department of Chemistry

Date: August 5, 1954

STERIC FACTORS IN ORGANIC CHEMISTRY

The object of this research program was to learn more about steric factors in the hydrolysis of esters of acetic acid. By comparing the rates of hydrolysis of a variety of primary alkyl acetates with that of ethyl acetate the steric effect of variations in the alkyl moiety can be evaluated if one assumes that substitutions on the β -carbon of the alkyl group have a negligible polar effect. In a similar way the steric effects in secondary alkyl acetates may be evaluated by comparing the rates of hydrolysis of secondary alkyl acetates with that of isopropyl acetate.


In this final report the synthesis of the alkyl acetates is described and the physical properties etc., of all compounds used in this research are tabulated.

The discussion of the significance of this work as well as the synthetic work involved is to be presented in two or three papers which are now being written for submission to the Journal of the American Chemical Society. Reprints of these papers will be mailed for distribution.

EXPERIMENTAL

Acetates of primary, secondary, and tertiary alcohols were prepared in good purity, and the rate of alkaline hydrolysis in 70% aqueous dioxane, or in water, at 20° and 30°C. was measured.


For the preparation of the primary and secondary alcohols and their acetates, the general methods used can be depicted as follows:




A. SYNTHESIS OF ACIDS

Six different methods were used for the preparation of the carboxylic acids or their esters. These methods are outlined below:


(1) Carboxylation of the Grignard Reagent:




(2) Alkylation of Malonic or Cyanoacetic Esters:




(3) 1,4-Addition of Grignard Reagent to Ethyl Alkylidene Malonate:





(4) Alkylation of Nitriles (Ziegler Method):



(5) Hypochlorite Oxidation of Methyl Ketones:



(6) Alkaline Rearrangement of α -Haloketones (Favorski Reaction):




In method (6) the α -bromoketones were prepared either (a) by direct bromination of the corresponding ketones or (b) through replacement of hydroxyl group in an acyloin by bromine. The ketones themselves were prepared either by oxidation of an appropriate alcohol or through treatment of a cadmium dialkyl with the appropriate acyl halide.

B. FORMATION OF ALCOHOLS

Lithium Aluminum hydride was used as a reducing reagent for converting carboxylic acids and esters into the corresponding primary alcohols, and ketones to corresponding secondary alcohols.

C. FORMATION OF ACETATES

The new alcohols were smoothly and conveniently converted into their respective acetates by treatment with: (1) acetyl chloride or acetic anhydride and pyridine; or (2) isopropenyl acetate and acid; or (3) acetyl chloride and magnesium metal in dry ether as follows:



D. RATES OF ALKALINE HYDROLYSIS

The rates of hydrolysis of the acetates with 0.01 N sodium hydroxide, using water or 70% aqueous dioxane as solvents, was followed titrimetrically at 20°C. and 30°C.

E. INFRARED SPECTRA

Spectra of all acids, ketones, alcohols and acetates were recorded between 5000 and 625 cm^{-1} with a Baird Infrared Recording Spectrophotometer Model B. All liquids were measured in a sandwich-type sodium chloride cell.

TABLE I. ACETATES OF ALCOHOLS OF FORMULA $R'' - C - CH_2OC(=O)CH_3$


$\begin{matrix} R' \\ | \\ R'' - C - CH_2OC(=O)CH_3 \\ | \\ R''' \end{matrix}$

No.	Alcohol	R'	R''	R'''	Carboxylic acid or ester		Alcohol formation,		Acetylation of alcohols	
					Method of preparation	Yield in %	Yield in %	in %	Yield in %	Method of formation
1.	2-Methyl-1-butanol	CH ₃	C ₂ H ₅	H	---	--	--	--	77	(1)
2.	2,3-Dimethyl-1-butanol	CH ₃	CH ₃ -CH	H	A-2	60	90	92		(2)
3.	2,3,3-Trimethyl-1-butanol	CH ₃	CH ₃ -C(CH ₃) ₂	H	A-6	40	68	89		(1)
4.	2-Ethyl-1-butanol	C ₂ H ₅	C ₂ H ₅	H	---	--	--	70		(1)
5.	3-Methyl-2-ethyl-1-butanol	C ₂ H ₅	(CH ₃) ₂ CH	H	A-2	78	83	90		(1)
6.	3,3-Dimethyl-2-ethyl-1-butanol	C ₂ H ₅	(CH ₃) ₃ C	H	---	--	96	97		(1)
7.	2-Ethyl-1-hexanol	C ₂ H ₅	n-C ₄ H ₉	H	---	--	--	93		(1)
8.	2-Isopropyl-3-methyl-1-butanol	(CH ₃) ₂ CH	(CH ₃) ₂ CH	H	A-2 A-6	50 83	87	72		(1)
9.	3,3-Dimethyl-1-butanol	(CH ₃) ₃ CCH ₂	H	H	A-3		83	81		(2)
10.	Cyclopropanemethanol	CH ₂ -CH		H	A-5	95	70	89		(1)
11.	Cyclobutanemethanol	CH ₂ -CH ₂ -CH ₂ -CH ₂		H	---	--	80	87		(1)
12.	Cyclopentanemethanol	CH ₂ -CH ₂ -CH ₂ -CH ₂ -CH ₂	(CH ₂) ₄	H	A-6	65	80	90		(1)
13.	Cyclohexanemethanol	CH ₂ -CH ₂ -CH ₂ -CH ₂ -CH ₂ -CH ₂	(CH ₂) ₅	H	---	--	--	77		(1)
14.	Neopentyl alcohol	CH ₃	CH ₃	CH ₃	---	--	91	80		(1)

TABLE I. (Continued)

No.	Alcohol	R'	R''	R'''	Carboxylic acid or ester		Alcohol formation, yield in %	Acetylation of alcohols	
					Method of preparation	Yield in %		Yield in %	Method of formation (1)
15.	2,2-Dimethyl-1-butanol	C ₂ H ₅	CH ₃	CH ₃	A-1	40	90	92	(1)
16.	2-Methyl-2-ethyl-1-butanol	C ₂ H ₅	C ₂ H ₅	CH ₃	A-6	76	83	75	(1)
17.	2,2-Diethyl-1-butanol	C ₂ H ₅	C ₂ H ₅	C ₂ H ₅	A-4	60	65	92	(1)

TABLE II. ACETATES OF SECONDARY AND TERTIARY ALCOHOLS



No.	Compound	R'	R''	R'''	Alcohol formation, yield in %		Acetylation of alcohols, Yield in %		Method of formation
					formation, yield in %		Yield in %		
18.	Pinacolyl acetate	CH ₃	(CH ₃) ₃ C	H	87		63		(1)
19.	Diisopropylcarbinyl acetate	(CH ₃) ₂ CH	(CH ₃) ₂ CH	H	93		92		(2)
							82		(1)
							90		(3)
20.	Diisobutylcarbinyl acetate	(CH ₃) ₂ CHCH ₂	CH ₃ > CH-CH ₂ CH ₃	H	--		90		(1)
21.	Tertiary-butyl acetate	CH ₃	CH ₃	CH ₃	--		52		(1)
22.	Triethylcarbinyl acetate	C ₂ H ₅	C ₂ H ₅	C ₂ H ₅	--		70		(3)

TABLE III. PHYSICAL PROPERTIES OF CARBOXYLIC ACIDS OR THEIR ESTERS

Compound	Structure	Formula	B.P.		Index of refraction	Density
			Temp., °C.	Pressure, mm	n_D^{25}	d_{25}^{25}
2,3-Dimethylbutyric acid	$\text{CH}_3\text{CH}(\text{CH}_3)\text{CH}(\text{CH}_3)\text{COOH}$	$\text{C}_6\text{H}_{12}\text{O}_2$	188-189	740	1.4127	---
Methyl 2,3,3-Trimethyl Butyrate	$\text{CH}_3\text{CH}(\text{CH}_3)\text{C}(\text{CH}_3)_2\text{CH}_2\text{COOCH}_3$	$\text{C}_8\text{H}_{16}\text{O}_2$	61	30	n_D^{25} 1.4188 n_D^{22} 1.4194	d_{22} 0.8885
Diisopropylacetic acid	$(\text{CH}_3)_2\text{CHCH}_2\text{COOH}$	$\text{C}_6\text{H}_{12}\text{O}_2$	214	740	n_D^{25} 1.4260	d_{25}^{25} 0.9125
Methyl diisopropylacetate	$(\text{CH}_3)_2\text{CHCH}_2\text{COOCH}_3$	$\text{C}_9\text{H}_{18}\text{O}_2$	72.5	25	n_D^{20} 1.4159	---
3,3-Dimethylbutyric acid	$(\text{CH}_3)_3\text{CCH}_2\text{COOH}$	$\text{C}_6\text{H}_{12}\text{O}_2$	5.6 181-181.5	740	n_D^{25} 1.4084	---
Cyclopropanecarboxylic acid		$\text{C}_4\text{H}_6\text{O}_2$	181-182	745	n_D 1.4363	d_{24}^{24} 1.0847

TABLE III. (Continued)

Compound	Structure	Formula	B.P.		Index of refraction	Density
			Temp., °C.	Pressure, mm		
Methyl cyclopentanecarboxylate		$C_7H_{12}O_2$	154	740	n_D 1.4353	
2,2-Dimethylbutyric acid		$C_6H_{12}O_2$	186	745		
Methyl 2,2-Diethylpropionate		$C_8H_{16}O_2$	62-63	25	n_D^{26} 1.4085	d_{25} 0.8824
Triethylacetic acid		$C_8H_{16}O_2$	131-132	20		
2-Ethyl isovaleric acid		$C_7H_{14}O_2$	196-200	740	n_D 1.4148	

TABLE IV. PHYSICAL PROPERTIES OF BRANCHED PRIMARY AND SECONDARY ALCOHOLS

No.	Compound	Formula	B.P.		n_D^{25}	d_{25}	$[\alpha_D]_D$	
			Temp., °C.	Pressure, mm			Calcd.	Found
1.	$\begin{array}{c} \text{C}_2\text{H}_5-\text{CH}-\text{CH}_2\text{OH} \\ \\ \text{CH}_3 \end{array}$	$\text{C}_5\text{H}_{12}\text{O}$	128	740	1.4104			
2.	$\begin{array}{c} \text{CH}_3\text{CH}-\text{CH}-\text{CH}_2\text{OH} \\ \quad \\ \text{CH}_3 \quad \text{CH}_3 \end{array}$	$\text{C}_6\text{H}_{14}\text{O}$	145-146	741	1.4173			
3.	$\begin{array}{c} \text{CH}_3 \quad \text{CH}_3 \\ \quad \\ \text{CH}-\text{C}-\text{CH}-\text{CH}_2\text{OH} \\ \\ \text{CH}_3 \end{array}$	$\text{C}_7\text{H}_{16}\text{O}$	158-159	740	1.4230	0.8238	36.05	35.86
4.	$\begin{array}{c} \text{C}_2\text{H}_5-\text{CHCH}_2\text{OH} \\ \\ \text{C}_2\text{H}_5 \end{array}$	$\text{C}_6\text{H}_{14}\text{O}$	146-147	741	1.4205			
5.	$\begin{array}{c} \text{CH}_3-\text{CH}-\text{CH}-\text{CH}_2\text{OH} \\ \quad \\ \text{CH}_3 \quad \text{C}_2\text{H}_5 \end{array}$	$\text{C}_7\text{H}_{16}\text{O}$	84-86	38	1.4234	0.8327	36.05	35.61
6.	$\begin{array}{c} \text{CH}_3 \\ \\ \text{CH}-\text{C}-\text{CH}-\text{CH}_2\text{OH} \\ \quad \\ \text{CH}_3 \quad \text{C}_2\text{H}_5 \end{array}$	$\text{C}_8\text{H}_{18}\text{O}$	88-89	38	1.4348	0.8425	40.66	40.32
7.	$\begin{array}{c} n\text{-C}_4\text{H}_9-\text{CH}-\text{CH}_2\text{OH} \\ \\ \text{C}_2\text{H}_5 \end{array}$	$\text{C}_8\text{H}_{18}\text{O}$	184	741	1.4280			

TABLE IV. (Continued)

No.	Compound	Formula	B.P.		n_D^{25}	d_{25}^{25}	$[\alpha]_D^{25}$	
			Temp., °C.	Pressure, mm			Calcd.	Found
8.		$C_8H_{18}O$	171-172	741	1.4342	0.8425	40.66	40.26
9.		$C_6H_{14}O$	144-145	740	1.4115	0.8097	31.43	31.35
10.		C_4H_8O	123	740	1.4297	0.9098		20.47
11.		$C_5H_{10}O$	140	740	1.4430			
12.		$C_6H_{12}O$	160	740	1.4550			
13.		$C_7H_{14}O$	180	741	1.4634			

TABLE IV. (Continued)

No.	Compound	Formula	B.P.		25 n _D	d 25	[R _L] _D	
			Temp., °C.	Pressure, mm.			Calcd.	Found
14.	$\begin{array}{c} \text{CH}_3 \\ \\ \text{CH}_3 - \text{C} - \text{CH}_2\text{OH} \\ \\ \text{CH}_3 \end{array}$	$\text{C}_5\text{H}_{12}\text{O}$	112-114	740	m.p. 51-51.6°			
15.	$\begin{array}{c} \text{CH}_3 \\ \\ \text{C}_2\text{H}_5 - \text{C} - \text{CH}_2\text{OH} \\ \\ \text{CH}_3 \end{array}$	$\text{C}_6\text{H}_{14}\text{O}$	134	740		1.4192		
16.	$\begin{array}{c} \text{CH}_3 \\ \\ \text{C}_2\text{H}_5 - \text{C} - \text{CH}_2\text{OH} \\ \\ \text{C}_2\text{H}_5 \end{array}$	$\text{C}_7\text{H}_{16}\text{O}$	152	740		1.4288		
17.	$\begin{array}{c} \text{CH}_3 \\ \\ (\text{C}_2\text{H}_5)_3\text{C} - \text{CH}_2\text{OH} \\ \\ \text{CH}_3 \end{array}$	$\text{C}_8\text{H}_{18}\text{O}$	92	25		1.4411		
18.	$\begin{array}{c} \text{CH}_3 \\ \\ \text{CH}_3 - \text{C} - \text{CHOHCH}_3 \\ \\ \text{CH}_3 \end{array}$	$\text{C}_6\text{H}_{14}\text{O}$	120-120.5	741		1.4153	0.8122	31.43 31.50
19.	$\begin{array}{c} \text{CH}_3 \quad \text{CH}_3 \\ \diagdown \quad \diagup \\ \text{CH} - \text{CHOHCH} \\ \diagup \quad \diagdown \\ \text{CH}_3 \quad \text{CH}_3 \end{array}$	$\text{C}_7\text{H}_{16}\text{O}$	139-139.5	740		1.4210	0.8245	36.05 35.71

TABLE IV. (Continued)

No.	Compound	Formula	B.P.		n_D^{25}	d_{25}	$[\alpha]_D$	
			Temp., °C.	Pressure, mm			Calcd.	Found
20.	$\begin{array}{c} \text{CH}_3 \\ \\ \text{CHCH}_2\text{CHOHCH}_2\text{CH} \\ \qquad \qquad \\ \text{CH}_3 \qquad \qquad \text{CH}_3 \end{array}$	$\text{C}_9\text{H}_{20}\text{O}$						
21.	$(\text{C}_2\text{H}_5)_3\text{COH}$	$\text{C}_7\text{H}_{16}\text{O}$	140-141	740	1.4256			

TABLE V. PHYSICAL PROPERTIES OF ACETATES OF HIGHLY-BRANCHED ALCOHOLS

No.	Acetate of:	Formula	B.P.		Pressure, mm	n_D^{25}	d_{25}	$[\alpha]_D^{25}$	
			°C	°C				Calcd.	Found
1.	2-Methyl-1-butanol	$C_7H_{14}O_2$	138-139		741	1.3996	0.8719	36.17	36.17
2.	2,3-Dimethyl-1-butanol	$C_8H_{16}O_2$	147		740	1.4068	0.8790	40.79	40.50
3.	2,3,3-Trimethyl-1-butanol	$C_9H_{18}O_2$	170-171		741	1.4125	0.8687	45.41	45.30
4.	2-Ethyl-1-butanol	$C_8H_{16}O_2$	160		740	1.4090	0.8764	40.79	40.68
5.	3-Methyl-2-ethyl-1-butanol	$C_9H_{18}O_2$	88-89		30	1.4156	0.8774	45.41	45.23
6.	3,3-Dimethyl-2-ethyl-1-butanol	$C_{10}H_{20}O_2$	100		40	1.4220	0.8770	50.0	49.90
7.	2-Ethyl-1-hexanol	$C_{10}H_{20}O_2$	104		40	1.4182	0.8688	50.0	49.97
8.	3,3-Dimethyl-1-butanol	$C_8H_{16}O_2$	156-157		741	1.4038	0.8683	40.79	40.60
9.	2-Isopropyl-3-methyl-1-butanol	$C_{10}H_{20}O_2$	130		110	1.4200	0.8803	50.0	49.5
10.	Cyclopropanemethanol	$C_6H_{10}O_2$	133.5		741	1.4156	0.9603		
11.	Cyclobutanemethanol	$C_7H_{12}O_2$	150		741	1.4245	0.9508		
12.	Cyclopentanemethanol	$C_8H_{14}O_2$	172.5		741	1.4340	0.9577		
13.	Cyclohexanemethanol	$C_9H_{16}O_2$	108		40	1.4422	0.9541		
14.	Neopentyl alcohol	$C_7H_{14}O_2$	127		740	1.3927	0.8539	36.17	36.36
15.	2,2-Dimethyl-1-butanol	$C_8H_{16}O_2$	152-153		740	1.4050	0.8704	40.79	40.60
16.	2-Methyl-2-ethyl-1-butanol	$C_9H_{18}O_2$	100		80	1.4150	0.8815	45.41	44.95

TABLE V. (Continued)

No.	Acetate of:	Formula	B.P.		n_D^{25}	d_{25}	$[\alpha]_D^{25}$	
			°C	Pressure, mm			Calcd.	Found
17.	2,2-Diethyl-1-butanol	$C_{10}H_{20}O_2$	103-104	40	1.4269	0.8900	50.00	49.68
18.	Pinacolyc alcohol	$C_8H_{16}O_2$	138-138.5	738	1.4002			
19.	Diisopropylcarbinol	$C_9H_{18}O_2$	161	745	1.4110*			
20.	Diisobutylcarbinol	$C_{11}H_{22}O_2$	125-126	60	1.4117			
21.	Trimethylcarbinol	$C_6H_{12}O_2$	96.5-97.0	745	1.3840			
22.	Triethylcarbinol	$C_9H_{18}O_2$	163	740	1.4270			

* n_D^{20}

TABLE VI. RATE OF SAPONIFICATION OF ACETATES OF PRIMARY ALCOHOLS

No.	Compound	Solvent	Normality of NaOH	Molarity of ester	Rate constant, k_2 ,		$\frac{k_{\text{ethyl acetate}}}{k_{\text{alkyl acetate}}}$
					$\frac{\text{g. mol. l.}}{20^\circ\text{C.}}$	$\frac{\text{l.}}{30^\circ\text{C.}}$	
1.	2-Methyl-1-butyl acetate	Water	0.012	0.0099	1.420	---	---
		70% Dioxane	0.015	0.0075	0.47	---	3.0
			0.0119	0.009	---	0.78	---
2.	2,3-Dimethyl-1-butyl acetate	70% Dioxane		0.009	0.35	---	---
			0.0133	0.008	---	0.72	3.3
			0.0119	0.009	---	---	---
3.	2,3,3-Trimethyl-1-butyl acetate	70% Dioxane	0.0123	0.007	0.324	---	3.7
				0.009	---	---	---
4.	2-Ethyl-1-butyl acetate	70% Dioxane	0.0133	0.0109	0.24	---	---
				0.0090	---	---	4.8
			0.0119	0.0093	---	0.49	---
				0.0100	---	---	---
5.	3-Methyl-2-ethyl-1-butyl acetate	70% Dioxane	0.015	0.010	0.23	---	---
				0.008	---	---	5.8
			0.0119	0.009	---	0.41	---
				0.010	---	---	---
6.	3,3-Dimethyl-2-ethyl-1-butyl acetate	70% Dioxane	0.015	0.0107	0.064	---	---
				0.0093	---	---	21.0
			0.0119	0.0095	---	0.11	---
				0.0100	---	---	---
7.	2-Ethyl-1-hexyl acetate	70% Dioxane	0.0125	0.0098	0.156	---	---
			0.0119	0.0102	---	0.36	6.5
				0.0085	---	---	---
8.	3,3-Dimethyl-1-butyl acetate	70% Dioxane	0.0123	0.0105	0.627	---	1.9
				0.0087	---	---	---

TABLE VI. (Continued)

No.	Compound	Solvent	Normality of NaOH	Molarity of ester	Rate constant, k_2 , $\frac{\text{g. mol. l.}}{\text{min. l.}} \cdot \frac{1}{30^\circ\text{C.}}$	$k_{\text{ethyl acetate}}$ $k_{\text{allyl acetate}}$
9.	2-Isopropyl-3-methyl-1-butyl acetate	70% Dioxane	0.0122	0.0087 0.0078	0.092	13.3
10.	Cyclopropanemethyl acetate	70% Dioxane	0.0133	0.0084 0.0100 0.0086 0.0084	1.03 ----- 2.28	1.004
11.	Cyclobutanemethyl acetate	70% Dioxane	0.0133	0.0084 0.0096 0.0090 0.0080	0.74 ----- 1.47	1.6
12.	Cyclopentanemethyl acetate	70% Dioxane	0.0133	0.0093 0.0087 0.0080 0.0096 0.0081	0.54 ----- 1.08	2.2
13.	Cyclohexanemethyl acetate	70% Dioxane	0.0135	0.0092 0.0100 0.0101 0.0098	0.33 ----- 0.70	3.4
14.	Neopentyl acetate	Water 70% Dioxane	0.0122 0.015 0.0119	0.0074 0.0069 0.0078 0.0083 0.0100	0.86 0.28 ----- 0.50	4.8

TABLE VI. (Continued)

No.	Compound	Solvent	Normality of NaOH	Molarity of ester	Rate constant, k_2 , g. mol. l./min. ⁻¹		$k_{\text{ethyl acetate}}$ $k_{\text{alkyl acetate}}$
					20°C.	30°C.	
15.	2,2-Dimethyl-1-butyl acetate	Water	0.012	0.007	0.70	----	
		70% Dioxane	0.015	0.0101	0.21	----	7.0
				0.0088			
			0.0119	0.0090	----	0.34	
				0.0082			
16.	2-Methyl-2-ethyl-1-butyl acetate	70% Dioxane	0.0123	0.0087	0.124	----	12.0
				0.0090			
17.	2,2-Diethyl-1-butyl acetate	70% Dioxane	0.0133	0.0092	0.050	----	
				0.0069			
			0.0119	0.0093	-----	0.108	22.5
				0.0080			

TABLE VII. RATE OF SAPONIFICATION OF ACETATES OF SECONDARY ALCOHOLS

No.	Compound	Solvent	Normality of NaOH	Molarity of ester	Rate constant, k_2 , $\frac{\text{g. mol. l.}}{\text{min. l.}}$		k	$\frac{k_{\text{isopropyl acetate}}}{k_{\text{alkyl acetate}}}$
					20°C.	30°C.		
1.	Pinacolyl acetate	70% Dioxane	0.0245	0.0100	0.035	-----	-----	11.3
			0.012	0.0108	-----	0.054	-----	
2.	Diisobutylcarbinyl acetate	70% Dioxane	0.025	0.0079	-----	0.0236	-----	25.0
				0.0091	-----		-----	
3.	Tert. butyl acetate	70% Dioxane	0.025	0.0075	-----	0.0381	-----	-----
				0.0090	-----		-----	
4.	Diisopropylcarbinyl acetate(*)	70% Dioxane	0.025	0.0079	-----	0.0109	-----	56.0

(*) Rate constant calculated graphically by extrapolating the line obtained from 30% to 80% hydrolysis.

Signature Page

to

FINAL Report
RF Project 497

on

Contract No. Nonr-495(04)
NR 055 298

Investigator S. Sarel Date 8-10-54
per in SN

Supervisor Melvin S. Newman Date 8-10-54

For The Ohio State University Research Foundation

Executive Director Oran C. Woodport Date 8/10/54
WOF

5 August 1954
RF 497

Distribution List for Final Report
Contract No. Nonr-495(04) NR 055 298
STERIC FACTORS IN ORGANIC CHEMISTRY

Copies

- 1 Director, ONR Branch Office
150 Causeway Street
Boston, Mass.
- 2 Director, ONR Branch Office
The John Crerar Library Bldg.
10th Floor, 86 E. Randolph St.
Chicago 1, Illinois
- 1 Director, ONR Branch Office
1000 Geary Street
San Francisco 9, Calif.
- 1 Director, ONR Branch Office
1030 N. Green Street
Pasadena 1, Calif.
- 2 Officer in Charge
ONR, Navy No. 100
Fleet Post Office
New York, New York
- 6 Director, Naval Research Lab.
Washington 25, D. C.
Attn: Technical Information
Officer
- 2 Chief of Naval Research
Washington 25, D. C.
Attn: Chemistry Branch
- 1 Dr. Ralph G. H. Sie, Research
Director
General Laboratories, QM Depot
2600 S. 20th Street
Philadelphia 45, Pennsylvania
- 1 Dr. Warren Stubblebine
Research Director
Chemical and Plastics Section RDB-MPD
Quartermaster General's Office
Washington 25, D. C.

Copies

- 1 Dr. A. Stuart Hunter,
Technical Director
Research and Development
Branch MPD
Quartermaster General's
Office
Washington 25, D. C.
- 1 Dr. A. Weissler
Department of The Army
Office of the Chief of
Ordnance
Washington 25, D. C.
Attn: ORDTB-PS
- 1 Research and Development
Group
Logistics Division, General
Staff
Department of The Army
Washington 25, D. C.
Attn: Dr. W. T. Read
Scientific Adviser
- 2 Director, Naval Research Lab.
Washington 25, D. C.
Attn: Chemistry Division
- 2 Chief of the Bureau of Ships
Navy Department
Washington 25, D. C.
Attn: Code 340
- 2 Chief of Bureau of Aeronautics
Navy Department
Washington 25, D. C.
Attn: Code TD-4
- 2 Chief of the Bureau of
Ordnance
Navy Department
Washington 25, D. C.
Attn: Code Rexd

5 August 1954
RF 497

Distribution List (Continued)

Copies

- 1 Dr. H. A. Zahl, Tech. Director
Signal Corps Engineering Labs.
Fort Monmouth, New Jersey
- 1 U. S. Naval Radiological Defense Lab.
San Francisco 24, Calif.
Attn: Technical Library
- 1 Naval Ordnance Test Station
Inyokern
CHINA LAKE, California
Attn: Head, Chemistry Div.
- 1 Office of Ordnance Research
2127 Myrtle Drive
Durham, North Carolina
- 1 Technical Command
Chemical Corps
Chemical Center, Maryland
- 1 U. S. Atomic Energy Commission
Research Division
Washington 25, D. C.
- 1 U. S. Atomic Energy Commission
Chemistry Division
Brookhaven National Laboratory
Upton, New York
- 1 U. S. Atomic Energy Commission
Library Branch, Tech. Info., ORE
P. O. Box E
Oak Ridge, Tennessee
- 1 University of California
Department of Chemistry
Los Angeles 24, California
Attn: Dr. S. Winstein
- 1 Columbia University
Department of Chemistry
New York 27, New York
Attn: Dr. L. P. Hammett
- 1 University of Colorado
Department of Chemistry
Boulder, Colorado
Attn: Dr. S. J. Cristol

Copies

- 1 Harvard University
Department of Chemistry
Cambridge 38, Massachusetts
Attn: Dr. P. D. Bartlett
- 1 Purdue University
Department of Chemistry
Lafayette, Indiana
Attn: Dr. Herbert C. Brown
- 1 Massachusetts Institute of
Technology
Department of Chemistry
Cambridge, Massachusetts
Attn: Dr. C. G. Swain
- 1 Pennsylvania State College
Department of Chemistry
State College, Pennsylvania
Attn: Dr. R. W. Taft, Jr.
- 1 University of Southern Calif.
Department of Chemistry
Los Angeles 7, California
Attn: Dr. R. F. Brown
- 5 ASTIA Document Service Center
Knott Building
Dayton 2, Ohio
- 1 Office of Technical Services
Department of Commerce
Washington 25, D. C.
- 1 Office of Secretary of Defense
Pentagon, Room 3D1041
Washington 25, D. C.
Attn: Library Branch
(R and D)
- 1 Dr. A. G. Horney
Office Scientific Research
R and D Command, USAF
Box 1395
Baltimore, Maryland
- 1 Director, ONR Branch Office
346 Broadway
New York 13, New York

Armed Services Technical Information Agency

Because of our limited supply, you are requested to return this copy WHEN IT HAS SERVED YOUR PURPOSE so that it may be made available to other requesters. Your cooperation will be appreciated.

AD

43632

NOTICE: WHEN GOVERNMENT OR OTHER DRAWINGS, SPECIFICATIONS OR OTHER DATA ARE USED FOR ANY PURPOSE OTHER THAN IN CONNECTION WITH A DEFINITELY RELATED GOVERNMENT PROCUREMENT OPERATION, THE U. S. GOVERNMENT THEREBY INCURS NO RESPONSIBILITY, NOR ANY OBLIGATION WHATSOEVER; AND THE FACT THAT THE GOVERNMENT MAY HAVE FORMULATED, FURNISHED, OR IN ANY WAY SUPPLIED THE SAID DRAWINGS, SPECIFICATIONS, OR OTHER DATA IS NOT TO BE REGARDED BY IMPLICATION OR OTHERWISE AS IN ANY MANNER LICENSING THE HOLDER OR ANY OTHER PERSON OR CORPORATION, OR CONVEYING ANY RIGHTS OR PERMISSION TO MANUFACTURE, USE OR SELL ANY PATENTED INVENTION THAT MAY IN ANY WAY BE RELATED THERETO.

Reproduced by
DOCUMENT SERVICE CENTER
KNOTT BUILDING, DAYTON, 2, OHIO

UNCLASSIFIED