

UNCLASSIFIED

AD NUMBER
AD481024
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; Feb 1966. Other requests shall be referred to U.S. Army Biological Center, Fort Detrick, MD.
AUTHORITY
BORL D/A ltr, 28 Sep 1971

THIS PAGE IS UNCLASSIFIED

AD

TECHNICAL MANUSCRIPT 285


481024

TREPTOMYCIN INHIBITION OF ELABORATION
OF STAPHYLOCOCCAL ENTEROTOXIC PROTEIN

Albert J. Rosenwald
Ralph E. Lincoln

FEBRUARY 1966

UNITED STATES ARMY
BIOLOGICAL CENTER
FORT DETRICK



Reproduction of this publication in whole or part is prohibited except with permission of Commanding Officer, U. S. Army Biological Center, ATTN: Technical Releases Section, Technical Information Department, Fort Detrick, Frederick, Maryland, 21701. However, DDC is authorized to reproduce the publication for United States Government purposes.

DDC AVAILABILITY NOTICES

Qualified requestors may obtain copies of this publication from DDC.

Foreign announcement and dissemination of this publication by DDC is not authorized.

Release or announcement to the public is not authorized.

DISPOSITION INSTRUCTIONS

Destroy this publication when it is no longer needed. Do not return it to the originator.

The findings in this publication are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.

U.S. ARMY BIOLOGICAL CENTER
Fort Detrick, Frederick, Maryland

TECHNICAL MANUSCRIPT 285

STREPTOMYCIN INHIBITION OF ELABORATION
OF STAPHYLOCOCCAL ENTEROTOXIC PROTEIN

Albert J. Rosenwald

Ralph E. Lincoln

Process Development Branch
AGENT DEVELOPMENT AND ENGINEERING LABORATORY

Project 1C522301A059

February 1966

ABSTRACT

Staphylococcus aureus, strain S-6, under certain cultural conditions, synthesizes and elaborates into the medium an enterotoxigenic protein. We have observed that streptomycin interferes with the elaboration of this protein by streptomycin-resistant staphylococci at drug levels permitting continued synthesis of those proteins necessary for growth and multiplication of the organisms. It is likely that the enterotoxigenic protein is incomplete or that it is not released from the cell because of streptomycin-induced alterations in the cell itself.

STREPTOMYCIN INHIBITION OF ELABORATION OF STAPHYLOCOCCAL ENTEROTOXIC PROTEIN

Staphylococcus aureus, strain S-6, under certain cultural conditions, synthesizes and elaborates into the medium an enterotoxigenic protein.* We have observed that streptomycin interferes with the elaboration of this protein by streptomycin-resistant staphylococci at drug levels permitting continued synthesis of those proteins necessary for growth and multiplication of the organisms. Enterotoxin production in the streptomycin-dependent strain, even at the point of maximum population increase, was severely depressed.

Streptomycin sulfate (E.R. Squibb & Sons, New York) was added at various levels to a medium composed of N-Z-Amine, Type A (Sheffield Chem., Norwich, New York), 4.0%; yeast extract (Difco Laboratories, Detroit, Michigan), 0.4%; K_2HPO_4 , 0.1%; and tap water, at pH 6.8 to 7.0. Twenty-five ml of this medium in a 250 ml Erlenmeyer flask was inoculated (1.0% vol/vol) with a culture of a phage-free isolate of Staphylococcus aureus, strain S-6. Isolates sensitive, resistant, and dependent, with regard to streptomycin, were used. The cultures were shaken (100 three-inch strokes/min) at 37 C for 24 hours after which optical density (1:10 dilution) and enterotoxin level were determined, the latter by the Oudin method.**

Figure 1 shows that enterotoxin level declined with optical density when the streptomycin-sensitive strain was treated with increasing concentration of streptomycin. Employing the resistant strain of the organism (Figure 2), enterotoxin level was decreased over a broad range of drug concentration in which there was no inhibition of culture growth. Although streptomycin at 250 to 1000 $\mu\text{g/ml}$ was sufficient to allow normal culture growth of the streptomycin-dependent strain (Figure 3), the maximum detectable enterotoxigenic protein was less than 10% of that obtained with the sensitive or resistant strains and occurred at the lowest level of drug permitting normal culture growth.

These data show that streptomycin limits the elaboration of immunologically detectable enterotoxin in both resistant and dependent organisms under conditions in which they apparently are functioning normally in

* Bergdoll, M.S.; Surgalla, M.J.; Dack, G.M. 1959. Staphylococcal enterotoxin. Identification of a specific precipitating antibody with enterotoxin-neutralizing property. J. Immunol. 83:334-338.

** Oudin, J. 1952. III. Techniques and analysis of the quantitative precipitin reaction. B. Specific precipitation in gels and its application to immunochemical analysis, p. 335-378, In A.C. Corcoran (ed.), Methods in Medical Research, Vol. 5. The Year Book Publishers, Inc., Chicago.

other respects. The reasons for this are not clear. Enterotoxin is not detectable until it reaches the cell surface.* Hypotheses based on the similarity of sites of protein synthesis and those of streptomycin action are tempting. The cell membrane would be of particular interest. It is unlikely under current concepts, that the protein suffers from miscoding. It is more likely that it is incomplete or that it is not released from the cell because of streptomycin-induced alterations in the cell itself.

* Friedman, M.E.; White, J.D. 1965. Immunofluorescent demonstration of cell-associated staphylococcal enterotoxin B. J. Bacteriol. 89:1155.




Figure 1. Growth and Synthesis of Enterotoxin by Streptomycin-Sensitive *S. aureus* in Presence of Streptomycin.




Figure 2. Growth and Synthesis of Enterotoxin by Streptomycin-Resistant *S. aureus* in Presence of Streptomycin.




Figure 3. Growth and Synthesis of Enterotoxin by Streptomycin-Dependent *S. aureus* in Presence of Streptomycin.