

AD658348

317000

DECOMPRESSION SICKNESS IN HIGH-ALTITUDE FLIGHT

WILLARD L. MEADER, Major, USAF, MC

D D C
RECEIVED
SEP 25 1967
A

March 1967

**USAF School of Aerospace Medicine
Aerospace Medical Division (AFSC)
Brooks Air Force Base, Texas**

Reproduced by the
CLEARINGHOUSE
for Federal Scientific & Technical
Information Springfield Va 22151

Decompression Sickness in High-altitude Flight

MAJ. WILLARD L. MEADER, USAF, MC

Review of 5 years of a WU-2 squadron's daily records shows 36 cases of decompression sickness entirely confined to reports of bends pain among 11 crewmembers in 958 flights. Generally, 40 minutes were spent in denitrogenation in a partial pressure suit and helmet before achieving cabin altitudes of close to 29,000 feet. The majority of bends occurred within the first 3.5 hours affecting the knee joints in almost two-thirds of the cases with a tendency to recur at the same joint. Only 1 case required descent to lower altitude to afford relief from pain.

THE ETIOLOGY of decompression sickness continues to be of fascination to aerospace medicine. Neither research nor case reports of altitude chamber or in-flight reactions have demonstrated clear and precise causes of predisposing factors, other than an exposure to altitude together with inadequate denitrogenation and the engaging in physical exertion. Many previous reports have emphasized the results of low-pressure chamber operations.^{1,6} Findings reported for a few in-flight reactions include itches, bends, chokes, visual changes, paralysis, unconsciousness^{2,3,7,9} and death.^{5,8} Few of the reports of decompression sickness encountered in an operational squadron in daily flying have

been published. With these thoughts in mind, the author received permission from the Surgeon, Strategic Air Command, to review the accumulated records of the Physiological Support Division (PSD) of the 803rd Medical Group at Davis-Monthan AFB, Arizona.

MATERIALS AND METHODS

PSD provided support to a high-altitude weather reconnaissance squadron, integrating equipment inspection and maintenance with aeromedical support.⁴ The aircraft flown was the WU-2, which achieves altitudes above 60,000 feet. The Consolidated Crew Mission Data Sheets covered the period of September 1959 through December 1964, and contained a preflight medical check, as well as a postflight debriefing. Data recorded prior to flight included the date, time of day, the crewmember's oral temperature, pulse, blood pressure, body weight, and whether he had eaten the available high-protein meal. Time spent in the prebreathing of oxygen was recorded, and debriefing included total flight time, maximum cabin altitude, duration of cabin altitude over 25,000 feet, and whether he had any symptoms or medical problems. It was also noted whether or not there were decompressions, and if so, the duration and to what cabin altitude. The preflight data was obtained by either a flight surgeon or medical technician, and the debriefing data by a personal equipment technician,

From the USAF School of Aerospace Medicine, Aerospace Medical Division (AFSC), Brooks Air Force Base, Texas.

Current address of Major Meader is Hq. SAC (SUP2), Offutt AFB, Nebraska 68113.

DECOMPRESSION SICKNESS IN HIGH-ALTITUDE FLIGHT—MEADER

and a flight surgeon if there was a medical problem. The information received from crewmembers was felt to be reliable. Many had flown with the squadron since its inception and had acquired considerable knowledge

TABLE 1. BENDS EXPERIENCE IN ONE SQUADRON DURING 5 YEARS

Subject (Total Flights) Incidence	Flight Number	Duration Prebreathing (Min.)	Season	Time of Day	Flight Duration (Hrs.)	Time of Onset of Symptoms (Hrs.)	Maximum Cabin Altitude (Ft.)	Duration Cabin Altitude >25,000 Ft.	Decompressions (No.)	Duration (Min.)	Location of Bends
A (81) 1.2%	1	40	W	AM	8.2	—	29	7.0	0	—	Right knee
B (20) 15.0%	2	50	Su	AM	7.0	4.5	29.5	5.5	0	—	Both knees
	3	40	Sp	AM	7.0	4	30	6.0	0	—	Both knees
	4	40	Sp	—	7.0	—	28.7	6.0	0	—	Left knee
C (122) 0.8%	5	40	Su	AM	7.2	4.5	28.5	6.0	0	—	—
D (44) 2.2%	6	40	Su	AM	6.8	4.5	29.5	4.5	0	—	Left shoulder
E (96) 1.0%	7	40	Su	AM	7.7	3.5	29	7.0	0	—	—
F (50) 2.0%	8	30	Su	AM	4.9	3.0	28.5	4.0	0	—	Left elbow and wrist
G (56) 1.8%	9	30	W	AM	6.7	—	>60	5.7	1	10	Right knee
	10	40	F	AM	7.4	—	29	6.5	0	—	Right knee
	11	40	Sp	PM	5.2	2.5	29	4.2	0	—	Both knees
	12	40	F	AM	8.3	—	>60	6.0	1	20	Right knee
	13	30	W	AM	11.4	2.0	33	10.2	0	—	Right knee
	14	40	F	AM	4.9	2.5	45	4.0	0	—	Right foot
	15	40	F	AM	6.1	—	30	5.0	0	—	Both knees
	16	40	F	AM	7.2	6.0	29	6.5	0	—	Both knees, Right ankle
	17	40	F	AM	7.0	5.0	35	6.0	0	—	Both knees
	18	40	W	AM	7.6	5.5	29	6.7	0	—	Right knee
	19	40	Sp	AM	8.2	3.0	29	7.5	0	—	Left knee and ankle
	20	40	W	AM	6.6	5.5	29	5.7	0	—	Right knee
	21	40	F	AM	7.1	5.5	29	6.2	0	—	Right foot
	22	40	F	AM	7.7	5.0	29	6.7	0	—	Left ankle
	23	40	W	AM	7.0	2.0	29	6.0	0	—	Ankle
	24	40	W	PM	5.7	3.0	28	4.7	0	—	Left foot
	25	50	W	AM	7.3	—	28	5.4	0	—	—
	26	60	W	AM	8.7	7.5	29	7.7	0	—	Right ankle
	27	60	W	AM	8.0	—	28	6.5	0	—	Knee and ankle
	28	60	W	AM	8.2	3.5	29	7.0	0	—	Both knees
	29	60	W	AM	8.3	—	28	7.7	0	—	Left wrist, both knees
	30	60	Sp	AM	8.0	3.0	28.5	7.0	0	—	Left arm
	31	60	Sp	AM	8.7	—	29	7.7	0	—	Left knee and ankle
	32	30	Su	AM	7.2	—	29	6.2	0	—	Right knee and ankle
I (140) 13.5%											
J (113) 0.8%	33	30	W	AM	7.2	3.5	28.5	3.5	0	—	Right knee
	34	40	Sp	AM	7.7	—	29	6.8	0	—	Right knee and ankle
K (97) 3.0%	35	30	W	AM	8.5	3.5	29	7.0	0	—	Right shoulder
	36	45	F	AM	6.0	3.0	28.7	5.0	0	—	—

of and appreciation for the physiology of high-altitude flight.

Crewmembers wore the MC-3A partial pressure suit with the MA-2 helmet, inflatable gloves, and unpressurized boots. Outer coveralls, a parachute, and under-arm life preservers were used routinely. The seatkit contained the F-2400 oxygen regulator and emergency oxygen, along with survival equipment. About one and one-half hours prior to flight, the pilot was seen in the flight surgeon's office for his medical check. He ate the high-protein meal, if desired, and then was dressed. A check of the suit and seatkit was done, and the pilot started denitrogenation, remaining on oxygen until completion of the flight. The cockpit of the WU-2 is no larger than that of most single engine jets; hence movement is considerably restricted.

Initially the records were screened for completeness. It was hoped that the data could be coded for computer analysis. Because of the paucity of information and the few cases of decompression sickness, it was decided to examine the flights of each crewmember who had one or more incidents of decompression sickness. For these flights, all at altitudes in excess of 45,000 feet, an attempt was made to correlate the incidents with the previously mentioned factors.

During the five years, there were 36 cases of decompression sickness in eleven crewmembers. However, 19 of these episodes involved one crewmember, who frequently took mild analgesics for joint aches and pains. He also wore spectacles while flying, and it is possible that he opened his face-plate to adjust his glasses on some of the longer flights. If this pilot is excluded, the total operational incidence of decompression sickness was 0.17 per cent, and for flights at altitudes greater than 45,000 feet, 0.20 per cent. All cases of decompression sickness were recorded as bends. None caused a mission to be aborted, and only one case required descent to a lower altitude to continue the mission.

In analyzing the flights of each individual, no correlations could be made between bends and body temperature, pulse, blood pressure or weight. Similarly, there was no relation to the preflight meal, duration of flight,




Fig. 1. Frequency of involvement by site (43 occurrences).

DECOMPRESSION SICKNESS IN HIGH-ALTITUDE FLIGHT—MEADER




Fig. 2. Time of onset of symptoms.

cabin altitude and duration above 25,000 feet. Only two cases of bends occurred in flights having decompressions (flights 9 and 12) and during only one of these did the pilot notice the onset of his symptoms at that time.

These eleven crewmembers flew 958 flights at altitudes greater than 45,000 feet. Of these, 116 or 12 per cent were in the last twelve hours of the day, while only 6 per cent of the bends occurred then (Table I). It is also of interest to note that more cases (38 per cent) occurred in the winter (21 December to 21 March) than in other seasons.

The relationship between bends and prebreathing of oxygen, excluding the one unduly susceptible individual, is listed in Table II.

As expected the incidence decreases with longer prebreathing.

The anatomical location of the bends in the eleven subjects is noted in Figure 1. Eighty-nine per cent of the cases occurred in the lower extremities and fourteen per cent in the upper extremities (both upper and lower extremity pain occurred in flight 29). Crewmembers who had multiple episodes of bends tended to have these recur in the same location.

This is demonstrated in Table III with three of the four subjects who had recurrences.

In 24 of the reported cases of bends, time of onset of symptoms was noted (Figure 2). Joint pain appeared as early as two hours, and as late as seven and one-half hours after take-off, with 50 per cent occurring in the first three and one-half hours.

DISCUSSION

The records of 958 WU-2 flights were examined and analyzed in close detail with respect to decompression sickness. Bends was the only manifestation of decompression sickness found, occurring 36 times in eleven different crewmembers. This is in contrast to previous reports^{3,7} in which symptoms ranging from itching or paresthesias to collapse were seen.

TABLE II. RELATIONSHIP OF TIME OF PREOXYGENATION TO INCIDENCE OF BENDS

Time of Preoxygenation	Number of Flights	Incidence
50 - 60 min.	114	.009
40 - 49 min.	556	.020
30 - 39 min.	139	.036

TABLE III. SITE OF RECURRENCE OF BENDS

Subject	Number of Episodes	Location	Number of Occurrences
Subject B	3	Right knee	2
		Left knee	3
Subject H	4	Right knee	4
		Left knee	1
Subject I	16	Left arm	1
		Left wrist	1
		Right knee	8
		Left knee	7
		Right ankle	3
		Left ankle	4
		Right foot	2
Subject K	2	Right shoulder	1
		Right knee	1
		Right ankle	1

Analysis of the data showed a significantly greater proportion of bends in the lower extremities as opposed to the upper extremities. Other investigators have found, in chamber flights, that the opposite was true^{1,6,9} and there is no ready explanation for this finding. There was a greater incidence in the morning hours, and during the winter months. When bends recurred, they tended to recur in the same joints. Though the usual time of preoxygenation only varied from 30 to 60 minutes, the additional time seemed to afford better protection against bends. However, when bends occurred, the majority appeared within three and one-half hours of take-off (Figure 2). Continued reporting of operational experiences is need to provide more complete information about inflight decompression sickness.

REFERENCES

- ADLER, H. F.: Dysbarism. *SAM Aeromed. Rev.*, 1-64, Feb. 1964.
- BERRY, C. A.: Severe Dysbarism in Air Force Operations and Training. *U. S. Armed Forces Med. J.*, 9:937, 1958.
- BERRY, C. A., and SMITH, M. R.: Recent USAF Experience with Inflight Dysbarism. *Aerospace Med.*, 33:995, 1962.
- BRUNEAU, R. J., and HERTZ, H. W.: Physiological Support of an Extreme High Altitude Flying Program. *Aerospace Med.*, 33:436, 1962.
- HAYMAKER, W., JOHNSTON, A. D., and DOWNEY, V. M.: Fatal Decompression Sickness During Jet Aircraft Flight. *J. Aviation Med.*, 27:2, 1956.
- HORNBERGER, W.: Decompression Sickness. In *German Aviation in World War II*, p. 354-394. The Surgeon General, U. S. Air Force Department of the Air Force, 1950.
- LEMESURIER, D. H., and BAXTER, R.: Evidence of Decompression Sickness During Flights at 20,000 to 24,000 Feet Above Base. *Med. J. Australia*, 1:188, 1964.
- ODLAND, L. T.: Fatal Decompression Sickness at an Altitude of 22,000 Feet. *Aerospace Med.*, 30:840, 1959.
- TOBIAS, C. A.: Decompression Sickness in Actual Flights. In *Fulton, J. F. (ed.). Decompression Sickness*, p. 360-377. Philadelphia: W. B. Saunders, 1951.

UNCLASSIFIED

Security Classification

DOCUMENT CONTROL DATA - R&D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) USAF School of Aerospace Medicine Aerospace Medical Division (AFSC) Brooks Air Force Base, Texas		2a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED <i>346</i>	
		2b. GROUP	
3. REPORT TITLE Decompression Sickness in High-altitude Flight			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)			
5. AUTHOR(S) (Last name, first name, initial) Meader, Willard L., Major, USAF, MC			
6. REPORT DATE March 1967		7a. TOTAL NO. OF PAGES 3	7b. NO. OF REFS 9
8a. CONTRACT OR GRANT NO. b. PROJECT NO. c. d.		8a. ORIGINATOR'S REPORT NUMBER(S) SAM TR 66-305 8b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
10. AVAILABILITY/LIMITATION NOTICES Distribution of this document is unlimited.			
11. SUPPLEMENTARY NOTES Reprinted from Aerospace Medicine 38(3): 301-303 (1967)		12. SPONSORING MILITARY ACTIVITY USAF School of Aerospace Medicine Aerospace Medical Division (AFSC) Brooks Air Force Base, Texas	
13. ABSTRACT Review of 5 years of a WU-2 squadron's daily records shows 36 cases of decompression sickness entirely confined to reports of bends pain among 11 crew-members in 958 flights. Generally, 40 minutes were spent in denitrogenation in a partial pressure suit and helmet before achieving cabin altitudes of close to 29,000 feet. The majority of bends occurred within the first 3.5 hours affecting the knee joints in almost two-thirds of the cases with a tendency to recur at the same joint. Only 1 case required descent to lower altitude to afford relief from pain.			

DD FORM 1473
1 JAN 64

UNCLASSIFIED

Security Classification

UNCLASSIFIED
Security Classification

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Physiology						
Decompression sickness						
Bends						

INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (*corporate author*) issuing the report.

2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. **REPORT DATE:** Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.

7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.

8a. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. **OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (*either by the originator or by the sponsor*), also enter this number(s).

10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through _____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (*paying for*) the research and development. Include address.

13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. **KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, rules, and weights is optional.