

AD 678366

ACUTE MORTALITY OF CHINCHILLAS EXPOSED TO MIXED
GAMMA-NEUTRON RADIATIONS OR 250 KVP X RAYS

T. A. STRIKE
L. J. SEIGNEUR


R. E. GEORGE
Commander, MSC, USN
Chairman
Radiation Biology Department


HUGH B. MITCHELL
Colonel, USAF, MC
Director

ARMED FORCES RADIOBIOLOGY RESEARCH INSTITUTE
Defense Atomic Support Agency
Bethesda, Maryland

Distribution of this document is unlimited

ACKNOWLEDGMENT

The authors wish to express their gratitude to the members of the National Chinchilla Breeders of America, Inc. who supported this study by their generous donation of the animals. The authors also gratefully acknowledge the technical assistance provided during the course of this study by B. L. Wilhelm and R. H. Crutcher.

TABLE OF CONTENTS

	Page
Foreword (Nontechnical summary)	iii
Abstract	v
I. Introduction	1
II. Materials and Methods	1
III. Results	8
IV. Discussion	11
V. Summary	13
References	15

LIST OF FIGURES

	Page
Figure 1. Chinchilla phantom showing placement of miniature tissue equivalent ionization chambers	3
Figure 2. Dose profiles in unilaterally exposed chinchilla phantoms .	4
Figure 3. X-ray exposure array for chinchillas	5
Figure 4. Reactor exposure array for chinchillas	6
Figure 5. Dose-response regression lines and associated 95 percent confidence bands for the chinchilla as calculated by probit analysis	8
Figure 6. Survival time regression lines and associated 95 percent confidence limits for the chinchilla	9

LIST OF TABLES

Table I. Chinchilla Mortality Data	2
Table II. Probit Analysis of Chinchilla 30-Day Mortality Data . .	9
Table III. Dose-Response Data of Rats and Chinchillas Exposed to Supralethal Doses of x rays	10
Table IV. Dose-Response Data of Rats and Chinchillas Exposed to Supralethal Doses of Mixed Gamma-Neutron Radiations	10

FOREWORD
(Nontechnical summary)

The acute mortality of chinchillas exposed to ionizing radiation was studied at the Armed Forces Radiobiology Research Institute (AFRRI) as part of an effort to characterize the biological effect of mixed gamma-neutron radiations from the AFRRI-TRIGA reactor.

Chinchillas were exposed to the mixed gamma-neutron radiations or to 250 kVp x rays at selected doses throughout the lethal range (that range of doses resulting in death of from 1 to 99 percent of the exposed animals within 30 days). In addition, some animals were irradiated at higher doses to obtain survival time data in the dose range expected to cause 100 percent mortality within 10 days. All exposures were whole body, unilateral, and delivered at approximately 20 rads/min. Midline tissue doses ranged from 200 to 639 rads for mixed gamma-neutron radiations and from 357 to 1786 rads for x rays.

Chinchilla deaths were recorded daily and the resulting data subjected to mortality and survival time analysis. A statistical method (probit analysis) was used to obtain dose-mortality response curves and associated parameters including the median lethal dose (that dose which will kill 50 percent of the animals in a large group).

In this study the 30-day median lethal doses ($LD_{50/30}$) for the mixed gamma-neutron radiations and for 250 kVp x rays were calculated to be 295 and 490 rads, respectively. By comparing the $LD_{50/30}$ values, the RBE (Relative Biological Effectiveness) of the reactor radiations was found to be 1.7.

In previous radiation lethality studies, survival time of Sprague-Dawley rats was reduced from approximately 10 days to less than 5 days by increasing the dose from the lethal range to about twice the LD_{50/30}. This result was not obtained when chinchillas were similarly treated.

ABSTRACT

The chinchilla's acute mortality response to mixed gamma-neutron radiations of the AFRRI-TRIGA reactor and to 250 kVp x rays was studied. Unilateral whole body irradiations were accomplished at doses from 200 to 639 rads of mixed gamma-neutron radiations and from 357 to 1786 rads of x rays. All radiations were delivered at approximately 20 rads/min, and doses are reported as midline tissue doses. The $LD_{50/30}$ values calculated for the mixed gamma-neutron radiations and for the x rays were 295 and 490 rads, respectively. Using 30-day median lethality as the end point, the RBE of the mixed gamma-neutron radiations was 1.7. The wide lethal dose range obtained was attributed to a high degree of variation in age of the chinchillas. In contrast to previous experience in rodents, the chinchilla showed a relative resistance to the classically described gastrointestinal modality of radiation death.

I. INTRODUCTION

A comparison study was initiated among several mammalian species to assess the biological effectiveness of mixed gamma-neutron radiations from the AFRRI-TRIGA Mark F reactor. The reference radiation was 250 kVp x rays, and the 30-day median lethal dose ($LD_{50/30}$) was selected as the biological end point for comparison. In the course of this research, an opportunity arose to study the response of chinchillas to ionizing radiations.

The chinchilla has many unique anatomical and physiological characteristics, therefore, its mortality response to ionizing radiations seemed especially worthy of study. This rodent has a 28-day estrous cycle and a 111-day gestation period,¹² a 12- to 20-year lifespan,⁷ fine hair that can be painlessly plucked with as many as 50-60 hairs exiting from each hair follicle,²⁵ and a metabolism which results in odorless urine and feces.¹⁴

The mortality results of chinchilla exposures to mixed gamma-neutron radiations from the TRIGA reactor or x rays are the subject of this report.

II. MATERIALS AND METHODS

Adult chinchillas* of Taniger strain were collected locally† from ranches throughout the United States. After a minimum conditioning period of 12 weeks, the chinchillas were transferred to environment-controlled animal rooms at this laboratory. They were individually caged and acclimatized an additional 4 weeks before

* These chinchillas represent stock discarded by the breeders because of their "fur chewing" tendencies. In all other respects these animals were normal and healthy.

† National Chinchilla Breeders of America, Inc., Chinchilla Industry Testing Center, Bethesda, Maryland

being irradiated. During this period, the chinchillas were weighed twice weekly and those animals exhibiting weight loss or symptoms of disease were excluded from the study. Food and water were available ad libitum. The diet consisted of guinea pig chow* and rough clover hay, supplemented twice weekly with apples.

In the initial series of exposures, a total of 512 chinchillas was used (Table I). Animals were assigned to dose groups in a random fashion, biased only by the fact that each group of 32 was equally divided as to sex. To test the reproducibility of the initial results, an additional 229 chinchillas were grouped randomly (16 or 20 animals

Table I. Chinchilla Mortality Data

MIDLINE TISSUE DOSE (RADS)*	DIED EXPOSED	POST-IRRADIATION DAY OF DEATH**																														PERCENT MORTALITY (30 DAYS)	SURVIVAL TIME (DAYS)		
		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30		MEAN	MEDIAN	
X RAYS																																			
0	0 32																																0		
0	0 16																																0		
12.5	3 16																																19	15.0	14.0
44.7	16 32																																30	17.7	17.5
53.6	17 32																																52	12.5	14.0
56	10 15																																67	11.4	12.0
62.5	24 32																																75	12.4	12.5
67.0	21 32																																84	12.1	11.0
71.4	25 32																																78	12.0	12.0
71	14 16																																68	12.3	12.7
73.0	24 32																																75	12.5	12.0
80.4	28 32																																91	11.2	12.0
84.0	31 32																																87	10.6	11.0
89.3	27 32																																84	10.3	10.0
91	15 15																																100	9.2	9.0
100	16 16																																100	7.3	7.0
100	16 16																																100	6.2	6.0
MIXED GAMMA-NEUTRON RADIATIONS																																			
0	0 32																																0		
0	0 20																																0		
200	5 32																																16	12.4	13.5
200	4 20																																40	13.9	13.5
200	14 20																																70	12.0	10.0
301	10 12																																31	12.0	12.0
325	17 20																																82	9.6	8.0
429	17 32																																53	11.6	11.0
477	20 30																																100	7.7	6.5
537	30 32																																90	11.2	11.5
597	16 16																																100	7.8	7.0
535	7 32																																100	6.5	6.0

*BOLD TYPE DENOTES THE INITIAL EXPOSURES AND ITALIC THE SECOND EXPOSURE SERIES
** FIGURES IN POSTIRRADIATION DAY OF DEATH COLUMNS INDICATE NUMBER OF DEATHS ON THAT DAY

* Ralston Purina Co., St. Louis, Missouri

per group) and were similarly irradiated (Table I). Exposures were extended into the supralethal dose range to permit a comparison with results obtained by exposing rats to comparable doses.

At the time of irradiation, the ages of the chinchillas ranged from 24 weeks to approximately 8 years* and their weights varied from 115 to 644 grams. All exposures were unilateral and employed minimal scatter conditions. Plexiglas† boxes (3-7/8" wide x 8-7/8" long x 5" high), constructed from sheets 1/8" thick, were used to confine the chinchillas during exposure. Depth dose measurements were made in chinchilla phantoms (Figure 1) using miniature tissue equivalent (T.E.) ionization chambers.⁶ The phantoms were fabricated from Plexiglas tubing (3-1/4" O.D. - 2-7/8" I.D.) and filled with tissue equivalent fluid.¹⁷ The phantoms were representative of




Figure 1. Chinchilla phantom showing placement of miniature tissue equivalent ionization chambers

* The precise ages of the chinchillas were unknown. The reported age range is based on the available growth curves¹⁹ and estimated longevity⁹ of these rodents.

† Acrylic plastic. Rohm and Haas Co., Philadelphia, Pennsylvania

the average weight and size of the chinchillas in this study. Two longitudinal grooves, diametrically opposing each other, were centered on the surface of the phantom. Each groove held three miniature T. E. chambers used to measure entrance or exit doses. Depth dose measurements at 1/4, 1/2, and 3/4 the diameter of the phantom were made with T.E. chambers positioned in Plexiglas tubes (5/16" O.D. - 3/16" I.D.) traversing the length of the cylinder. All depth dose measurements in the phantoms were made using the same exposure conditions as for the animal irradiations.

The results of depth dose measurements made in the chinchilla phantoms are shown in Figure 2. Categorizing the irradiations according to degree of uniformity of absorbed dose within the volume of interest,¹¹ both the x-ray and mixed gamma-neutron irradiations were Class B nonuniform exposures.




Figure 2. Dose profiles in unilaterally exposed chinchilla phantoms

The x-ray exposures were accomplished with the 360° radial beam from a 250 kVp x-ray generator operated at 30 mA. The inherent (1.2 mm beryllium) and added (0.95 mm copper) filtration resulted in a half value layer (HVL) of 1.9 mm of copper. Exposure boxes were placed in the radiation field so that the midline of each animal was 1 meter from the x-ray target (Figure 3).




Figure 3. X-ray exposure array for chinchillas

The absorbed dose at the center of the animal was calculated from three factors. First, a Victoreen Roentgen Chamber was used to determine the exposure, free-in-air, at the position to be occupied by the center line of the animal. Positions were selected so that the variation in this quantity was less than 4 percent from the mean. Second, the ratio of the dose at the midline of a phantom (simulating the

chinchilla in the irradiation position) to the exposure free-in-air, was obtained using miniature ionization chambers. Third, the conversion factor ($\bar{f}$) of 0.95 for muscle and for this quality of radiation was obtained from the ICRU Report 10b.¹⁰ The product of these three factors gave the absorbed dose at the center line of the animal. The dose rate in all exposures was approximately 21 rads/min. Dose rate was monitored continuously during each exposure with a Victoreen rate meter in order to detect any changes in the output of the x-ray unit.

Figure 4 illustrates the array used for the exposures to mixed gamma-neutron radiations. The exposure boxes were positioned so that the midline of each animal was on an arc 292 cm from the vertical center line of the reactor core. This arc, located approximately in the middle of the exposure room, was in an exposure field in which the tissue kerma, free-in-air, did not vary by more than 4 percent from the mean.



Figure 4. Reactor exposure array for chinchillas

The absorbed dose at the midline of the animal was calculated from two factors. First, the tissue kerma, free-in-air, was calculated from measurements with a 50 cm³ cavity tissue equivalent plastic* walled ionization chamber. Second, the ratio of the absorbed dose in the center of the phantom (simulating the chinchilla in the irradiation position) to the tissue kerma, free-in-air, was obtained. The product of these two quantities gave the absorbed dose at the center line of the animal. The absorbed dose rate for all exposures was approximately 19 rads/min.

Approximately 60 percent of the tissue kerma, free-in-air, is attributed to gamma rays, 30 percent to neutrons of energies greater than 10 keV and 10 percent to neutrons of lower energies. The effective energy† of the gamma component was between 1 and 2 MeV. Details of the reactor characteristics and methods of dosimetry used in this mixed radiation field have been previously described.^{8,20}

Chinchilla deaths were recorded daily for 30 days following irradiation. The LD_{50/30} values were calculated by subjecting the resulting data to probit analysis using a maximum likelihood method programmed for a digital computer. The resultant regression lines from the initial and second series of exposures were tested for homogeneity and parallelism, and the LD_{50/30} values were tested for differences. The mean survival time for the decedents of each group was plotted and a "least squares" method employed to determine the best fitting lines. The 95 percent

* Plastic supplied by Dr. F. R. Shonka, St. Procopius College, Lisle, Illinois. (Composition by weight: 76.1 percent carbon, 10.1 percent hydrogen, 5.2 percent oxygen, 3.5 percent nitrogen, 1.0 percent silicon, 2.0 percent calcium and 2.0 percent fluorine.)

† Depth dose studies, using Plexiglas rat phantoms, indicate that the deposition of energy by the gamma component of the reactor radiations was similar to that of ⁶⁰Co gamma rays.

confidence band for each regression line as a whole was computed using the method described by Natrella.¹⁵

No significant differences were found when the dose response regression lines and LD_{50/30} values from the initial exposures were tested against their counterparts from the second series of exposures, nor was any significant difference found between the radiosensitivity of males and females. Since the results of the initial and the second series of exposures were similar, the data were combined and analyzed to simplify presentation.

III. RESULTS

Table I on page 2 summarizes the mortality data for the chinchillas. The raw data used for probit analysis and the resultant dose-response regression lines with their 95 percent confidence bands are displayed graphically in Figure 5.




Figure 5. Dose-response regression lines and associated 95 percent confidence bands for the chinchilla as calculated by probit analysis. Plotted points represent raw data and arrows indicate the doses associated with 100 percent mortality.

The results of probit analysis and the calculated relative biological effectiveness (RBE) for the mixed gamma-neutron radiations are shown in Table II. The mean survival time regression lines and their 95 percent confidence bands for chinchillas exposed to x rays and mixed gamma-neutron radiations are shown in Figure 6.

Table II. Probit Analysis of Chinchilla 30-Day Mortality Data

Radiation source	Calculated lethal dose values*					Slope of regression line	RBE† for LD ₅₀ /30's
	LD ₁₀	LD ₃₀	LD ₅₀	LD ₇₀	LD ₉₀		
x ray	260 (209-335)‡	390 (323-438)	490 (435-531)	616 (574-658)	858 (780-971)	5.3	-
mixed gamma-neutron radiations	168 (69-227)	234 (137-292)	295 (212-360)	370 (299-487)	515 (413-899)	5.3	1.7

* Midline tissue dose (rads)

† 250 kVp x rays used as standard reference source

‡ 95 percent confidence limits shown in parentheses




Figure 6. Survival time regression lines and associated 95 percent confidence limits for the chinchilla. The plotted points represent the mean values for each dose group listed in Table I.

In Table III, the mortality response of chinchillas exposed to supralethal doses of x rays is compared with the respective mortality response previously obtained in rats. A similar comparison after exposure to mixed gamma-neutron radiations is shown in Table IV.

Table III. Dose-Response Data of Rats and Chinchillas Exposed to Supralethal Doses of x rays

MIDLINE TISSUE DOSE (RADS)	NUMBER OF ANIMALS	POSTIRRADIATION DAY OF DEATH					30-DAY PERCENT MORTALITY	MEDIAN SURVIVAL TIME (DAYS)
		5	10	15	20	25		
2052	48	41 7					100.0	4.0
1231	48	44 4					100.0	4.0
1026	48	21 13	5 4	1 2			100.0	5.0
1785	16	1 7	4				100.0	6.0
1072	16	3 2	3 5	1 1			100.0	7.0
893	47	1 1	2 6	4 7	6 2	3 5	89.4	9.5

Table IV. Dose-Response Data of Rats and Chinchillas Exposed to Supralethal Doses of Mixed Gamma-Neutron Radiations

MIDLINE TISSUE DOSE (RADS)	NUMBER OF ANIMALS	POSTIRRADIATION DAY OF DEATH					30-DAY PERCENT MORTALITY	MEDIAN SURVIVAL TIME (DAYS)
		5	10	15	20	25		
734	48	24 24					100.0	4.5
649	48	16 30	2				100.0	5.0
548	48	2 3	13 9	3 4	1 3		100.0	7.0
639	32	1 1	2 12	2 4	6 2	1 2	100.0	8.0
601	19	1 3	4 2	1 1	4 2	1	100.0	7.0
531	32		2 1	2 2	1 9	3 8	93.8	11.0

IV. DISCUSSION

Dose-mortality and survival-time values for the chinchilla indicate that the response of this rodent to doses of ionizing radiations in the lethal range is similar to that of mice and rats.²² The lethal dose range for the chinchillas used in this study was much wider than for mice and rats. The increased width of the lethal range indicates that the chinchillas represented a heterogeneous population. This heterogeneity is reflected in the slopes of the probit regression lines (Figure 5) and the width of the associated 95 percent confidence bands (Table II). The slopes of the probit regression lines are less than half of the values reported for rats²² and mice.^{2,22} It has been shown that the radiosensitivity of mice changes with age.¹ Assuming that this is also true for other rodents, the wide range of ages of the chinchillas in this study could account for much of the heterogeneity that was observed.

The 16 early deaths (prior to the 5th postirradiation day) observed in chinchillas exposed to x rays or to mixed gamma-neutron radiations (Table I) in the lethal dose range apparently resulted from a natural characteristic of the species, rather than an effect of radiation. Chinchillas are extremely sensitive to some forms of stress. An animal may appear quite normal and within minutes die from shock induced by conditions ordinarily not considered lethally stressful. The gross pathology seen at the necropsy of animals dying early was similar to that described by T-W- Fiennes²³ for chinchillas which died from shock and was not characteristic of radiation injury. These early deaths were included in the probit analysis but had no significant effect on the mortality values calculated in this study.

A unique characteristic of the chinchilla found in this study is its apparent radioresistance to the gastrointestinal modality of death when compared to the Sprague-Dawley rat. For example, it has been our repeated observation that Sprague-Dawley rats are, within the 500- to 700-rad dose range of mixed gamma-neutron radiations (Table IV), quite susceptible to the gastrointestinal modality of death. (Other investigators have made similar observations in mice following exposure to neutron or modified fission spectrum radiation doses of less than 500 rads.^{3,24}) However, the chinchilla did not demonstrate this susceptibility when subjected to the same mixed gamma-neutron radiations and comparable dose range. The resistance of the chinchilla to the gastrointestinal mode of death was tested further by irradiating animals at doses as high as 1786 rads of 250 kVp x rays (Table III). The mortality results indicated that the supralethal doses of x rays used did not shift the chinchilla deaths into the gastrointestinal temporal range. The mortality response of the chinchilla in the supralethal dose range does not agree with the results reported for mice and rats wherein classical gastrointestinal deaths are found in small rodents exposed to 1000 R or more of gamma or x rays.^{4,13,16,18}

Mean survival time for the decedents of each exposed group was plotted against dose and regression lines fitted to the data points for each radiation type (Figure 6). For those portions of the regression lines where equal doses can be compared, the chinchillas exposed to radiations from the reactor have significantly shorter survival times than the exposed to x rays. Similar results were reported for mice^{3,5,22,24} and rats.²²

V. SUMMARY

Chinchillas were unilaterally exposed to whole body doses of mixed gamma-neutron (reactor) radiations or 250 kVp x rays. Dose rates were approximately 20 rads/min. Midline tissue doses from 200 to 639 rads of mixed gamma-neutron radiations and from 357 to 1786 rads of x rays were used. The $LD_{50/30}$ values were calculated to be 290 rads for the mixed gamma-neutron radiations and 490 rads for the x rays. Using the $LD_{50/30}$ value as the end point, the relative biological effectiveness (RBE) of the reactor radiations was 1.7. The heterogeneity in age of the chinchillas was interpreted as being responsible for the rather wide variations in mortality response. At comparable supralethal doses, chinchilla survival times were not as markedly reduced as the rat survival times.

REFERENCES

1. Abrams, H. L. Influence of age, body weight and sex on susceptibility of mice to the lethal effects of X-radiation. *Proc. Soc. Exptl. Biol. Med.* 76:729-737, 1951.
2. Ainsworth, E. J., Leong, C. F., Kendall, K., and Alpen, E. L. The lethal effects of pulsed neutron or gamma irradiation in mice. *Radiation Res.* 31: 75-85, 1964.
3. Bond, V. P. Comparison of the mortality response of different mammalian species to X-rays and fast neutrons. In: *Biological Effects of Neutron and Proton Irradiations*, Vol. II, pp. 365-377. Vienna, International Atomic Energy Agency, 1964.
4. Bonét-Maury, P. and Patti, F. Lethal irradiation of mice with high doses of roentgen and gamma rays. *Radiology* 57:419-423, 1951.
5. Carter, R. E., Bond, V. P., and Seymour, P. H. The relative biological effectiveness of fast neutrons in mice. *Radiation Res.* 4:413-423, 1956.
6. Chambers, F. W., Jr. Miniature tissue equivalent ionization chambers and their use. *Aerospace Med.* 34:193-196, 1963.
7. Crandall, L. S. *Management of Wild Mammals in Captivity*. Chicago, Illinois, University of Chicago Press, pp. 254-256, 1964.
8. Dowling, J. H. Experimental determination of dose for the monkey in a reactor pulse environment. Bethesda, Maryland, Armed Forces Radiobiology Research Institute Scientific Report SR66-3, 1966.
9. Graham-Jones, O. Clinical approach to chinchillas and monkeys. *British Small Animal Vet. Assoc. Congress Proc.*, 1959 (Jones, B., editor), pp. 113-124. London, Pergamon Press Ltd., 1960.
10. International Commission on Radiological Units and Measurements (ICRU) Report 10b, 1962. Physical aspects of irradiation. National Bureau of Standards Handbook No. 85. Washington, D. C., U. S. Government Printing Office, 1964.
11. International Commission on Radiological Units and Measurements (ICRU) Report 10e, 1962. Radiobiological Dosimetry. National Bureau of Standards Handbook No. 88. Washington, D. C., U. S. Government Printing Office, 1963.

12. Kitchen, H. and Kitchen, Y. Reproduction, breeding and management of chinchillas. *Lab. Animal Digest* 3(4):3-7, October 1967.
13. Langham, W., Woodward, K. T., Rothermel, S. M., Harris, P. S., Lushbaugh, C. C., and Storer, J. B. Studies of the effect of rapidly delivered massive doses of gamma-rays on mammals. *Radiation Res.* 5:404-432, 1956.
14. McKay, C. M. Chinchillas. *Lancet* 2:350, 1959.
15. Natrella, M. G. Experimental Statistics. National Bureau of Standards Handbook No. 91, Washington, D. C., U. S. Government Printing Office, 1963.
16. Quastler, H. Studies on roentgen death in mice. I. Survival time and dosage. *Am. J. Roentgenol.* 54:449-456, 1945.
17. Rossi, H. H. and Failla, G. Tissue-equivalent ionization chambers. *Nucleonics* 14(2):32-37, February 1956.
18. Rothermel, S. M., Woodward, K. T., and Storer, J. B. The effect of massive doses of neutrons on the median survival time of mice. *Radiation Res.* 5:433-440, 1956.
19. Rubacky, E. P. Weighing chinchillas. Chinchilla Fur Farming Series, File 80.90 pp. 1-4, October 1963.
20. Sayeg, J. A., compiler. Neutron and gamma dosimetry measurements at the AFRRI-DASA TRIGA Reactor. Bethesda, Maryland, Armed Forces Radiobiology Research Institute Contract Report CR65-6, 1965 (originally issued as Edgerton, Germeshausen and Grier, Inc. Report S-260-R, Santa Barbara, California, 1964).
21. Strike, T. A. (unpublished results).
22. Strike, T. A., Seigneur, L. J., and Stanley, R. E. Acute mortality of mice and rats exposed to mixed gamma-neutron radiations or to x rays. Bethesda, Maryland, Armed Forces Radiobiology Research Institute Report SR68-6 (in press).
23. T-W-Fiennes, R. N. Observations on the pathology of chinchillas and monkeys. *British Small Animal Vet. Assoc. Congress Proc.*, 1959 (Jones, B., editor) pp. 108-112. London, Pergamon Press Ltd., 1960.
24. Vogel, H. H., Jr., Clark, J. W., and Jordan, D. L. Comparative mortality following single whole-body exposures of mice to fission neutrons and Co^{60} gamma rays. *Radiology* 68:386-398, 1957.
25. Wilcox, H. W. Histology of the skin and hair of the adult chinchilla. *Anat. Rec.* 108:385-397, 1950.

UNCLASSIFIED
Security Classification

DOCUMENT CONTROL DATA - RSP		
(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)		
1 ORIGINATING ACTIVITY (Corporate author) Armed Forces Radiobiology Research Institute Defense Atomic Support Agency Bethesda, Maryland 20814		2a REPORT SECURITY CLASSIFICATION UNCLASSIFIED
3 REPORT TITLE ACUTE MORTALITY OF CHINCHILLAS EXPOSED TO MIXED GAMMA-NEUTRON RADIATIONS OR 250 KVP X RAYS		2b GROUP N/A
4 DESCRIPTIVE NOTES (Type of report and inclusive dates)		
5 AUTHOR(S) (Last name, first name, initial) Strike, T. A. and Seigneur, L. J.		
6 REPORT DATE April 1968	7a TOTAL NO OF PAGES 24	7b NO OF REFS 25
8a CONTRACT OR GRANT NO.	9a ORIGINATOR'S REPORT NUMBER(S) AFRRI SR68-7	
b PROJECT NO. c R MD 3 9002	9b OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
10 AVAILABILITY/LIMITATION NOTICES Distribution of this document is unlimited		
11 SUPPLEMENTARY NOTES	12 SPONSORING MILITARY ACTIVITY Defense Atomic Support Agency Washington, D. C. 20305	
13 ABSTRACT The chinchilla's acute mortality response to mixed gamma-neutron radiations of the AFRRI-TRIGA reactor and to 250 kVp x rays was studied. Unilateral whole body irradiations were accomplished at doses from 200 to 639 rads of mixed gamma-neutron radiations and from 357 to 1786 rads of x rays. All radiations were delivered at approximately 20 rads/min, and doses are reported as midline tissue doses. The LD _{50/30} values calculated for the mixed gamma-neutron radiations and for the x rays were 295 and 490 rads, respectively. Using 30-day median lethality as the end point, the RBE of the mixed gamma-neutron radiations was 1.7. The wide lethal dose range obtained was attributed to a high degree of variation in age of the chinchillas. In contrast to previous experience in rodents, the chinchilla showed a relative resistance to the classically described gastrointestinal modality of radiation death.		

DD FORM 1 JAN 64 1473

UNCLASSIFIED
Security Classification

UNCLASSIFIED
Security Classification

KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
acute mortality lethality, median RBE mixed gamma-neutron radiations chinchillas						

INSTRUCTIONS

1. ORIGINATING ACTIVITY: Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (*corporate author*) issuing the report.

2a. REPORT SECURITY CLASSIFICATION: Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. GROUP: Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. REPORT TITLE: Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parentheses immediately following the title.

4. DESCRIPTIVE NOTES: If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. AUTHOR(S): Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. REPORT DATE: Enter the date of the report as day, month, year; or month, year. If more than one date appears on the report, use date of publication.

7a. TOTAL NUMBER OF PAGES: The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. NUMBER OF REFERENCES: Enter the total number of references cited in the report.

8a. CONTRACT OR GRANT NUMBER: If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, 8d. PROJECT NUMBER: Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. ORIGINATOR'S REPORT NUMBER(S): Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. OTHER REPORT NUMBER(S): If the report has been assigned any other report numbers (*either by the originator or by the sponsor*), also enter this number(s).

10. AVAILABILITY/LIMITATION NOTICES: Enter any limitations on further dissemination of the report, other than those imposed by security classification, using standard statements such as:

(1) "Qualified requesters may obtain copies of this report from DDC."

(2) "Foreign announcement and dissemination of this report by DDC is not authorized."

(3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."

(4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."

(5) "All distribution of this report is controlled. Qualified DDC users shall request through _____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. SUPPLEMENTARY NOTES: Use for additional explanatory notes.

12. SPONSORING MILITARY ACTIVITY: Enter the name of the departmental project office or laboratory sponsoring (paying for) the research and development. Include address.

13. ABSTRACT: Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS) (S) (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. KEY WORDS: Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical content. The assignment of links, rules, and weights is optional.