

AD 685029


FOREIGN TECHNOLOGY DIVISION



SYNTHESIS OF CERTAIN HOMOLOGS OF ADAMANTANE FROM
ADAMANTANE CARBOXYLIC-1 ACID

by

S. Landa, Y. Burkhard, and Y. Weiss



Distribution of this document is unlimited. It may be released to the Clearinghouse, Department of Commerce, for sale to the general public.

Reproduced by the
CLEARINGHOUSE
for Federal Scientific & Technical
Information Springfield Va. 22151

EDITED TRANSLATION

SYNTHESIS OF CERTAIN HOMOLOGS OF ADAMANTANE
FROM ADAMANTANE CARBOXYLIC-1 ACID

By: S. Landa, Y. Burkhard, and Y. Weiss

English pages: 6

Source: Neftekhimiya (Fuel Chemistry),
Vol. 8, No. 3, 1968, pp. 323-325.

Translated by: D. Koolbeck/TDBRO-2

THIS TRANSLATION IS A RENDITION OF THE ORIGINAL FOREIGN TEXT WITHOUT ANY ANALYTICAL OR EDITORIAL COMMENT. STATEMENTS OR THEORIES ADVOCATED OR IMPLIED ARE THOSE OF THE SOURCE AND DO NOT NECESSARILY REFLECT THE POSITION OR OPINION OF THE FOREIGN TECHNOLOGY DIVISION.

PREPARED BY:

TRANSLATION DIVISION
FOREIGN TECHNOLOGY DIVISION
WP-APB, ONIS.

DATA HANDLING PAGE

01-ACCESSION NO. TP8002081		98-DOCUMENT LOC		39-TOPIC TAGS lithium, hydrocarbon, gas chromatography	
09-TITLE SYNTHESIS OF CERTAIN HOMOLOGS OF ADAMANTANE FROM ADAMANTANE CARBOXYLIC-1 ACID					
47-SUBJECT AREA 07					
43-AUTHOR/CO-AUTHORS LANDA, S.; 16-BURKHARD, Y.; 15-WEISS, Y.				10-DATE OF INFO -----68	
43-SOURCE NEFTEKHIMIYA (RUSSIAN)				FTD- 66-DOCUMENT NO. HT-23-1007-68	
				69-PROJECT NO. 50403	
63-SECURITY AND DOWNGRADING INFORMATION UNCL, 0			64-CONTROL MARKINGS NONE		97-HEADER CLASS UNCL
76-REEL/FRAME NO. 1887 1866	77-SUPERSEDES	78-CHANGES	40-GEOGRAPHICAL AREA UR	NO OF PAGES 6	
CONTRACT NO.	X REF ACC. NO. 65-	PUBLISHING DATE 94-00	TYPE PRODUCT TRANSLATION	REVISION FREQ NONE	
STEP NO. 02-UR/0204/68/008/003/0323/0325			ACCESSION NO.		


ABSTRACT

The reaction of alkyl lithium with adamantane carboxylic-1 acid is used to synthesize ketones, which are then reduced by the Clemmenson method to produce homologs of adamantane. Also, carbinols obtained as by-products of the initial reactions are reduced by the Clemmensen method to produce a series of hydrocarbons. It is stated that the method outlined in the article gives higher yields of reaction products than previously published procedures. Also, the previously indescribed 1-tret-kutyl adamantane was synthesized. Orig. art. has: 5 formulas and 2 tables.

**SYNTHESIS OF CERTAIN HOMOLOGS OF ADAMANTANE FROM
ADAMANTANE CARBOXYLIC-1 ACID**

S. Landa, Y. Burkhard, and Y. Weiss

The literature contains little information about the synthesis of ketones from free acids and alkyl lithium [1]. It is known that the yield of ketones from the reactions of lower carboxylic acids is low. A higher yield is obtained in the reaction of acids in which the number of carbon atoms is greater than five. We found that adamantane carboxylic-1 acid reacts well with R'Li, as is shown on diagram (1):




However, in this case besides the desired ketone there is incidental formation of a small quantity of tertiary alcohol [1, 2], especially when a large excess of alkyl lithium is applied (diagram 2):



The reaction mixture was reduced by the Clemmensen method:



During reduction the olefin and corresponding saturated hydrocarbon are formed from the tertiary alcohol produced during the reaction:




We did not examine the mechanism of formation of these hydrocarbons in detail.

Identification of the by-products of the reaction was conducted by gas-liquid chromatography, by comparison with synthetic standard preparations. Separation of the hydrocarbon mixture was realized with the help of preparative gas-liquid chromatography.

Only the reaction of adamantane carboxyl-1 acid with n-alkyl lithium gives a satisfactory yield; the reaction does not proceed with isopropyl lithium.

The reactions described in this work are suitable in general for syntheses of 1-n-alkyl adamantanes. In contrast to works published earlier [3, 4], the method ensures a higher yield of reaction products.

The following method was used to synthesize tert-butyl adamantane: the reaction of the methyl ester of adamantane carboxylic-1 acid with methyl magnesium iodide yielded 2-(1-adamantyl)-propanol-2, from which the corresponding bromide was synthesized with thionyl bromide; the bromide was then transformed into tert-butyl adamantane during reaction with methyl magnesium iodide. However, the content of this hydrocarbon in the reaction products amounted to only 22%; the remaining 78% was 2-(1-adamantyl)-propene. Since 2-(1-adamantyl)-2-bromopropane introduced into the reaction [chart(5)] was not analyzed, but was only purified of excess thionyl bromide, it is difficult to say whether 2-(1-adamantyl)-propene appears during the synthesis of the bromide or during the reaction of the bromide with methyl magnesium iodide.



Separation of the reaction mixture was accomplished with the help of preparative gas-liquid chromatography.

Experimental Section

Alkyl lithium was prepared by the usual method [5]. The reaction with adamantane carboxylic-1 acid was conducted in absolute ether in a nitrogen atmosphere at a temperature from -40 to -35°C . Yields of reaction products are given in Table 1. Properties of the synthesized hydrocarbons are shown in Table 2.

Table 1. Yield of reaction products.

Alkyl lithium	Yield of wet ketone, calc. on acid taken, wt. %	Content of tertiary alcohol in wet ketone, mol. %	Unreacted acid, wt. %
CH_3Li	80.3	5.0	15.8
$\text{C}_2\text{H}_5\text{Li}$	36.7	20.0	41.5
$n\text{-C}_3\text{H}_7\text{Li}$	74.8	6.0	9.0
$\text{iso-C}_3\text{H}_7\text{Li}$	—	—	94.4

Table 2. Properties of synthesized hydrocarbons

Hydrocarbon	T _{melt} °C	T _{boil} °C	n _D ²⁰	Calc., %		Found, %	
				C	H	C	H
1-ethyl adamantane	-57.7	218	1.4950	87.73	12.27	87.80	12.22
1-n-propyl adamantane	+7.5	218.2*[6]	1.4950*	87.56	12.44	87.61	12.47
1-n-butyl adamantane	-12.0	253	1.4923	87.43	12.57	87.46	12.51
3-(1-adamantyl)-pentane	not determined	262*[7]	1.4917*	87.30	12.70	87.41	12.78
3-(1-adamantyl)-pentene-2	not determined	not determined	1.5000*[8]	88.16	11.84	88.22	11.87
1-tret-butyl adamantane	111.7- 113.0	-	-	87.43	12.57	87.49	12.53
2-(1-adamntyl)-propene	not determined	not determined	1.5106	88.56	11.44	88.62	11.46
*Literature data.							

Conclusions

Methyl-(1-adamantyl)-, ethyl-(1-adamantyl)-, and n-propyl-(1-adamantyl)-ketones were synthesized by the reaction of adamantane carboxylic-1 acid with alkyl lithium; by reduction of these products by the Clemmensen method the following hydrocarbons were obtained: 1-ethyl adamantane, 1-n-propyl adamantane, and 1-n-butyl adamantane.

The following carbinols were formed as by-products of the reaction of adamantane carboxylic-1 acid with alkyl lithium: 2-(1-adamantyl)-propanol-2, 3-(1-adamantyl)-pentanol-3, and 4-(1-adamantyl)-heptanol-4. Their reduction by the Clemmensen method will yield 2-(1-adamantyl)-propene and 2-(1-adamantyl)-propane, 3-(1-adamantyl)-pentene-2 and 3-(1-adamantyl)-pentane, 4-(1-adamantyl)-heptene-3, and 4-(1-adamantyl)-heptane. These hydrocarbons were identified with the help of gas-liquid chromatography, by comparison with synthesized standard preparations. Besides this, 3-(1-adamantyl)-pentene-2 and 3-(1-adamantyl)-pentane were separated by means of preparative gas-liquid chromatography.

As compared to procedures published earlier, this method gives higher yields of the reaction products.

In the work we also synthesized the previously undescribed 1-tret-butyl adamantane, formed together with 2-(1-adamantyl)-propene.

References

1. Tegner, C. Acta Chem. Scand., 6, 782, 1952.
2. Gilman, H. and P. R. van Ness. J. Amer. Chem. Soc., 55, 1258, 1933.
3. Landa, S. and S. Hála. Coll. Czechoslov. Chem. Communs., 24, 93, 1959.
4. Spengler, G. and F. Wunsch. Erdöl und Kohle, 9, 702, 1962.
5. Gilman, H., J. A. Beel, et al. J. Amer. Chem. Soc., 71, 1499, 1949.
6. Landa, S. and S. Hála. Coll. Czechoslov. Chem. Communs., 25, 2692, 1960.

7. Hoek, W., J. Strating, and H. Wynberg. Recueil. Trav. Chim., 85, 1045, 1966.

8. Hála, S. and S. Landa. Unpublished data.

9. Grob, C. A., W. Schwarz, and H. P. Fischer. Helv. Chim. Acta, 47, 1385, 1964.

Prague Institute of Chemical Technology

Received
6 December 1967