


SAM-TR-70-15

6265020A
ADD705979

**INFLUENCE OF TEMPERATURE ON THE EFFECTS OF
ENTEROTOXIN AND ENDOTOXIN IN MICE**

WILLIAM G. GLENN, Ph. D.



**USAF School of Aerospace Medicine
Aerospace Medical Division (AFSC)
Brooks Air Force Base, Texas**

March 1970

**This document has been approved for public release and sale;
its distribution is unlimited.**

Qualified requesters may obtain copies of this report from DDC. Orders will be expedited if placed through the librarian or other person designated to request documents from DDC.

When U. S. Government drawings, specifications, or other data are used for any purpose other than a definitely related Government procurement operation, the Government thereby incurs no responsibility nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise, as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use, or sell any patented invention that may in any way be related thereto.

ACCESSION FOR		
CFSTI	WHITE SECTION	<input checked="" type="checkbox"/>
DDC	DIFF SECTION	<input type="checkbox"/>
UN ANNOUNCED		<input type="checkbox"/>
JUSTIFICATION		
BY		
DISTRIBUTION AVAILABILITY CODES		
DIST.	AVAIL.	DDC OR SPECIAL
1		

INFLUENCE OF TEMPERATURE ON THE EFFECTS OF ENTEROTOXIN AND ENDOTOXIN IN MICE

WILLIAM G. GLENN, Ph. D.

This document has been approved for public release and sale;
its distribution is unlimited.

FOREWORD

This research was accomplished in the Microbiology-Immunology Branch, under task No. 775309, between January and May 1969. The paper was submitted for publication on 12 January 1970.

The animals involved in this study were maintained in accordance with the "Guide for Laboratory Animal Facilities and Care" as published by the National Academy of Sciences-National Research Council.

For this research, the B-type enterotoxin (SEB) was supplied by Dr. E. J. Schantz, Ft. Detrick, Frederick, Md.; and the lipopolysaccharide endotoxin (LPS) was secured from the Difco Laboratories, Detroit, Mich.

The author expresses his appreciation to Alton J. Rahe, Biometrics Division, for the statistical analyses.

This report has been reviewed and is approved.


JOSEPH M. QUASHNOCK
Colonel, USAF, MC
Commander

ABSTRACT

For optimum health, the host must control the proliferation of endogenous bacteria and neutralize harmful toxins. Staphylococcal enterotoxin B (SEB) produced by gram-positive bacteria and lipopolysaccharide endotoxin (LPS) produced by certain gram-negative bacteria can exert synergistic effects when injected into homothermic animals (mice). In both acclimatized and nonacclimatized animals, exposure to temperature extremes of 4° C. or 35° C.—except in one instance—either hastens or increases death in the presence of SEB and LPS when compared to room temperature (24° C.). This research determines experimentally that ambient temperature is a meaningful variable in the resistance or susceptibility of a mammal to certain common bacterial toxins.

INFLUENCE OF TEMPERATURE ON THE EFFECTS OF ENTEROTOXIN AND ENDOTOXIN IN MICE

I. INTRODUCTION

Staphylococcal enterotoxin (SE) and lipopolysaccharide endotoxin (LPS), separately and in synergistic combination, are the subject of this research. In man, SE is a common cause of food poisoning (1); and LPS, produced by certain gram-negative bacteria, causes fever. The bacteria that produce these toxins are common genera which may inhabit man and other animals. In these hosts, effective concentrations of the toxins may occur simultaneously.

No observable effect occurs when SE is injected into mice, and LPS is only slightly incapacitating; but the combination, when given in sequence at certain concentrations, is highly lethal.

II. MATERIALS AND METHODS

Toxins

The SE was a lyophilized extract of *Staphylococcus* spp. Fractionation of this extract resulted in a B-type enterotoxin, generally referred to as SEB (2). The LPS endotoxin was a commercial preparation of lipopolysaccharide obtained from *Salmonella enteritidis*. The animals were administered, intraperitoneally (I.P.): 12 μ g. of SEB in 0.1 ml. of saline; and, approximately 4 hours later, 150 μ g. of LPS in 0.1 ml. of saline. To minimize circadian variability, the SEB injections were given between 8:00 and 9:00 a.m. and the LPS injections between 12:00 and 1:00 p.m. (3).

Animals and housing

Male mice, of the Swiss-Webster strain, were used in this research. They ranged in

weight from 20 to 26 gm. The animals were housed in groups of two in plastic cages (28 cm. x 18 cm. x 13 cm.) having perforated metal lids and pellet troughs. Small groups of mice were preferred in order to increase the liberation of body heat at the higher temperature and to decrease any effects of huddling at the lower temperature. Food and water were provided ad libitum. Sawdust bedding was kept to a minimum to limit the burrowing tendency expected at the lower temperature.

Environmental conditions

Three temperatures were selected for study: 35° C., 24° C., and 4° C. The 24° C. environment was the normal laboratory condition and served as a control. (Acclimatization to 24° C., when stated in figure legends, signifies that the mice were not moved from any other temperature to 24° C.) The high temperature was maintained by placing the cages in an incubator room; 4° C. was the temperature of the cold room. Temperature variation was approximately $\pm 2^\circ$ C. The conditions are summarized in table I.

III. RESULTS

In figure 1 are shown the lethality curves for mice exposed to 35° C., 24° C., and 4° C. under acclimatized (48 hours) and nonacclimatized conditions. These mice were injected with saline and, 4 hours later, with LPS. Evidently the trauma of sudden ambient temperature change increases the susceptibility of nonacclimatized mice to LPS endotoxin (fig. 1: exp. 1 vs. exps. 2, 3) at both 4° C. and 35° C.

When the animals were acclimatized for 48 hours before being injected with saline and

TABLE I

Ambient conditions and injections used on mice to explore influence on effects of SEB and LPS

Exp. No.	Temp. (deg. C.)	Acclimatized	SEB	LPS	Remarks
1	24	No	No	Yes	Saline for SEB
2*	35				
3*	4				
4	4	48 hr.	No	Yes	
5	35				
6	24	No	Yes	Yes	Control
7	4				
8	35				
9	4	48 hr.	Yes	Yes	—
10	35				
11*	35	No	Yes	No	Saline for LPS
12*	4				
13*	35	No	No	No	Saline-temp. control
14*	4				

*Only 10 mice per experiment. Each of the other experiments had 20 mice.

later with LPS, only the 35° C. temperature (exp. 5) increased their susceptibility to a statistically significant probability level.

Experiments 11 and 12, in which SEB is used before saline, were conducted as additional controls. (These experiments are not included in figure 2 because no deaths occurred at 4° C. or 35° C.)

As shown in figure 2, both the temperature-acclimatized (exp. 9, 10) and the nonacclimatized (exp. 7, 8) mice die at a faster rate at 4° C. (exp. 7, 9) and 35° C. (exp. 8, 10) than at 24° C. (exp. 6). Moreover, in the nonacclimatized group, at 35° C. (exp. 8), the increased deaths at 3.5 hours were statistically

significant compared to those at 24° C. (exp. 6) and at 4° C. (exp. 7).

IV. DISCUSSION

The mechanism and effects of endotoxin from gram-negative bacteria have stimulated considerable research for two decades (4-6). In addition to vascular effects, other alterations (e.g., pyrexia, leukopenia, leukocytosis, hyper- and hypoglycemia, changed resistance to infections, and tumor necrotizing capacity) have been associated with LPS (7). In the extreme, these symptoms lead to death of the host.

According to one report at least, oxygen tensions and cell anoxia resulting from metabolic events are responsible for endotoxin




FIGURE 1

Total deaths vs. hours for mice after last injections of 0.1 ml. of saline (I.P.) and, 4 hours later, of 150 μ g./0.1 ml. of LPS (I.P.). [Before injection, the acclimatized groups were held 48 hours at the temperatures shown. Statistically significant (SS) differences and probability levels are indicated. Experiments 2 and 3 had 10 animals per group. The number of deaths was doubled to facilitate graphing.]

shock. Glycorticoids from the adrenal cortex seem to be effective in protecting some animals from endotoxin (8). Some evidence indicates that adrenal hypertrophy occurs in hamsters kept at 5° C. (9), and it is reasonable to suspect that the same occurs in mice. Such hypertrophy would increase the corticosteroids and




FIGURE 2

Total deaths vs. hours for mice after last injections of 12 μ g./0.1 ml. of SEB (I.P.) and, 4 hours later, of 150 μ g./0.1 ml. of LPS (I.P.). [Before injection, the acclimatized groups were held 48 hours at the temperatures shown. Statistically significant (SS) differences and probability levels are indicated.]

thus increase resistance. As for saline and LPS only, ambient temperature affects the rate of death and eventual overall mortality in both acclimatized and nonacclimatized animals, except in one instance (fig. 1): acclimatized mice seem to combat toxin effects better at 4° C. than at 35° C.

REFERENCES

1. Dack, G. M. *Staphylococcus enterotoxin: A review.* Jap. J. Med. Sci. Biol. 16:1-12 (1963).
2. Wagman, J., R. C. Edwards, and E. J. Schantz. Molecular size, homogeneity, and hydrodynamic properties of purified staphylococcal enterotoxin B. *Biochemistry (Wash.)* 4:1017-1023 (1965).
3. Becker, R. E., and W. G. Glenn. Differential resistance of mice at various time levels to the lethality of rattlesnake venom (*Crotalus atrox*). *Texas Rep. Biol. Med.* 25:360-364 (1967).
4. Thomas, L. Physiologic and pathologic alterations produced by the endotoxins of Gram negative bacteria. *Arch. Int. Med. (Chicago)* 101: 452-461 (1958).
5. Sugiyama, H., et al. Enhancement of bacterial endotoxin lethality by staphylococcal enterotoxin. *J. Infect. Dis.* 114:111-118 (1964).
6. Zweifach, B. W., B. Benacerraf, and L. Thomas. The relationship between vascular manifestations of shock. *J. Exp. Med.* 106:403-414 (1957).

7. Gilbert, R. P. Mechanisms of the hemodynamic effects of endotoxin. *Physiol. Rev.* 40:245-279 (1960).
8. Berry, J. L. *In* Discussion. Influence of cold on host-parasite interactions, part III, p. 444. USAF Arctic Aeromedical Laboratory, Ft. Wainwright, Alaska, 1963.
9. Teodoru, C. V., and G. Schwartzman. Endocrine factors in pathogenesis of experimental poliomyelitis in hamsters. Role of inoculatory and environmental stress. *Proc. Soc. Exp. Biol. Med.* 91:181-187 (1956).

Unclassified

Security Classification

DOCUMENT CONTROL DATA - R & D		
<i>(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)</i>		
1. ORIGINATING ACTIVITY (Corporate author) USAF School of Aerospace Medicine Aerospace Medical Division (AFSC) Brooks Air Force Base, Texas		2a. REPORT SECURITY CLASSIFICATION Unclassified
		2b. GROUP
3. REPORT TITLE INFLUENCE OF TEMPERATURE ON THE EFFECTS OF ENTEROTOXIN AND ENDOTOXIN IN MICE		
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Progress report January - May 1969		
5. AUTHOR(S) (First name, middle initial, last name) William G. Glenn		
6. REPORT DATE March 1970	7a. TOTAL NO. OF PAGES 4	7b. NO. OF REFS 9
8a. CONTRACT OR GRANT NO.	9a. ORIGINATOR'S REPORT NUMBER(S) SAM-TR-70-15	
b. PROJECT NO. 7753		
c. Task No. 775309	9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d.		
10. DISTRIBUTION STATEMENT This document has been approved for public release and sale; its distribution is unlimited.		
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY USAF School of Aerospace Medicine Aerospace Medical Division (AFSC) Brooks Air Force Base, Texas
13. ABSTRACT For optimum health, the host must control the proliferation of endogenous bacteria and neutralize harmful toxins. Staphylococcal enterotoxin B (SEB) produced by gram-positive bacteria and lipopolysaccharide endotoxin (LPS) produced by certain gram-negative bacteria can exert synergistic effects when injected into homothermic animals (mice). In both acclimatized and non- acclimatized animals, exposure to temperature extremes of 4° C. or 35° C.-- except in one instance--either hastens or increases death in the presence of SEB and LPS when compared to room temperature (24° C.). This research determines experimentally that ambient temperature is a meaningful variable in the resistance or susceptibility of a mammal to certain common bacterial toxins.		

DD FORM 1 NOV 65 1473

Unclassified

Security Classification

Unclassified

Security Classification

14 KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Immunology Microbiology Toxicology Endotoxin, temperature Bacterial toxin, temperature Temperature influence on endotoxin and enterotoxin						

Unclassified

Security Classification