

AD734829

AD

**EDGEWOOD ARSENAL
TECHNICAL REPORT**

EATR 4588

**SYNTHESIS OF
SOME 1-SUBSTITUTED-4-FORMYLPYRIDINIUM
OXIMES AND OXIMATES**

by


**David W. Reger
Edward J. Poziomek**

December 1971



Reproduced by
**NATIONAL TECHNICAL
INFORMATION SERVICE**
Springfield, Va. 22151

**DEPARTMENT OF THE ARMY
EDGEWOOD ARSENAL
Chemical Laboratory
Edgewood Arsenal, Maryland 21010**



UNCLASSIFIED

Security Classification

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) CO, Edgewood Arsenal ATTN: SMUEA-CI-PD Edgewood Arsenal, Maryland 21010		2a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED	
		2b. GROUP NA	
3. REPORT TITLE SYNTHESIS OF SOME 1-SUBSTITUTED-4-FORMYLPYRIDINIUM OXIMES AND OXIMATES			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) This work was performed in February 1970.			
5. AUTHOR(S) (First name, middle initial, last name) David W. Reger Edward J. Poziomek			
6. REPORT DATE December 1971		7a. TOTAL NO. OF PAGES 15	7b. NO. OF REFS 3
8a. CONTRACT OR GRANT NO.		8b. ORIGINATOR'S REPORT NUMBER(S) EATR 4588	
8c. PROJECT NO.			
8d. Task No. 1B662710AD2901		8e. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
10. DISTRIBUTION STATEMENT Approved for public release; distribution unlimited.			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY NA	
13. ABSTRACT In connection with a broad study to find new reagents for applications in chemical detection, a number of new pyridinium oximes and oximates were synthesized. 1-Substituted-4-formylpyridinium chloride oximes were synthesized by alkylation of isonicotinaldehyde oxime. Corresponding iodide salts were obtained by treating the chlorides with an excess of methyl iodide. Oximate internal salts were prepared by the reaction of the oxime halides with silver oxide. All the oximates were found to be stable at 100°C for at least 8 hours.			
14. KEYWORDS Pyridinium compounds Oximes Oximates Defense systems Detection Army research			

DD FORM 1473

REPLACES DD FORM 1473, 1 JAN 64, WHICH IS OBSOLETE FOR ARMY USE.

15

UNCLASSIFIED

Security Classification

TEST	WHITE SECTION	☐
UC	DIFF SECTION	☐
		☐
DISTRIBUTION/AVAILABILITY CODES		
DIST.	AVAIL.	SPECIAL
A		

Distribution Statement

Approved for public release; distribution unlimited.

Disclaimer

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

Disposition

Destroy this report when no longer needed. Do not return it to the originator.

EDGEWOOD ARSENAL TECHNICAL REPORT

EATR 4588

SYNTHESIS OF SOME 1-SUBSTITUTED-4-FORMYLPYRIDINIUM
OXIMES AND OXIMATES

by

David W. Reger
Edward J. Poziomek

Defense Research Branch

December 1971

Approved for public release; distribution unlimited.

Task 1B662710AD2901

DEPARTMENT OF THE ARMY
EDGEWOOD ARSENAL
Chemical Laboratory
Edgewood Arsenal, Maryland 21010

FOREWORD

The work described in this report was authorized under Task 1B662710AD2901, Chemical Detection and Identification Technology, Detection and Identification Concepts. The work was performed in February 1970. The results are recorded in notebook 8328.

Reproduction of this document in whole or in part is prohibited except with permission of the Commanding Officer, Edgewood Arsenal, ATTN: SMUEA-TS-R, Edgewood Arsenal, Maryland 21010; however, DDC and The National Technical Information Service are authorized to reproduce the document for United States Government purposes.

Acknowledgments

Elemental analyses were performed by the Analytical Chemistry Branch, Chemical Laboratory, Edgewood Arsenal, Maryland 21010.

DIGEST

In connection with a broad study to find new reagents for applications in chemical detection, a number of new pyridinium oximes and oximates were synthesized. 1-Substituted-4-formylpyridinium chloride oximes were synthesized by alkylation of isonicotinaldehyde oxime. Corresponding iodide salts were obtained by treating the chlorides with an excess of methyl iodide. Oximate internal salts were prepared by the reaction of the oxime halides with silver oxide. All the oximates were found to be stable at 100°C for at least 8 hours.

CONTENTS

	Page
I INTRODUCTION	7
II. MATERIALS AND METHODS	7
A. Chloride Oximes	7
B. Iodide Oximes	7
C. Oximate Inner Salts	7
III. RESULTS	7
DISTRIBUTION LIST	11

Preceding page blank

SYNTHESIS OF SOME 1-SUBSTITUTED-4-FORMYLPYRIDINIUM OXIMES AND OXIMATES

I. INTRODUCTION.

As part of a broad study on the use of pyridinium derivatives in chemical detection, a series of 1-substituted-4-formylpyridinium chloride (and iodide) oximes and 1-substituted-4-formylpyridinium oximates were synthesized and characterized. Analytical data, colors, and melting points for the chloride oximes, the iodide oximes, and the oximates are given in tables I, II, and III. No attempt was made to optimize yields, but they were generally well over 50%.

II. MATERIALS AND METHODS.

A. Chloride Oximes.

The 1-substituted-4-formylpyridinium chloride oximes were prepared by dissolving equimolar quantities of isonicotinaldehyde oxime (mp 132° to 133°C) and the appropriate benzyl chloride in 75 ml of acetone and heating the solutions overnight at 80°C. It was found convenient to use 200-ml heavy walled polymer bottles, which were capped before placing them in the 80° oven. Colorless solids formed during the heating process. The bottles were allowed to cool to room temperature; the reaction mixtures were filtered. One recrystallization was performed from methanol-ether. CAUTION! Because of the lachrymal properties of the alkylating agents, all procedures were carried out in a fume hood. 1-(*n*-Dodecyl)-4-formylpyridinium bromide oxime was prepared in a similar fashion from isonicotinaldehyde oxime and *n*-dodecyl bromide.

B. Iodide Oximes.

The 1-substituted-4-formylpyridinium iodide oximes were prepared by adding an excess of methyl iodide to a methanolic solution of the appropriate chloride oxime and allowing the solution to stand for at least 24 hours. The solutions were protected from light during the reaction period. Addition of ether effected the precipitation of the iodide oximes. One recrystallization was performed from methanol-ether.

C. Oximate Inner Salts.

The 1-substituted-4-formylpyridinium oximate inner salts were prepared by treating a methanolic solution of the appropriate halide oxime with excess silver oxide. (To ensure dryness of the reaction medium, 3 ml of triethylorthoformate were added prior to adding the silver oxide.) The resulting mixture was stirred vigorously for 5 minutes and then filtered. Slow addition of ethyl ether to the filtrate generally caused the oximate to crystallize. When immediate crystallization did not occur (as with the dodecyl and *p*-methoxybenzyl derivatives), the solution was evaporated to dryness. Ethyl ether then was added to the residue, and the mixture was allowed to stand in a refrigerator until solidification occurred. The mixture was filtered, and the product was dried over calcium sulfate in an Abderhalden apparatus at 100°C under vacuum for at least 8 hours.

III. RESULTS.

In contrast to the reported instability of 1-methyl-4-formylpyridinium oximate,* the oximates reported here were stable under the drying conditions of 100°C for at least 8 hours. Also, none of the oximates showed any signs of decomposition after several months at room temperature. Pyridinium oximates are of interest because of their internal charge-transfer properties.

*Engelhard, N., and Werth, B. *Tetrahedron Letters* (10), 661 (1963).

Table 1. 1-Substituted-4-Formylpyridinium Chloride Oximes





Table II. 1-Substituted 4-Formylpyridinium Iodide Oximes




R	mp ^a °C	Color	Calcd						Found					
			C	H	Cl	I	N	O	C	H	Cl	I	N	O
<i>n</i> -Dodecyl ^b	94-96	Yellow	51.7	7.5			6.7	3.8	51.6	7.3			6.9	3.7
Benzyl	113-117	Yellow	45.9	3.9		37.3		4.7	45.6	4.1		37.0		
<i>p</i> -Chlorobenzyl	198-199	Yellow	41.6	3.2	9.5	33.9			41.4	3.0	9.6	33.8		
<i>o</i> -Chlorobenzyl	165-167	Yellow	41.6	3.2			7.5	4.3	41.8	3.3			7.5	4.6
2,4-Dichlorobenzyl	207-208	Yellow	38.2	2.7		31.0	6.9	3.9	38.3	2.8		31.1	6.8	3.8
3,4-Dichlorobenzyl	211-213	Yellow	38.2	2.7		31.0	6.9	3.9	38.3	3.0			7.0	4.2
<i>p</i> -Methoxybenzyl	176-178	Yellow	45.4	4.1			7.6	8.7	45.7	4.3			7.6	8.9
<i>p</i> -Nitrobenzyl	184-187	Yellow	40.5	3.1			10.9	12.5	40.3	3.1			10.7	
<i>m</i> -Nitrobenzyl	192-193	Yellow	40.5	3.1		33.0			40.3	3.4		32.9		12.3

^aDecomposition points except for the *n*-dodecyl derivative. A Thomas Hoover capillary melting point apparatus was used.

^bPreviously reported in Cohen, W., and Erlanger, B. F. J. Amer. Chem. Soc. 82, 3928 (1960).

Table III. 1-Substituted-4-Formylpyridinium Oximates



R	mp ^a	Color	Calcd					Found				
			C	H	Cl	N	O	C	H	Cl	N	O
	°C											
<i>n</i> -Dodecyl	103-105	Black	74.4	10.4	9.7	5.5		74.1	10.5	9.8		
Benzyl	121-123	Purple-black	73.6	5.7	13.2			73.3	5.8	13.3		14.3
<i>p</i> -Chlorobenzyl	135-138	Brown	63.3	4.5			14.4	62.9	4.5			14.1
<i>o</i> -Chlorobenzyl	128-131	Black	63.3	4.5			14.4	62.9	4.6			24.9
2,4-Dichlorobenzyl	132-133	Black-brown	55.5	3.6			25.2	55.3	3.8			25.3
3,4-Dichlorobenzyl	152-154	Dark purple	55.5	3.6			25.3	55.6	3.6			14.6
<i>p</i> -Methoxybenzyl	105-109	Black	69.4	5.8			14.5	69.1	5.6			
<i>p</i> -Nitrobenzyl ^b	113-134	Purple-black	59.4	4.3				59.5	4.4			
<i>m</i> -Nitrobenzyl	136-138	D- ⁺ , brown	60.6	4.3	16.3			60.5	4.3	16.2		

^aDecomposition points. A Thomas Hoover capillary melting point apparatus was used.^bMonohydrate.