


AD 745191



TECHNICAL / SCIENTIFIC REPORT


17-72

CANINE LEPTOSPIROSIS IN CAIRO

by


R.R. Maronpot, I.S. Barsoun and E. Ezzat

NATIONAL TECHNICAL
INFORMATION SERVICE



U. S. NAVAL MEDICAL RESEARCH UNIT No.3
(CAIRO, THE ARAB REPUBLIC OF EGYPT)
FPO NEW YORK 09527

DISCLAIMER NOTICE



THIS DOCUMENT IS BEST
QUALITY AVAILABLE. THE COPY
FURNISHED TO DTIC CONTAINED
A SIGNIFICANT NUMBER OF
PAGES WHICH DO NOT
REPRODUCE LEGIBLY.

UNCLASSIFIED

Security Classification

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) U.S. Naval Medical Research Unit No. 3 FPO New York 09527		2a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED	
		2b. GROUP	
3. REPORT TITLE CANINE LEPTOSPIROSIS IN CAIRO			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) M Interim Report			
5. AUTHOR(s) (First name, middle initial, last name) R.R. Maronpot, I.S. Barsoum and E. Ezzat			
6. REPORT DATE May 1970		7a. TOTAL NO. OF PAGES 3	7b. NO. OF REFS 22
8a. CONTRACT OR GRANT NO.		9a. ORIGINATOR'S REPORT NUMBER (S) NAIRU-3-TR.17-72	
b. PROJECT NO. MF12-524.009-3024		9b. OTHER REPORT NO (S) (Any other numbers that may be assigned this report)	
c.			
d.			
10. DISTRIBUTION STATEMENT Distribution of this report is unlimited			
11. SUPPLEMENTARY NOTES Published in J. Infect. Dis 123(5):548-550, 1970		12. SPONSORING/MONITORING ACTIVITY Bureau of Medicine and Surgery Department of the Navy Washington, D.C. 20390	
13. ABSTRACT Sera of 170 dogs in Cairo were tested for leptospiral antibodies by the microscopic agglutination test. Forty-one sera had detectable leptospiral agglutinins, most reacted against <u>Leptospira</u> serotype <u>canicola</u>. Subacute interstitial nephritic lesions were present in renal tissue from 27 seropositive and two seronegative dogs. <u>Leptospira</u> serotype <u>canicola</u> was isolated in cultures from urine of two dogs.			

14 KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
<u>Leptospira serotype canicola</u> Leptospirosis Canine Leptospiral antibodies Interstitial nephritis Egypt						

Reprinted for private circulation from
THE JOURNAL OF INFECTIOUS DISEASES • VOL. 123, NO. 5 • MAY 1971
© 1971 by the University of Chicago. All rights reserved.

Canine Leptospirosis in Cairo

R. R. Maronpot, I. S. Barsoum and E. Ezzat

From the Veterinary Medicine Department,
United States Naval Medical Research Unit No. 3,
and the Veterinary Department,
Ministry of Agriculture,
Cairo, United Arab Republic

The worldwide distribution of canine leptospirosis has been documented by isolation of *Leptospira*¹ and serologic surveys [1-4]. Frequencies of leptospiral antibodies in canine populations range from 3 to 65% [1-4]. Infections of dogs with *Leptospira* serotype *canicola* and *Leptospira* serotype *icterohemorrhagiae* produce serum agglutinins which may be detectable for years [5].

Naturally occurring canine leptospirosis has been associated with interstitial nephritis [6, 7], but this relationship is not always easily demonstrated experimentally [8-11]. Recently, Anderson [12] and Taylor et al. [13] have produced subacute interstitial nephritis following experimental infection of dogs with *canicola*. However, canine leptospiral infection has not been proved to cause chronic interstitial nephritis [14, 15].

Canine leptospirosis in Egypt, U.A.R., has been poorly studied. McGuire and Myers [16] reported that five of 30 sera from Cairo dogs, when tested by complement fixation (CF), had antibody titers of 1:8 to *Leptospira* but were negative by the microscopic agglutination test.

More recently, Hamed [17] documented that prevalence of leptospiral seropositivity in 98 dogs in Cairo was 15%. Leptospiral agglutinating antibodies have been reported in man and in non-canine domesticated and wild mammals in Egypt [16-19].

We undertook this study to assess the serologic status of dogs in Cairo with respect to *Leptospira* and to determine whether there is any consistent relationship between their serologic status and interstitial nephritic lesions.

Methods

Jugular blood was obtained from 170 unselected stray dogs from different areas of Cairo; sera were harvested and stored at -20 C until used for leptospiral serology. Blood from 68 of these dogs was cultured by aseptic addition of two drops of whole blood to two tubes each of Fletcher's and Ellinghausen's media. All dogs were killed and necropsied. Urine was aseptically obtained from the urinary bladders of the same 68 dogs immediately after death and was inoculated with and without dilution into three tubes each of Fletcher's and Ellinghausen's media. Kidney and liver specimens were fixed in 10% formalin buffered at neutral pH and processed routinely. Sections were stained with hematoxylin and eosin; Warthin-Starry stains were made on kidney sections.

All sera were tested by the microscopic agglutination (MA) test [20, 21] with use of the following serotypes of *Leptospira*: *canicola*, *austumnalis*, *sentot*, *djasimana*, *bataviae*, *australis*, *javanica*, *grippotyphosa*, *icterohemorrhagiae*, *wolffi*, *hebdomadis*, *pomona*, *tarassovi*, *pyrogenes*, *ballum*, *butembo*, and *cynopteri*. The highest final twofold dilution of serum with at least 50% agglutination was the endpoint.

Significant interstitial nephritis is defined as active diffuse inflammation producing histologic

Received for publication May 26, 1970 and in revised form December 3, 1970.

The opinions and assertions in this report are those of the authors and do not necessarily reflect the official views of the U. S. Department of the Navy, the naval services at large, or the Egyptian Ministry of Agriculture. This research was supported by Bureau of Medicine and Surgery work unit no. MF12.524.009-3024B.

The experiments reported herein were conducted according to the principles set forth in "Guide for Laboratory Animal Facilities and Care" prepared by the National Academy of Science, National Research Council.

Please address requests for reprints to Medical Editor, NAMRU-3, FPO, New York 09527.

¹ Galton, M. M. 1966. Leptospiral Serotype Distribution Lists. Communicable Disease Center Zoonoses Surveillance, U. S. Department of Health, Education, and Welfare, Public Health Service, Atlanta, Georgia, 130 p.

[illegible]

significant interstitial nephritis for this study were established to avoid the probability of associating cases of spontaneous or noninfectious nephritis with leptospiral seropositivity.

This study was intended to document the prevalence of leptospiral seropositivity and associated interstitial nephritis in dogs in Cairo, not to prove or disprove a causal relationship between naturally occurring canine leptospirosis and interstitial nephritis. The results do document a higher prevalence of severe interstitial nephritis associated with leptospiral seropositivity than with seronegativity. Although specimens that are positive by the Warthin-Starry stain and isolation of leptospires from seropositive dogs with severe nephritis are not absolute proof of a causal relationship between leptospiral infection and canine interstitial nephritis, such a relationship is at least suggested. Other factors may causally lead to interstitial nephritis.

Summary

Sera of 170 dogs in Cairo were tested for leptospiral antibodies by the microscopic agglutination test. Forty-one sera had detectable leptospiral agglutinins, most reacted against *Leptospira* serotype *canicola*. Subacute interstitial nephritic lesions were present in renal tissue from 27 seropositive and two seronegative dogs. *Leptospira* serotype *canicola* was isolated in cultures from urine of two dogs.

References

1. Rosenberg, B. L. 1951. Canicola fever. Review, with report of two new cases. *Amer. J. Med.* 11:75-91.
2. Byrne, R. J. 1955. Canine leptospirosis and public health. *Public Health Rep.* 70:1229-1236.
3. Alexander, A. D., C. A. Gleiser, P. Malnati, and H. Yoder. 1957. Observations on the prevalence of leptospirosis in canine populations of the United States. *Amer. J. Hyg.* 65:43-56.
4. White, F. H., H. F. Stoliker, and M. M. Galton. 1961. Detection of leptospires in naturally infected dogs, using fluorescein-labeled antibody. *Amer. J. Vet. Res.* 22:650-654.
5. McIntyre, W. I. M., and H. F. Seiler. 1953. Epidemiology of canicola fever. *J. Hyg. (Camb.)* 51:330-339.
6. Bloom, F. 1941. The histopathology of canine leptospirosis. *Cornell Vet.* 31:266-288.
7. Ohshima, K., and S. Miura. 1966. Pathological studies of canine leptospirosis in naturally occurring cases. *J. Fac. Agric. Iwate Univ.* 8:15-49.
8. Monlux, W. S. 1948. Leptospirosis. IV. The pathology of canine leptospirosis. *Cornell Vet.* 38:199-208.
9. Low, D. G., C. W. Hiatt, C. A. Gleiser and E. N. Bergman. 1956. Experimental canine leptospirosis. I. *Leptospira icterohemorrhagiae* infections in immature dogs. *J. Infect. Dis.* 98:249-259.
10. Gleiser, C. A. 1957. Experimental canine leptospirosis. III. Histopathologic changes. *J. Infect. Dis.* 100:249-256.
11. Menges, R. W., M. M. Galton and R. T. Habermann. 1960. Culture and serologic studies on four dogs inoculated with two leptospiral serotypes, *Leptospira pomona* and *Leptospira canicola*. *Amer. J. Vet. Res.* 21:371-376.
12. Anderson, L. J. 1967. Experimental reproduction of canine interstitial nephritis. *J. Comp. Path.* 77:413-418.
13. Taylor, P. L., L. E. Hanson and J. Simon. 1970. Serologic, pathologic, and immunologic features of experimentally induced leptospiral nephritis in dogs. *Amer. J. Vet. Res.* 31:1033-1049.
14. Low, D. G. 1964. Current status of canine leptospirosis. *Allied Vet. (Jan-Feb.)*:3-11.
15. Low, D. G., G. W. Mather, D. R. Finco and N. V. Anderson. 1967. Long-term studies of renal function in canine leptospirosis. *Amer. J. Vet. Res.* 28:731-739.
16. McGuire, C. D., and D. M. Myers. 1957. Leptospirosis in Egypt. *Amer. J. Trop. Med.* 6:532-539.
17. Hamed, O. M. 1968. Leptospirosis in animals and man in U.A.R. Thesis. Cairo University.
18. Hamdy, A. H., W. J. Brownlow and J. D. Dedeaux. 1962. Leptospirosis in bovines and their human contacts in Egypt (U.A.R.) *Amer. J. Trop. Med.* 11:98-101.
19. Brownlow, W. J., and J. D. Dedeaux. 1964. Leptospirosis in animals of upper Egypt. *Amer. J. Trop. Med.* 13:311-318.
20. Schüffner, W., and A. Mochtar. 1927. Versuche zur Aufteilung von Leptospirenstämmen, mit einleitenden Bemerkungen über den Verlauf von Agglutination und Lysis. *Zbl. Bakt. Orig.* 101:405-413.
21. Gochenour, W. S., R. H. Yager, P. W. Wetmore and J. A. Hightower. 1953. Laboratory diagnosis of leptospirosis. *Amer. J. Public Health.* 43:405-410.
22. Turner, I. H. 1968. Leptospirosis. II. Serology. *Trans. Roy. Soc. Trop. Med. Hyg.* 62:880-899.