

4

745284

**AD** 

**SR72-3** 

FRR

# ALTERATION OF RADIATION EFFECTS BY 2-(n-DECYLAMINO) ETHANETHIOSULFURIC ACID (WR1607) IN THE MONKEY

C. L. Turbyfill R. M. Roudon R. W. Young V. A. Kieffer



NATIONAL TECHNICAL INFORMATION SERVICE

ARMED FORCES RADIOBIOLOGY RESEARCH INSTITUTE Defense Nuclear Agency Bethesda, Maryland

Approved for public release; distribution unlimited



ļ

ł

- H. . . . .

1

ing a state



sa traaciati

\_\_\_\_\_ł\_\_

· . ,

<u>.</u>...,.....

eren (h. 1999) en el en filo Marin e restances company en el el

ŧ 

e. P 1.1

<u>\_</u> 1



. ಕೆಳ್ಳಾ

14

. . 1.1

Socionative contract on indexing annotative   OPECHNERIC Composite authous   Armed Forces Radiobiology Research Institute   Defense Nuclear Agency   Bethesda, Maryland 20014   ALTERATION OF RADIATION   2 -(n-DECYLAMINO) ETHANE   ACD (WR1607) IN THE   ALTERATION OF RADIATION   2 -(n-DECYLAMINO) ETHANE   ACD (WR1607) IN THE   AUTHOR(S) (First name, middle initial, lest name)   C. L. Turbyfill, R. M. Roudon, R. W. Young and   AUTHOR(S) (First name, middle initial, lest name)   C. L. Turbyfill, R. M. Roudon, R. W. Young and   AUTHOR(S) (First name, middle initial, lest name)   C. L. Turbyfill, R. M. Roudon, R. W. Young and   AUTHOR(S) (First name, middle initial, lest name)   C. L. Turbyfill, R. M. Roudon, R. W. Young and   AUTHOR(S) (First name, middle initial, lest name)   AUTHOR(S) (	Iden must be entered when the overall report is classified;   Iden must be entered when the overall report is classified;   Iden must be entered when the overall report is classified;   Iden must be entered when the overall report is classified;   Iden must be entered when the overall report is classified;   Iden must be entered when the second procession;   Iden V. A. Kieffer
ORIGINATING ACTIVITY (Corporate author)   Armed Forces Radiobiology Research Institute   Defense Nuclear Agency   Bethesda, Maryland 20014   AREPORT TITLE   ALTERATION OF RADIATION   2-(n-DECYLAMINO) ETHANE   ACID (WR1607) IN THE   OSSCRIPTIVE NOTES (Type of report and inclusive dates)   AUTHOR(S) (First name, middle initial, last name)   C. L. Turbyfill, R. M. Roudon, R. W. Young and   REPORT DATE   March 1972   CONTRACT OR GRANT NO.   B. PROJECT NO. NWER XAXM   A. Work Unit 04   OUSTAILOUTION STATEMENT   Approved for public release; distribution unlimited   TABLEMENTARY NOTES   12. SP   2-(n-decylamino) ethanethiosulfuric acid ( untrained monkeys (10 mg/kg) 30 minutes prior on behavior and physiology during the early trant to monitor aortic and venous pressures, heart to cally implanted in the untrained monkeys. The form a shock motivated visual discrimination ta	Image: All of the second se
Defense Nuclear Agency   Bethesda, Maryland 20014   ALTERATION OF RADIATION   2-(n-DECYLAMINO) ETHANE   ACID (WR1607) IN THE   OESCRIPTIVE NOTES (Type of report and inclusive dates)   AUTHOR(S) (Flist name, middle initial, last name)   C. L. Turbyfill, R. M. Roudon, R. W. Young and   MARCON 1972   * CONTRACT OR GRANT NO   *. PROJECT NO. NWER XAXM   *. PROJECT NO. NWER XAXM   *. Work Unit 04   * OISTAIEUTION STATEMENT   Approved for public release; distribution unlimits   * SAUSTRACT   2-(n-decylamino) ethanethiosulfuric acid ( untrained monkeys (10 mg/kg) 30 minutes prior on behavior and physiology during the early trant to monitor aortic and venous pressures, heart to cally implanted in the untrained monkeys. The form a shock motivated visual discrimination ta	ted
Bethesda, Maryland 20014   ALTERATION OF RADIATION   2-(n-DECYLAMINO) ETHANE   ACID (WR1607) IN THE   OESCRIPTIVE NOTES (Type of report and inclusive dates)   AUTHOR(S) (First name, middle initial, fast name)   C. L. Turbyfill, R. M. Roudon, R. W. Young and   March 1972   C. L. Turbyfill, R. M. Roudon, R. W. Young and   March 1972   C. CONTRACT ON GRANT NO   b. PROJECT NO. NWER XAXM   c. Task and Subtask C 903   obstateution statement   Approved for public release; distribution unlimited   'SUPPLEMENTARY NOTES   12. SP   S. ABSTRACT   2-(n-decylamino) ethanethiosulfuric acid ( untrained monkeys (10 mg/kg) 30 minutes prior on behavior and physiology during the early trant to monitor aortic and venous pressures, heart to cally implanted in the untrained monkeys. The form a shock motivated visual discrimination ta	N/A   DN EFFECTS BY   ETHIOSULFURIC   E MONKEY   and V. A. Kieffer   OTAL NO. OF PAGES   24   16   RIGIN STOR'S REPORT NUMBERIES   AFRRI SR72-3   THER REPORT NOIS) (Any other numbers that may be seeled   PONSORING MILLITARY ACTIVITY   Director   Defense Nuclear Agency   Washington, D. C. 20305   (WR 1607) was given to trained and   r to irradiation to investigate its effects
ALTERATION OF RADIATION 2-(n-DECYLAMINO) ETHANE ACID (WR1607) IN THE OESCRIPTIVE NOTES (Type of report and inclusive dates) AUTHOR(S) (First hame, middle initial, last name) C. L. Turbyfill, R. M. Roudon, R. W. Young and REPORT DATE March 1972 CONTRACT OF GRANT NO. PROJECT NO. NWER XAXM A PROJECT NO. NWER XAXM A Work Unit 04 OISTAIOUTION STATEMENT Approved for public release; distribution unlimited DISTAIOUTION STATEMENT 2-(n-decylamino) ethanethiosulfuric acid ( untrained monkeys (10 mg/kg) 30 minutes prior on behavior and physiology during the early tran- to monitor aortic and venous pressures, heart is cally implanted in the untrained monkeys. The form a shock motivated visual discrimination taken 2-(n-decylamino) ethaletion in the intrained monkeys. The form a shock motivated visual discrimination taken 2-(n-decylamino) ethaletion in the intrained monkeys. The form a shock motivated visual discrimination taken C. D. D. D. C. L. C.	DN EFFECTS BY ETHIOSULFURIC E MONKEY and V. A. Kieffer The NO. OF PAGES 24 BIGINATOR'S REPORT NUMBERIES AFRRI SR72-3 THER REPORT NOISI (Any other numbers that may be easily is report) ted PONSORING MILLITARY ACTIVITY Director Defense Nuclear Agency Washington, D. C. 20305 (WR 1607) Was given to trained and r to irradiation to investigate its effects
2-(n-DECYLAMINO) ETHANE   ACID (WR1607) IN THE   OESCRIPTIVE NOTES (Type of report and inclusive dates)   AUTHOR(S) (First name, middle initial, isst name)   C. L. Turbyfill, R. M. Roudon, R. W. Young and   March 1972   * CONTRACT OF GRANT NO.   * PROJECT NO. NWER XAXM   * Task and Subtask C 903   * Work Unit 04   * DISTRIBUTION STATEMENT   Approved for public release; distribution unlimited   * SUPPLEMENTARY NOTES   12. SP   * ABSTRACT   2-(n-decylamino) ethanethiosulfuric acid ( untrained monkeys (10 mg/kg) 30 minutes prior on behavior and physiology during the early trant to monitor aortic and venous pressures, heart is cally implanted in the untrained monkeys. The form a shock motivated visual discrimination task	ETHIOSULFURIC E MONKEY and V. A. Kieffer otal NO. OF PAGES 24 15. NO. OF PAGES 24 16 RIGINATOR'S REPORT NUMBER(S) AFRRI SR72-3 THER REPORT NOIS) (Any other numbers that may be seeled its mport) ted PONSORING MILLITARY ACTIVITY Director Defense Nuclear Agency Washington, D. C. 20305 (WR 1607) was given to trained and r to irradiation to investigate its effects
ACID (WR1607) IN THE OESCRIPTIVE NOTES (Type of report and inclusive dates) AUTHOR(3) (First name, middle initial, lest name) C. L. Turbyfill, R. M. Roudon, R. W. Young and REPORT DATE March 1972 CONTRACT OR GRANT NO. B. PROJECT NO. NWER XAXM A. Task and Subtask C 903 Work Unit 04 OISTALEUTION STATEMENT Approved for public release; distribution unlimited OISTALEUTION STATEMENT 2-(n-decylamino) ethanethiosulfuric acid ( untrained monkeys (10 mg/kg) 30 minutes prior on behavior and physiology during the early trant to monitor aortic and venous pressures, heart in cally implanted in the untrained monkeys. The form a shock motivated visual discrimination taken AUTHOR(3) (First name, middle initial, lest name) C. L. Turbyfill, R. M. Roudon, R. W. Young and AUTHOR(3) (First name, middle initial, lest name) C. L. Turbyfill, R. M. Roudon, R. W. Young and March 1972 (12. 50) (12. 50) (13. 50) (14. 50) (15. 50) (1	ted PONSORING MILLITARY ACTIVITY Director Defense Nuclear Agency Washington, D. C. 20305 (WR 1607) was given to trained and r to irradiation to investigate its effects
AUTHOR(3) (First name, middle initial, last name) AUTHOR(3) (First name, middle initial, last name) C. L. Turbyfill, R. M. Roudon, R. W. Young and March 1972 CONTRACT OF GRANT NO. PROJECT NO. NWER XAXM A PROJECT NO. NWER XAXM A Work Unit 04 OISTAIOUTION STATEMENT Approved for public release; distribution unlimited DISTAIOUTION STATEMENT Approved for public release; distribution unlimited DISTAIOUTION STATEMENT 2-(n-decylamino) ethanethiosulfuric acid ( untrained monkeys (10 mg/kg) 30 minutes prior on behavior and physiology during the early tran- to monitor aortic and venous pressures, heart is cally implanted in the untrained monkeys. The form a shock motivated visual discrimination taken a second a transmission and the second	ted PONSORING MILITARY ACTIVITY Director Defense Nuclear Agency Washington, D. C. 20305 (WR 1607) was given to trained and r to irradiation to investigate its effects
C. L. Turbyfill, R. M. Roudon, R. W. Young and MEPONT CATE (70. TO March 1972 C. CONTRACT OR GRANT NO. D. PROJECT NO. NWER XAXM A. Task and Subtask C 903 Work Unit 04 DISTALEUTION STATEMENT Approved for public release; distribution unlimited DISTALEUTION STATEMENT APPROVENTION STATEMENT A	ted PONSORING MILITARY ACTIVITY Director Defense Nuclear Agency Washington, D. C. 20305 (WR 1607) was given to trained and r to irradiation to investigate its effects
C. L. Turbyfill, R. M. Roudon, R. W. Young and REPORT DATE March 1972 CONTRACT OF GRANT NO. PROJECT NO. NWER XAXM A PROJECT NO. NWER XAXM A Task and Subtask C 903 Work Unit 04 OISTAILOUTION STATEMENT Approved for public release; distribution unlimited DISTAILOUTION STATEMENT Approved for public release; distribution	ted PONSORING MILITARY ACTIVITY Director Defense Nuclear Agency Washington, D. C. 20305 (WR 1607) was given to trained and r to irradiation to investigate its effects
ABEPORT DATE 10. TO   March 1972 10. TO   A. CONTRACT OR GRANT NO. 90. OR   B. PROJECT NO. NWER XAXM A   c. Task and Subtask C 903 90. OR   a. Work Unit 04 90. OR   O DISTAILOUTION STATEMENT 90. OR   Approved for public release; distribution unlimits 10. SPECTRACT   2 - (n-decylamino) ethanethiosulfuric acid ( untrained monkeys (10 mg/kg) 30 minutes prior on behavior and physiology during the early trant to monitor aortic and venous pressures, heart to cally implanted in the untrained monkeys. The form a shock motivated visual discrimination ta	ted PONSORING MILITARY ACTIVITY Director Defense Nuclear Agency Washington, D. C. 20305 (WR 1607) was given to trained and r to irradiation to investigate its effects
March 1972   A GATALLY OR GRANT NO.   B. PROJECT NO. NWER XAXM   A. Task and Subtask C 903   Work Unit 04   D OSTALEUTION STATEMENT   Approved for public release; distribution unlimited   D OSTALEUTION STATEMENT   A Proved for public release; distribution unlimited   D OSTALEUTION STATEMENT   A Proved for public release; distribution unlimited   D OSTALEUTION STATEMENT   A Proved for public release; distribution unlimited   D OSTALEUTION STATEMENT   A Proved for public release; distribution unlimited   D OSTALEUTION STATEMENT   2 - (n-decylamino) ethanethiosulfuric acid ( untrained monkeys (10 mg/kg) 30 minutes prior on behavior and physiology during the early trant to monitor aortic and venous pressures, heart to cally implanted in the untrained monkeys. The form a shock motivated visual discrimination tast	24 IG INIGINATOR'S REPORT NUMBERIES AFRRI SR72-3 ITHER REPORT NOIS) (Any other numbers that may be seeled is report) ted PONSORING MILITARY ACTIVITY Director Defense Nuclear Agency Washington, D. C. 20305 (WR 1607) Was given to trained and r to irradiation to investigate its effects
Approved for public release; distribution unlimite Approved for public release; distribution unlimite - OFFICE - CONTRACT - OFFICE - OFFIC	AFRRI SR72-3 THEN REPORT NOIS) (Any other numbers that may be seeled ted PONSORING MILLITARY ACTIVITY Director Defense Nuclear Agency Washington, D. C. 20305 (WR 1607) was given to trained and r to irradiation to investigate its effects
A PROJECT NO. NWER XAXM Task and Subtask C 903 Work Unit 04 DISTRIBUTION STATEMENT Approved for public release; distribution unlimite UPPLEMENTARY NOTES 2-(n-decylamino) ethanethiosulfuric acid ( untrained monkeys (10 mg/kg) 30 minutes prior on behavior and physiology during the early tran to monitor aortic and venous pressures, heart is cally implanted in the untrained monkeys. The form a shock motivated visual discrimination ta	AFRRI SR72-3 THEN AEPORT NOIS) (Any other numbers that may be easily ted PONSORING MILITARY ACTIVITY Director Defense Nuclear Agency Washington, D. C. 20305 (WR 1607) was given to trained and r to irradiation to investigate its effects
A A A A Task and Subtask C 903 Work Unit 04 Distance for public release; distribution unlimited Distance for public release; distrib	ted PONSORING MILITARY ACTIVITY Director Defense Nuclear Agency Washington, D. C. 20305 (WR 1607) was given to trained and r to irradiation to investigate its effects
Approved for public release; distribution unlimite Approved for public release; distribution unlimite Distribution statement 2-(n-decylamino) ethanethiosulfuric acid untrained monkeys (10 mg/kg) 30 minutes prior on behavior and physiology during the early tran to monitor aortic and venous pressures, heart is cally implanted in the untrained monkeys. The form a shock motivated visual discrimination ta	ted PONSORING MILITARY ACTIVITY Director Defense Nuclear Agency Washington, D. C. 20305 (WR 1607) was given to trained and r to irradiation to investigate its effects
A Work Unit 04 Approved for public release; distribution unlimite Approved for public release; distribution unlimite UNPPLEMENTARY NOTES 2-(n-decylamino) ethanethiosulfuric acid ( untrained monkeys (10 mg/kg) 30 minutes prior on behavior and physiology during the early tran- to monitor aortic and venous pressures, heart is cally implanted in the untrained monkeys. The form a shock motivated visual discrimination ta	ted Director Defense Nuclear Agency Washington, D. C. 20305 (WR 1607) was given to trained and r to irradiation to investigate its effects
Approved for public release; distribution unlimite USUPPLEMENTARY NOTES 2-(n-decylamino) ethanethiosulfuric acid ( untrained monkeys (10 mg/kg) 30 minutes prior on behavior and physiology during the early tran to monitor aortic and venous pressures, heart is cally implanted in the untrained monkeys. The form a shock motivated visual discrimination ta	ted PONSORING MILITARY ACTIVITY Director Defense Nuclear Agency Washington, D. C. 20305 (WR 1607) was given to trained and r to irradiation to investigate its effects
2-(n-decylamino) ethanethiosulfuric acid ( untrained monkeys (10 mg/kg) 30 minutes prior on behavior and physiology during the early tran to monitor aortic and venous pressures, heart cally implanted in the untrained monkeys. The form a shock motivated visual discrimination to	Washington, D. C. 20305 (WR1607) was given to trained and r to irradiation to investigate its effects
2-(n-decylamino) ethanethiosulfuric acid ( untrained monkeys (10 mg/kg) 30 minutes prior on behavior and physiology during the early tran to monitor aortic and venous pressures, heart to cally implanted in the untrained monkeys. The form a shock motivated visual discrimination to	Washington, D. C. 20305 (WR 1607) was given to trained and r to irradiation to investigate its effects
2-(n-decylamino) ethanethiosulfuric acid ( untrained monkeys (10 mg/kg) 30 minutes prior on behavior and physiology during the early tran to monitor aortic and venous pressures, heart cally implanted in the untrained monkeys. The form a shock motivated visual discrimination to	(WR 1607) was given to trained and r to irradiation to investigate its effects
2-(n-decylamino) ethanethiosulfuric acid ( untrained monkeys (10 mg/kg) 30 minutes prior on behavior and physiology during the early tran to monitor aortic and venous pressures, heart a cally implanted in the untrained monkeys. The form a shock motivated visual discrimination ta	(WR1607) was given to trained and r to irradiation to investigate its effects
a significant increase occurred in aortic pressurate. No significant changes in percent correct injection. The untrained injected animals were or 15,000 rads of mixed gamma-neutron radiat decrease in aortic pressure occurred from 2-4 from 1-2 minutes in the 15,000-rad group. A increased significantly in all injected groups. With a pulse of 2500 or 4000 rads did not disple the early incapacitation period (0-20 minutes preceiving the drug displayed a decrease in per response level in both the 2500- and 4000-rad of treated and untreated animals were not sign	ansient incapacitation period. Catheters rate and respiratory rate were surgi- e trained monkeys were trained to per- task. Following the injection of WR1607 sure with a significant decrease in heart et trials or latency occurred following e irradiated with a pulse of 2500, 4000 ation. After irradiation a significant -4 minutes in the 2500-rad group and After irradiation the respiratory rate Trained injected monkeys irradiated olay a decrease in performance during postirradiation). Trained monkeys not erformance below a 90 percent correct and groups. The physiology and behavior mificantly different after the first

₩÷@seeterse setter interfetter

100

 -----

Security Classification

<u>بيد اند فيه من المحمد من ا</u>

\*

.....

r stære

5-1-1-1-2

معمرين وحجر بالاحيا

ł

ia

ध्य-२०१४:अन्द्रात् स्टब्स् स्टब्स् केल्क्स केल्क्स केल्क्स केल्क्स

and the second second

 $\mathrm{ad}^{-1} \mathrm{H}^{-1}$ 

Ł

6

۰.

LIN	LINKA LINKB			LIN	LINKC			
ROLE	<b>₩</b> Т	ROLE	WT	ROLE	w			
				. :				
				1				
1								
		l –						
		1						
		[						
				}				
		}						
					ŀ			
	•							
				1 1				
ļ								
1								
		Li		I				
	UNC	LASSIF	IED					
	Security	Classifi	cation					

AFRRI SR72-3 March 1972

- 14

ģ

# ALTERATION OF RADIATION EFFECTS BY

nn-in thisisten an sin na state an air an an air an air

# 2-(n-DECYLAMINO) ETHANETHIOSULFURIC

ACID (WR1607) IN THE MONKEY

C. L. TURBYFILL R. M. ROUDON R. W. YOUNG V. A. KIEFFER

Davis, J W. F. DAVIS, JR.

bairman Behavioral Sciences Department

MYRON I. VARON Captain MC USN Director

and the second states of the second

ARMED FORCES RADIOBIOLOGY RESEARCH INSTITUTE Defense Nuclear Agency Bethesda, Maryland

Approved for public release; distribution unlimited i b

# ACKNOWLEDGMENT

The authors express their gratitude to M. H. Heiffer, Department of Pharmacology and D. E. Davidson, Jr., Division of Medicinal Chemistry, Walter Reed Army Medical Center, for furnishing WR1607 and their consultation during the study. Also recognized for their valuable assistance in training the animals are B. A. Dennison, P. Mannon, J. R. Harrison and G. G. Kessell, Behavioral Sciences Department, AFRRI.

# TABLE OF CONTENTS

•

1

非不可<del>这些就是那些化效地和能力和严格</del>的是是我们的

and and the state of the state

and more a surface during the second

1

-----

ł

																								Page
For	eword (No	nteo	chr	nica	l s	um	ma	ry)	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	iii
Absi	tract .	•	•	•	•	•	•	·	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	v
I.	Introduct	ion	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	1
ц.	Materials	s an	ld I	Met	hoc	ls	•	•	٠	•	•	•	•	•	•	•	•	•	•	•	•	•	•	1
ш.	Results	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	4
IV.	Discussio	n	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	10
v.	Summary	•	٠	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	11
Refe	erences.	•	•	•	•	•	٠	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	13

id

10.44

Ţ.

# AND AND AND ALL AND ALL

# LIST OF FIGURES

1

Figure 1.	Physiology of control animals injected with 2-(n-decylamino) ethanethiosulfuric acid 6
Figure 2.	Physiology of animals injected with 2-(n-decylamino) ethanethiosulfuric acid and irradiated 6
Figure 3.	Physiology of animals injected with 2-(n-decylamino) ethanethiosulfuric acid and irradiated
Figure 4.	Physiology of animals injected with 2-(n-decylamino) ethanethiosulfuric acid and irradiated
Figure 5.	Behavioral response of animals injected with 2-(n-decylamino) ethanethiosulfuric acid and irradiated with 2500 or 4000 rads of mixed gamma-neutron radiation

# LIST OF TABLES

Table	I.	Experimental Groupings	٠	2
Table	п.	Average Behavioral Response with Ranges before and following Irradiation	•	9

### FOREWORD (Nontechnical summary)

Numerous chemicals have been investigated to determine their ability to attenuate the effects of radiation injury. Most of the chemicals have been tested at radiation doses in the  $LD_{50/30}$  range using increased survival as indicators of protection. Sulfur containing N-substituted aliphatic amines have been found to be among the most promising radioprotective agents. One of these compounds 2-(n-decylamino) ethanethiosulfuric acid (WR1607) has been found to also alleviate the early transient incapacitation (ETI) observed in the monkey following a dose of 10,000, 20,000 or 40,000 rads. This indicates that WR1607 is capable of attenuating radiation effects over a very broad dose range.

In the present study WR1607 was given to monkeys prior to irradiation to investigate the effects of the compound on behavior and physiology during the ETI period. WR1607 was injected intravenously (10 mg/kg) 30 minutes before irradiation. Following the injection of WR1607 the animals appeared to become very apprehensive with increased movement and vocalization in the restraining chair. This period of increased activity lasted for 7-10 minutes postinjection and usually ended with the animal vomiting. After injection of WR1607 the aortic pressure increased and the heart rate decreased. However, the animals' ability to perform a learned task was not impaired during the period following injection.

The animals that were surgically implanted to monitor aortic and venous pressure, heart rate and respiratory rate were injected with WR1607 and irradiated with a pulse of 2500, 4000 or 15,000 rads of mixed gamma-neutror radiation. After irradiation, the respiratory rate increased in all radiation groups. The aortic pressure decreased below the elevated postirjection values and was significantly lower than the preinjection values from 2-4 minutes postirradiation in the 2500-rad group and from 1-2 minutes postirradiation in the 15,000-rad group. The aortic pressure values in the 4000-rad group were not significantly different from preinjection values.

The animals trained to perform a learned task were irradiated with a pulsed dose of 2500 or 4000 rads of mixed gamma-neutron radiation. Half of the trained animals in each group were injected with WR 1607. Following irradiation the injected groups of animals maintained performance values near preirradiation levels until 60-80 minutes postirradiation. The noninjected groups displayed a performance decrement during the first 20 minutes postirradiation. After the first 20-minute behavioral session the performance of the noninjected animals and WR 1607 injected animals was not significantly different.

The compound WR1607 is a radioprotective agent which appears capable of alleviating the ETI period observed in the monkey following a supralethal dose of radiation. The mechanism by which WR1607 prevents ETI or the mechanism by which it attenuates radiation injury in the lethal dose range is at present speculative. The mechanisms which may be involved and physiological and behavioral changes found in the present study are discussed.

iv

#### ABSTRACT

and the second se

2-(n-decylamino) ethanethiosulfuric acid (WR 1607) was given to trained and untrained monkeys (10 mg/kg) 30 minutes prior to irradiation to investigate its effects on behavior and physiology during the early transient incapacitation period. Catheters to monitor aortic and venous pressures, heart rate and respiratory rate were surgically implanted in the untrained monkeys. The trained monkeys were trained to perform a shock motivated visual discrimination task. Following the injection of WR1607 a significant increase occurred in aortic pressure with a significant decrease in heart rate. No significant changes in percent correct trials or latency occurred following injection. The untrained injected animals were irradiated with a pulse of 2500, 4000 or 15,000 rads of mixed gamma-neutron radiation. After irradiation a significant decrease in aortic pressure occurred from 2-4 minutes in the 2500-rad group and from 1-2 minutes in the 15,000-rad group. After irradiation the respiratory rate increased significantly in all injected groups. Trained injected monkeys irradiated with a pulse of 2500 or 4000 rads did not display a decrease in performance during the early incapacitation period (0-20 minutes postirradiation). Trained monkeys not receiving the drug displayed a decrease in performance below a 90 percent correct response level in both the 2500- and 4000-rad groups. The physiology and behavior of treated and untreated animals were not significantly different after the first 20-minute postirradiation period.

v

Many compounds have been investigated to determine their ability to attenuate radiation effects in animals receiving a control  $LD_{50/30}$  to  $LD_{99/30}$  radiation dose.<sup>9, 15</sup> Several sulfur containing N-substituted aliphatic amines have been shown to have radioprotective properties.<sup>3, 5</sup> One of these compounds 2-(n-decylamino) ethanethiosulfuric acid (WR1607) was found to also alleviate the early transient incapacitation (ETI) observed in the monkey following a dose of 10,000, 20,000 or 40,000 rads.<sup>10</sup> This compound (WR1607) prevented adrenergic vasodilation by phenoxybenzamine in the dog and restored the pressor effect of epinephrine following alpha adrenergic blockade. General vasoconstriction was evident following WR1607 treatment in the dog as a decrease in pulse pressure was observed without a change in systolic pressure.<sup>4</sup>

In the present study WR1607 was given to monkeys prio. to irradiation to investigate its effects on behavior and physiology during the period associated with radiationinduced early transient incapacitation.<sup>13</sup> It was anticipated that WR1607 would alleviate the behavioral and physiological decrement observed during this period.

#### II. MATERIALS AND METHODS

Monkeys were used in this study to evaluate the ability of WR1607 to attenuate the early transient incapacitation induced by radiation. One group of 23 female monkeys (<u>Macaca mulatta</u>) was utilized to study the effects of WR1607 on aortic pressure, venous pressure, heart rate and respiratory rate both preirradiation and postirradiation. A second group of 24 male monkeys was utilized to evaluate the effects of WR1607 on the animals' ability to perform a learned task preivradiation and postirradiation. The two groups were subdivided as shown in Table 1

Programmed dose (rads)	Average midline tissue dose <u>+</u> S. E.* (rads)	WR1607 10 mg/kg	Number of animals		
Physiology					
0	Control (sham irradiated)	yes	5		
2500	2230 ± 34	yes	6		
4000	3680 ± 41	yes	6		
15,000	14,830 ± 55	yes	6		
Behavior					
2500	2450 ± 56	yes	6		
2500	2630 ± 41	no	6		
4000	4000 ± 26	yes	6		
4000	4620 ± 10	no	6		

#### Table I. Experimental Groupings

\* S. E. = Standard error of the mean

Carlow No. 1900 States and the second se

The animals in the group selected for physiological studies were placed in primate restrain . :hairs for 3 days before surgery. A 24-hour fast preceded all surgery. Atropine sulfate (0.05 mg/lb, I. M.) was given before surgery and anesthesia was induced with phencyclidine HCl (1 mg/kg, I. M.). A surgical plane was maintained with sodium thiopental. The surgical procedure consisted of placing catheters (PE166 polyvinyl tubing) into the femoral artery and vein and advancing the respective catheters into the aorta and vena cava. Paired frontal and occipital dural stainless steel screws were also placed on the calvarium to record the electroencephalogram (EEG). At this time only the cardiovascular data will be presented. The catheters were flushed daily with physiological saline containing 0.1 percent heparin. Postoperative antibiotic merapy (Bicillin,\* 1 cc, I. M.) was initiated and continued for 5 days. The surgery was performed 5 days before irradiation. The aortic and \* Wyeth Laboratories, Philadelphia, Pennsylvania venous pressure, heart rate and respiratory rate were monitored for 2 hours on the day before irradiation to establish a preirradiation base line.

The behavioral groups of animals were trained to perform a simultaneous tisk using shock motivation. The apparatus and training procedures have been previously described.<sup>1</sup> Trials were presented at 10-second intervals, with the animal given 5 seconds to respond to the correct key. If the animal failed to respond within 5 seconds or made an incorrect response, it received a brief electrical shock. Each test period consisted of 100 trials and lasted 16.7 minutes. The test periods started at 20-minute intervals giving the animal a brief rest period following each 100-trial session.

Food was withheld from the monkeys in all groups for 16 hours before irradiation; however, water was available <u>ad libitum</u> except for a period of approximately 3 hours. This 3-hour period without water included preparation of the animal for irradiation and the time spent in the exposure room. Approximately 90 minutes before irradiation the animals were transported to the exposure room and positioned at a distance from the radiation source calculated to give the desired dosc. The behavioral animals were monitored while in an isolation cubicle within the exposure room. The appropriate connections were then made for monitoring either the physiology<sup>\*</sup> or behavior.

The monkeys were positioned in the exposure room of the AFRRI-TRIGA reactor to receive the calculated dose. The average midline tissue dose (MTD) and the number of animals in each group are presented in Table I. The MTD was obtained

3

- - And dela

<sup>\*</sup> Pressure transducers: arterial, Type P23Db; venous, Type P23BB, Statham Laboratories, Inc., Hato Rey, Puerto Rico Eight-channel recorder, Model 7700, Hewlett-Packard Company, Rockville, Maryland

by determining the tissue kerma, free-in-air, at the midline exposure volume and multiplying this value by an experimentally derived factor (0.85). For these exposures the operation of the reactor and the characteristics of the radiation field were as previously described.<sup>2</sup>

Before injection of the WR1607 via the venous catheter (physiology) or saphenous vein (behavior) the animals were monitored for 20 minutes or a 100-trial session, respectively. A 10 mg/kg dose of WR1607 was injected 30 minutes before irradiation (prepared by dissolving the powder in 0.1 N NaOH and bringing the solution to volume with physiological saline, 5 mg/ml pH 10.5 to 11.0). The postinjection behavioral session was started 20 minutes before irradiation. The physiology animals were monitored for 1 hour postirradiation and the behavioral groups for 2 hours or six behavioral sessions postirradiation.

The behavioral data were tabulated at 10-trial intervals with a 90 percent correct response as the minimum acceptable criterion. The WR1607 injected animals' performance was compared to the noninjected animals' performance using the "exact" test.<sup>8, 12</sup> Latency values of the two groups were analyzed using the Wilcoxon signed ranks test.<sup>12</sup>

The physiological data were statistically analyzed using the analysis of variance with a one-way layout for equal or unequal group size  $^{11}$  Values were considered significantly different when p < .05.

#### III. RESULTS

The aortic pressure, venous pressure, heart rate and respiratory rate of the groups of animals used in this study are shown in Figures 1-4.

Following the injection of WR 1607 a significant (p < .05) increase occurred in the aortic pressure with a significant decrease in heart rate in all groups when compared to preinjection values. With the increase in aortic pressure a decrease in pulse pressure occurred due to a large increase in diastolic pressure. The venous pressure and respiratory rate also increased but not to a significant level over preinjection values. All indices, except for aortic pressure, tended to return toward preinjection levels after 15 minutes postinjection. The aortic pressure remained elevated for approximately 70 minutes postinjection.

After irradiation the respiratory rate increased significantly in all injected groups when compared to preinjection or control nonirradiated injected values. The aortic pressure decreased to significant levels in the 2500-rad group from 2-4 minutes postirradiation and in the 15,000-rad group from 1-2 minutes postirradiation when compared to preinjected values; but no significant change was observed in the 4000-rad group. When compared to the WR1607 injected controls, a significant difference in aortic pressure occurred from 1-30 minutes postirradiation in the 2500rad group, from 1-5 minutes postirradiation in the 4000-rad group, and from 1-60 minutes postirradiation in the 15,000-rad group. The venous pressure and heart rate did not change significantly from preinjection values following irradiation. The heart rate in the 2500-rad group was significantly higher than the control injected group up to 10 minutes postirradiation.

The percent correct and latency data are presented in Figure 5 and Table II. Each point in Figure 5 represents either the percentage of correct responses or the average time required to complete a response during a 10-trial problem.











The second secon







No significant changes in percent correct trials or latency were observed in the session following the injection of WR 1607. After irradiation a nonsignificant decrease in performance and a nonsignificant increase in latency occurred during the first four 10-trial blocks of problems of the first postirradiation session. Animals which did not receive WR 1607 displayed changes to below acceptable values in these indices and did not return to preirradiation values during the first 100-trial session (Figure 5).

att ...

į

۱



Figure 5. Behavioral response of animals injected with 2-(n-decylamino) ethanethiosulfuric acid and irradiated with 2500 or 4000 rads of mixed gamma-neutron radiation

		Tri (percent	als correct)		Latency (seconds/trial)						
[nterval	2500-rad	2500-rad	4000-rad	4000-rad	2500-rad	2500-rad	4000-rad	4000-rad			
	control	injected	control	injected	control	injected	control	injected			
Preinjection	99.2	97.8	96.8	98.0	1,13	1.25	1.32	1.49			
	98-100	94-100	95-100	95-100	.98-1.98	.95-1.69	.83-1.87	.88-1.89			
Postinjection		99.5 99-100		98.3 98-100		1.31 1.02-1.79		1,61 ,94-1,96			
Pulse											
0-10 minutes	71.3	99.5	29.6**	91.5	2.16 <sup>*</sup>	1.79	3.42 <sup>*†</sup>	1,79			
	34-98	92-100	10-64	76-100	1.11-2.64	.98-2.63	1,06-5.00	1,10-3,11			
10-20 minutes	73.4	99.7	36.6* <sup>**</sup>	98.8	2.09 <sup>+</sup>	1.43	3.75**	1,43			
	0-100	98-100	0-96	96-100	1.76-2.64	.92-2.01	1.10-5.00	.80-1.76			
20-40 minutes	99.0 98-100	99.0 97-100	72.3 <sup>+</sup> 3-94	98.9 98-99	$1,21^{+}$ ,97-1,40	1.49 .94-1.81	2.18 <sup>+</sup> .78-4.71	1.56 .98-2.05			
40-60 minutes	94.5	93.2	63.8 <sup>*</sup>	82.0	$1.79^{+}$	1.81 <sup>+</sup>	2.49 <sup>†</sup>	2.31 <sup>†</sup>			
	83-100	84-98	20-97	59-96	1.05-2.34	1.18-2.24	1.19-4.10	1.49-3.21			
	97.7	87.2	67.0	$54.2^+$ 0-97	1.84 <sup>*</sup>	2.06 <sup>+</sup>	2,53 <sup>†</sup>	3,15*			
60-80 minutes	95-99	41-100	0-98		1.29-2.48	1.27-2.71	1,48-5,00	1,48-5,00			
80-100 minutes	98.8 96-100	83.3 10-100	70.6 1-98	48.8 0-98	$1.67^{+}$ .99-2.21	2.14 <sup>+</sup> 1.01-4.48	2.34 .86-4.32	3.36 <sup>†</sup> 1.43-5.00			
100-120 minutes	98.7	81.3	79.2	69.0	1,60 <sup>+</sup>	1,98 <sup>+</sup>	2,23 <sup>†</sup>	2.70 <sup>+</sup>			
	96-100	0-100	38-99	5-9ນ	1,05-2,21	1,14-5.00	.89-4.20	.99-5.00			

# Table II. Average Behavioral Response with Ranges before and following Irradiation

\* Significantly different from WR1607 treated group at .05 percent level \* Significantly different from own preinjection group at .05 percent level

The animals receiving WR1607 maintained percent correct and latency values near preirradiation levels until the third session postirradiation (40-60 minutes). During the fourth session following irradiation (60-80 minutes), in both the 2500- and 4000-rad WR1607 treated groups, the percent correct responses decreased. The WR1607 injected animals' values were not significantly different from the control groups' values for any behavioral session after the first 100-trial session postirradiation (0-20 minutes).

Following the injection of WR1607 the animal appeared to become very apprehensive with increased movement in the restraining chair. In most instances the animal vocalized during the injection. This period of increased activity lasted for 7-10 minutes postinjection and usually culminated in the animal vomiting 8-10 minutes postinjection. After the animal vomited the period of apprehension appeared to end with no further overt effects of the drug being evident.

Following irradiation the WR 1607 treated animals appeared much more alert than nontreated animals for 30 to 40 minutes. After 40 minutes postirradiation there was no discernible difference.

#### IV. DISCUSSION

The data obtained in this study show WR1607 to be beneficial in eliminating the early transient incapacitation (ETI) period observed in monkeys receiving a single pulse of 2500 or 4000 rads of radiation.

The physiological mechanisms which produce the ETI period in the monkey after a supralethal dose of radiation are unknown. Therefore, it is not possible to define the mechanisms by which WR1607 is able to attenuate the radiation effect in the monkey. From the data presented in this study it would appear that a state of hypertension was induced by WR1607 through the beta adrenergic-like properties attributed to the compound.<sup>4</sup> Other investigations have indicated that the ETI may be initiated by a change in blood flow to various areas of the brain.<sup>\*, 14, 16</sup> Pretreatment with WR1607 because of its adrenergic properties may inhibit these changes, thus alleviating the ETI period.

<sup>\*</sup>Unpublished: Turbyfill, C. L. and Flinton, J. H., Armed Forces Radiobiology Research Institute, Bethesda, Maryland

The chemical properties of WR1607 must also be considered to elucidate the mechanisms by which this class of compounds attenuate radiation injury. Since the aminoethanethiols can be cleaved hydrolytically, the radioprotective mechanism of WR1607 may be due to a release of sulfides by hydrolysis, freeing the thiol group to form mixed disulfides.<sup>6,7</sup> However, the radioprotective mechanisms of WR1607 are at this time speculative and further research in this area is needed.

Although WR1607 shows promise as a radioprotective compound, it has properties which may limit its use in man.<sup>4</sup> It must be given prior to irradiation in order to obtain the radioprotective effect. In addition, side effects such as cardiac toxicity and vasoconstriction occur following its administration. In the present study a dose of 10 mg/kg was used which is approximately one-third the LD<sub>50</sub> dose. At this level the undesirable effects were observed but the degree of physiological or behavioral impairment did not exceed acceptable limits. However, due to the toxic properties of WR1607, future investigative effort will be conducted with less toxic compounds, such as S-2-(3-aminopropyl) aminoethyl phosphorothioic acid (WR2721AF).

#### V. SUMMARY

2-(n-decylamino) ethanethiosulfuric acid (WR1607) was given to monkeys (10 mg/kg) 30 minutes prior to irradiation to investigate its effects on behavior and physiology during the early incapacitation period. Following the injection of WR1607 a significant increase occurred in aortic pressure with a significant decreas in heart rate. No significant changes in percent correct trials or latency occurred following the injection of WR1607 in trained monkeys. After irradiation with 2500, 4000, or 15,000 rads the WR1607 injected animals displayed a decrease in aortic pressure from the

elevated levels following injection. The decrease in aortic pressure was significantly different from preirradiation values 2-4 minutes postirradiation in the 2500-rad group and from 1-2 minutes in the 15,000-rad group. The respiratory rate increased significantly in all irradiated groups postirradiation. Trained monkeys receiving WR 1607 did not display a decrease in performance during the usual FTI period (0-20 minutes postirradiation) following a dose of 2500 or 4000 rads. Trained monkeys not receiving the drug all displayed a decrease in performance below a 90 percent correct response level in both the 2500- and 4000-rad groups. The physiology and behavior of WR 1607 treated and untreated animals were not significantly different after the first 20-minute postirradiation period.

#### REFERENCES

- de Haan, H. J., Germas, J. E. and Kaplan, S. J. Visual discrimination performance: a training procedure for the restrained monkey (<u>Macaca mulatta</u>). Bethesda, Maryland, Armed Forces Radiobiology Research Institute Technical Note TN68-5, 1968.
- Dowling, J. H. Experimental determination of dose for the monkey in a reactor pulse environment. Bethesda, Maryland, Armed Forces Radiobiology Research Institute Scientific Report SR66-3, 1966.
- Hamilton, H. E., Melville, G. S., Jr. and Stork, E. J. Radioprotection of primates with 2-(1-decylamino)ethanethiosulfuric acid in dimethyl sulfoxide. Brooks Air Force Base, Texas, U. S. Air Force School of Aerospace Medicine Report TR68-137, 1968.
- Heiffer, M. H., Mundy, R. L., Demaree, G. E., Jacobus, D. P., Herman, E. H. and Brockenton, R. E. Beta adrenergic blocking properties of 2-(n-decylaminc) ethanethiosulfuric acid. Washington, D. C., Walter Reed Army Institute of Research (personal communication), 1971.
- 5. Jacobus, D. P., Klayman, D. L., Rothe, W., Grenan, M. M., Henderson, E. and Davidson, D. E. N-substituted aminoethanethiosulfuric acids as potential anti-radiation agents. Pharmacologist 8:226 (Abstract 311), 1966.
- Kelley, J. J., Hamilton, N. F. and Friedman, O. M. Studies on latent derivatives of aminoethanethiols as potentially selective cytoprotectants. III. Reactions of cysteamine-S-sulfate in biologic media. Cancer Res. 27:143-147, 1967.
- Kelley, J. J., Herrington, K. A., Ward, S. P., Meister, A. and Friedman, O. M. Studies on latent derivatives of aminoethanethiols as potentially selective cytoprotectants. II. In vivo distribution of cysteamine liberated in rat tissues, Cancer Res. 27:137-142, 1967.
- 8. Mainland, D., Herrera, L. and Sutcliffe, M. I. Statistical Tables for Use with Binomial Samples. New York, N. Y., New York University College of Medicine, Department of Medical Statistics, 1956.
- 9. Plzak, V. and Doull, J. A further survey of compounds for radiation protection. Brooks Air Force Base, Texas, U. S. Air Force School of Aerospace Medicine Report TR69-1, 1969.

- Sharp, J. C., Kelly, D. D. and Brady, J. V. The radio-attenuating effects of n-decylaminoethanethiosulfuric acid in the Rhesus monkey. In: Use of Nonhuman Primates in Drug Evaluation, A Symposium, pp. 338-346. Austin and London, University of Texas Press, 1967.
- 11. Snedecor, G. W. and Cochran, W. G. Statistical Methods, 6th ed. Ames, Iowa, The Iowa State University Press, 1967.
- 12. Steel, R. G. D. and Torrie, J. H. Principles and Procedures of Statistics. New York, N. Y., McGraw-Hill Book Company, Inc., 1960.
- Turbyfill, C. L., Kieffer, V. A. and Dewes, W. A. Cardiovascular response of monkeys to supralethal doses of mixed gamma-neutron radiation. Bethesda, Maryland, Armed Forces Radiobiology Research Institute Scientific Report SR70-10, 1970.
- Turbyfill, C. L., Roudon, R. M. and Kieffer, V. A. Behavior and physiology of the monkey (Macaca mulatta) following 2500 rads of pulsed mixed gammaneutron radiation. Aerospace Med. 43:41-45, 1972.
- Turns, J. E., Doyle, T. F. and Curran, C. R. Norepinephrine effects on early postirradiation performance decrement in the monkey. Bethesda, Maryland, Armed Forces Radiobiology Research Institute Scientific Report SR71-16, 1971.
- Watters, J. W., Wailly, L. F. and Carter, P. B. Radiation induced changes in intracranial pressure. Brooks Air Force Base, Texas, U. S. Air Force School of Aerospace Medicine Report (in press).