

AD-770 560

**CORRELATION OF EYE-LEVEL BLOOD FLOW
VELOCITY AND BLOOD PRESSURE DURING +GZ
ACCELERATION**

Robert W. Krutz, Jr., et al

**School of Aerospace Medicine
Brooks Air Force Base, Texas**

November 1973

DISTRIBUTED BY:

NTIS

**National Technical Information Service
U. S. DEPARTMENT OF COMMERCE
5285 Port Royal Road, Springfield Va. 22151**

NOTICES

This final report was submitted by personnel of the Biodynamics Branch, Environmental Sciences Division, USAF School of Aerospace Medicine, AFSC, Brooks Air Force Base, Texas; Ames Research Center, National Aeronautics and Space Administration, Sunnyvale, California; and University of Santa Clara, Santa Clara, California, under job order 7930-03-25.

When U.S. Government drawings, specifications, or other data are used for any purpose other than a definitely related Government procurement operation, the Government thereby incurs no responsibility nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise, as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use, or sell any patented invention that may in any way be related thereto.

The voluntary informed consent of the subjects used in this research was obtained as required by Air Force Regulation 80-33.

This report has been reviewed and cleared for open publication and/or public release by the appropriate Office of Information (OI) in accordance with AFR 190-17 and DODD 5230.9. There is no objection to unlimited distribution of this report to the public at large, or by DDC to the National Technical Information Service (NTIS).

This technical report has been reviewed and is approved for publication.

Robert W. Krutz Jr.

ROBERT W. KRUTZ, JR., Major, USAF, BSC
Project Engineer/Scientist

Evan R. Goles

EVAN R. GOLES, Colonel, USAF, MC
Commander

ACCESSION for	
NTIS	WFO Section <input checked="" type="checkbox"/>
DOC	Ref Section <input type="checkbox"/>
UNCLASSIFIED	
JUSTIFICATION	
BY _____	
DISTRIBUTION/AVAILABILITY CODES	
DECL.	AVAIL. FOR/OF SPECIAL
A	

16

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER SAM-TR-73-36	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER AD 770 560
4. TITLE (and Subtitle) CORRELATION OF EYE-LEVEL BLOOD FLOW VELOCITY AND BLOOD PRESSURE DURING +G_z ACCELERATION		5. TYPE OF REPORT & PERIOD COVERED Final (Jul 72-Jun 73)
7. AUTHOR(s) Robert W. Krutz, Jr., S. A. Rositano, and R. E. Mancini		6. PERFORMING ORG. REPORT NUMBER SAM-TR-73-36
9. PERFORMING ORGANIZATION NAME AND ADDRESS USAF School of Aerospace Medicine (VNB) Aerospace Medical Division (AFSC) Brooks Air Force Base, TX 78235		8. CONTRACT OR GRANT NUMBER(s)
11. CONTROLLING OFFICE NAME AND ADDRESS USAF School of Aerospace Medicine (VNB) Aerospace Medical Division (AFSC) Brooks Air Force Base, TX 78235		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 7930-03-25
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE November 1973
		13. NUMBER OF PAGES 5
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Doppler flowmeter +G_z tolerance Noninvasive technique		
Reproduced by NATIONAL TECHNICAL INFORMATION SERVICE U S Department of Commerce Springfield VA 22151		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) <p>Eye-level blood flow and blood pressure changes were correlated on the USAFSAM human centrifuge during both rapid onset (ROR, 1 G/sec) and gradual onset runs (GOR, 0.1 G/sec). A transcutaneous Doppler ultrasonic flowmeter was used to monitor temporal artery blood flow (\dot{Q}_{ta}); direct blood pressure was obtained by cannulation of a radial artery and measured at eye level with a Statham P-37 miniature transducer. Eye-level mean blood pressure (P_a) decreased to 20 mm Hg and zero forward \dot{Q}_{ta} occurred 6 sec (range 4-9 sec) prior to blackout in experienced centrifuge subjects during RORs. The same degree of correlation was not seen during GORs.</p>		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

CORRELATION OF EYE-LEVEL BLOOD FLOW VELOCITY AND BLOOD PRESSURE DURING $+G_z$ ACCELERATION

INTRODUCTION

The need for a reliable objective means for monitoring $+G_z$ tolerance has long been recognized by acceleration physiologists. Loss of eye-level arterial pressure has been correlated with cessation of blood flow to the retina (1) and remains the most reliable indication of cardiovascular status during exposure to $+G_z$ acceleration. This technique, however, has the distinct disadvantage of being invasive, requiring cannulation of a radial artery.

This study was designed to examine the efficacy of a transcutaneous Doppler flowmeter (2) monitoring eye-level (temporal artery) blood flow velocity during $+G_z$ acceleration and to correlate results with direct arterial pressure referenced to eye level.

This noninvasive technique would quantitate man's ability to withstand G without using invasive monitoring techniques, and thus altering his normal response to G stress because of the instrumentation.

METHODS

Seven healthy male volunteer subjects (age range 21 to 25) were studied. All had recently passed a USAF Class II flying physical examination and had extensive centrifuge experience.

Subjects were instrumented with miniature 8 MHz Doppler sensors (2 x 1 x 0.5 cm) secured to the skin above the maximum palpable impulse from both the right and left frontal branches of the temporal arteries to detect blood flow velocity from back-scattered ultrasound. A directional signal processor was used with one sensor while a nondirectional processor was used with the other. With this arrangement, retrograde flow was graphically portrayed. Audio recordings of the unprocessed Doppler-shift were also made. The right radial artery was cannulated, and eye-level arterial blood pressure was measured using a Statham (P-37) miniature strain gauge transducer mounted at eye level. Mean arterial blood pressure was obtained by electronically damping the arterial pressure wave with appropriate filtering.

EKG was continuously monitored, and audiovisual communication was maintained with the subject at all times.

Both rapid onset runs (ROR, 1 G/sec) and gradual onset runs (GOR, 0.1 G/sec) were used to stress subjects to the point of visual failure on the USAFSAM human centrifuge. RORs were begun at peak 2.5 G for 15 sec and increased in increments of 0.5 G until peripheral light loss (PLL) occurred; at this time, runs were increased by 0.2 G until the end-point of blackout (50% loss of central light). Subjects then underwent a GOR to blackout. Adequate time was allowed between runs for the subject to return to a normal physiologic state.

RESULTS

When blackout $+G_z$ level (range 2.7 to 4.6 G) was approached during rapid onset runs, eye-level arterial blood pressure began to fall concomitant with the occurrence of retrograde flow in the temporal artery during diastole (Fig. 1). This occurrence of retrograde flow has been verified using both directional and nondirectional Doppler systems (3). It can be easily recognized with audio recordings of the Doppler-shift. Zero forward temporal artery flow (\dot{Q}_{ta}) was determined by both graphic and audio recordings 6 sec prior to blackout (range 4-9 sec). Eye-level mean arterial pressure (\bar{P}_a) decreased to 20 mm Hg when zero forward \dot{Q}_{ta} was recorded. Arterial pressure and \dot{Q}_{ta} increased simultaneously during centrifuge deceleration with a characteristic increase in arterial pressure and flow occurring postrun when compared to prerun values.

The correlation of mean arterial pressure, temporal artery blood flow, and visual symptoms recorded during rapid onset runs was not duplicated to the same degree during gradual onset runs. Although changes in mean arterial pressure and \dot{Q}_{ta} occurred simultaneously, sustained zero forward flow was recorded in only 2 subjects prior to blackout. In these subjects, the onset of sustained zero forward \dot{Q}_{ta} occurred simultaneously with a decrease in mean arterial pressure to 20 mm Hg and occurred 5 sec prior to blackout (Fig. 2).

In a third subject, mean arterial pressure was sustained at 20 mm Hg and below for 9 sec prior to blackout, and zero net forward flow occurred for the same duration; the nondirectional flowmeter was not functioning. In the four remaining subjects, zero sustained forward flow was never attained and mean arterial pressure was maintained at higher levels; nevertheless, the retrograde flow portion of the \dot{Q}_{ta} wave had become progressively dominant.




Figure 1. Eye-level arterial pressure and blood flow responses during ROR (1 G/sec).




Figure 2. Eye-level arterial pressure and blood flow responses during GOR (0.1 G/sec).

Reproduced from
best available copy.

DISCUSSION

Coburn (3), as recently as 1970, voiced his concern regarding the "reliability and repeatability of a large portion of the data reported in the literature." Many of his objections to subjective endpoints have been verified by other acceleration physiologists. An inexperienced centrifuge subject may terminate a run because of fear or a misunderstanding of the desired endpoint. Even an experienced, highly motivated subject will terminate a run prematurely if unduly fatigued from previous runs. It was the purpose of this study to determine if a correlation could be established between the objective measurements of temporal artery blood flow velocity, mean arterial pressure, and subjective blackout in highly trained, experienced centrifuge subjects.

During the rapid onset runs, zero forward flow and a mean arterial pressure of 20 mm Hg occurred 6 sec prior to blackout. It is assumed that at this eye-level arterial pressure, the critical closing pressure of branches of the temporal artery is approached (4).

Duane (5) using direct ophthalmoscopic observations and Leverett and Newsom (1) using retinal photography and fluorescence angiography found that subjective blackout coincides with cessation of flow in the retinal circulation. They also found that visual failure at blackout occurred when head-level arterial pressure had fallen below 20 mm Hg. Our findings using transcutaneous ultrasound suggest that blood flow velocity changes in the retinal circulation are reflected in flow velocity changes in frontal branches of the temporal arteries during rapid onset +G_z acceleration.

During the gradual onset runs, blackout occurred before a sustained zero forward flow and mean pressure of 20 mm Hg had been reached in 4 of 7 subjects. This could, in part, be attributed to a combination of hypoxic hypoxia, resulting from reduced oxygenation of the blood during these prolonged +G_z exposures (6) and stagnant hypoxia, since the retrograde component of the flow wave gradually increased until blackout was reached.

CONCLUSIONS

The transcutaneous Doppler ultrasonic flowmeter monitoring \dot{Q}_{ta} appears to be a reliable tool for measuring cardiovascular status and predicting visual failure during rapid onset $+G_z$ acceleration. Cardiovascular status as indicated by mean arterial pressure is well correlated with changes in temporal artery blood flow and visual symptoms reported by experienced centrifuge subjects. Cardiovascular status during gradual onset runs also appears to be reflected by simultaneous changes in temporal artery blood flow and mean arterial pressure; however, the accurate prediction of visual failure during these prolonged runs requires further investigation. Work is continuing using this new noninvasive technique to assess cardiovascular status during high, sustained $+G_z$ acceleration and in the evaluation of new $+G_z$ protective devices and techniques.

REFERENCES

1. Leverett, S. D., Jr., and W. A. Newson. Photographic observations of the human fundus oculi during $+G_z$ blackout on the USAF School of Aerospace Medicine Centrifuge. *In* Lunc, M. (ed.). XIXth International Astronautical Congress Bioastronautics - Book 4. Oxford: Pergamon, 1971.
2. Rositano, S. A., et al. Noninvasive determination of retrograde eye-level blood flow as a $+G_z$ tolerance indicator. Proceedings of the 44th Annual Scientific Meeting, Aerospace Medical Association, Las Vegas, Nev., 7-10 May 1973.
3. Coburn, K. R. Physiological endpoints in acceleration research. *Aerosp Med* 41:5 (1970).
4. Nichol, J., et al. Fundamental instability of the small blood vessels and critical closing pressures in vascular beds. *Am J Physiol* 164:330 (1951).
5. Duane, T. D. Observations on the fundus oculi during black-out. *Arch Ophthalmol* 51:343 (1954).
6. Michaelson, E. D. Blood oxygenation in man during high, sustained $+G_z$. Proceedings of the 43rd Annual Scientific Meeting, Aerospace Medical Association, Bal Harbour, Fla., 8-11 May 1972.