

JUL 2 1973

UNCLASSIFIED

Security Classification

AD773468

DOCUMENT CONTROL DATA - R &

(Security classification of title, body of abstract and indexing annotation must be entered)

1. ORIGINATING ACTIVITY (Corporate author)

The Chemical Laboratory
 University of Sussex
 Falmer, Brighton BN1 9QJ, England

2a. REPORT SECURITY CLASSIFICATION

Unclassified

2b. GROUP

3. REPORT TITLE

COMPUTER ANALYSIS OF THERMO CHEMICAL DATA: SILICON COMPOUNDS

4. DESCRIPTIVE NOTES (Type of report and inclusive dates)

Scientific. Interim.

5. AUTHOR(S) (First name, middle initial, last name)

J. B. Pedley, A. Kirk, S. Seilman, and L. G. Heath

6. REPORT DATE

September 1972

7a. TOTAL NO. OF PAGES

7b. NO. OF REFS

79

8a. CONTRACT OR GRANT NO.

AFOSR-72-2220

9a. ORIGINATOR'S REPORT NUMBER(S)

b. PROJECT NO.

9537

c.

61102F

d.

681303

9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)

AFOSR - TR - 73 - 1741

10. DISTRIBUTION STATEMENT

Approved for public release; distribution unlimited

11. SUPPLEMENTARY NOTES

12. SPONSORING MILITARY ACTIVITY

Air Force Office of Scientific
 Research/NC, 1400 Wilson Blvd.
 Arlington, Va. 22209

13. ABSTRACT

Thermodynamic data on silicon compounds from numerous sources were stored, sorted and processed by computer. Enthalpies of reactions were determined by the computer from the available data by a least squares procedure. Computer controlled automatic typesetting was used to prepare these tables of enthalpies.

Reproduced by
 NATIONAL TECHNICAL
 INFORMATION SERVICE
 U S Department of Commerce
 Springfield VA 22151



14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
COMPUTER ANALYSIS						
HEATS OF FORMATION						
SILICON COMPOUNDS: STANDARD HEATS OF FORMATION						
THERMOCHEMICAL DATA						

ia/



Introduction

The first half of each section of the CATCH tables consists of a list of pure substances and aqueous solutions in the standard order of arrangement described in detail in ref.1. The standard enthalpies of formation and the associated uncertainties have been calculated from the standard enthalpy changes of the numbered reactions listed in the second half.

The following symbols are used to indicate the states of the substances involved.

- (c) crystalline solid
- (l) liquid
- (g) gaseous
- (am) glassy or amorphous
- (a) completely dissociated species at infinite aqueous dilution
- (ao) undissociated species at infinite aqueous dilution

For solutions which are not infinitely dilute, the number of moles of water associated with one mole of solute is indicated in brackets.

Elements in their standard reference states are indicated by the symbols (cs), (ls) and (gs) and these, with the hydrogen ion at infinite dilution and the gaseous electron, are defined to have zero enthalpy of formation. (N.B. the electron is omitted from the equations corresponding to the processes in which it is involved).

Since compounds are identified by the computer by their molecular weights, isomers have been distinguished by an appropriate number after the symbol defining the state of the substance.

Some of the compounds at the end of the section do not belong to that set of compounds being explicitly considered, but they are present in the reactions. The standard enthalpies of formation of these species have been derived from other sections of the CATCH tables where this is possible, or from references 1, 2 or 3 in that order of preference. Data on organic species have been obtained from reference 4. The uncertainties are assigned as zero unless they are derived from other CATCH tables in which case they are the computer processed values.

In the reactions section, the reaction number is that referred to in the compounds section as contributing to the standard enthalpy of formation of a specific compound. The reactions form an overdetermined set of equations for the calculation of standard enthalpies of formation, so the data was analysed by standard least squares procedure using the reciprocal of the uncertainty on the enthalpy of reaction as a weight. The residuals associated with the reactions are the differences between the experimental value of the standard enthalpy of reaction and that calculated from the derived standard enthalpies of formation of the species in the reaction. These residuals are printed only when greater than 0.001kcal/mol and are a measure of the compatibility of a specific reaction with the others. The uncertainties (errors) on the standard enthalpies of formation were calculated from the uncertainties (errors) on the standard enthalpies of reaction by standard statistical methods (see, for example, ref.4, p.43).

Any species already considered by the CODATA task group (ref.1) are assumed to have well defined standard enthalpies of formation and no attempt has been made to reanalyse the data leading to those values. The formation reaction of the species from the elements are included (labelled COD), and the weight of that equation in the least squares process has been increased to such an extent that the residual is always less than 0.001kcal/mol.

To make the list of compounds as comprehensive as possible, values of the enthalpy of the (often) hypothetical formation reaction of some compounds from their elements have been taken from reference 2 (and labelled NBS). Because the data sources leading to the selection of NBS values are at present unlisted, no further attempts at critical assessment have been made, and the uncertainty interval is assumed to be ten times the last figure given in reference 2. However, wherever

enthalpies of formation can be derived from critically assessed data without resort to unassessed values, this procedure has been adopted and the NBS data disregarded. It is intended that the number of references to NBS will be gradually reduced as critical assessment becomes possible. NBS data has also been used to provide enthalpies of solution and dilution.

Also included in the list of reactions are equations defining the equivalence of the enthalpies of a completely ionised solute at infinite dilution and the ions which are assumed to constitute the molecule of the electrolyte.

The uncertainties on many of the experimental standard enthalpies of reaction have been increased compared with the values quoted in the literature, when it was felt by the selector that large systematic errors may possibly be present. This is a very subjective operation and suggestions concerning the reliability of the data in the tables would be welcomed by the selector.

References

- 1 (COD) Report of CODATA Task Group on Key Values for Thermodynamics, Nov. 1971. CODATA BULLETIN No. 6, 1971.
- 2 (NBS) Wagman, Evans, Parker, Harlow, Bailey, and Schumm, *Selected Values of Chemical Thermodynamic Properties*, Natl. Bur. Stand. Tech. Note 270-3 U.S. Government Printing Office, Washington D.C., 1968.
- 3 Rossini, Wagman, Evans, Levine and Jaffe, *Selected Values of Chemical Thermodynamic Properties*, Natl. Bur. Stand. Circ. 500.
- 4 Cox and Pilcher, *Thermochemistry of Organic and Organometallic Compounds*, Academic Press, London, 1970.

J. B. Pedley,
School of Molecular Sciences,
University of Sussex.

Silicon compounds

Compound	Formula weight	ΔH_f° (kcal/mol)	Error	ΔH_f° (kj/mol)	Reactions for ΔH_f°
Si(cs)	28.086	0.000	0.000	0.000	0
Si(am)	28.086	1.000	0.101	4.184	2 20 23
Si(l)	28.086	11.586	0.101	48.475	1 115 128 130
Si(g)	28.086	107.920	0.863	451.536	3 36 37 39 40 41 123 125 127 129 216
Si ⁺ (g)	28.086	297.370	0.869	1244.195	4
Si ₂ (g)	56.172	141.840	3.754	593.457	5
Si ₃ (g)	84.258	152.759	10.330	639.145	6
SiO(g)	44.085	-24.775	1.590	-103.658	7 8 9 10
SiO ⁺ (g)	44.085	224.225	10.126	938.158	11
SiO ₂ (c1)	60.085	-217.997	0.271	-912.101	7 8
Alpha, quartz					12 13 14 47
SiO ₂ (c2)	60.085	-217.494	0.339	-909.995	10 13
Alpha, tridymite					14 44
SiO ₂ (c3)	60.085	-217.397	0.289	-909.590	9 12 17 45
Alpha, cristobalite					46
SiO ₂ (am)	60.085	-215.744	0.269	-902.674	15
SiO ₂ (l)	60.085	-216.047	0.406	-903.942	16
SiO ₂ (g)	60.085	-74.397	10.004	-311.278	17 81
SiO ₂ (a)	60.085	-217.244	0.288	-908.950	82 192 193 194
Colloidal solution					
SiO ₂ ⁺ (g)	60.085	195.603	14.145	818.402	18
SiH(g)	29.094	90.023	3.443	376.655	19
SiH ₂ ⁺ (g)	30.102	285.291	7.073	1193.658	21 25
SiH ₃ (g)	31.110	54.250	11.191	226.984	26
SiH ₄ ⁺ (g)	31.110	234.749	5.010	982.189	22
SiH ₄ (g)	32.118	8.302	0.319	34.735	20 21
					25
Si ₂ H ₅ ⁺ (g)	61.212	229.896	10.007	961.885	27
Si ₂ H ₆ (l)	62.220	14.499	0.619	60.664	24
Si ₂ H ₆ (g)	62.220	19.099	0.364	79.911	23 25
Si ₃ H ₈ (l)	92.322	22.200	1.158	92.885	29
Si ₃ H ₈ (g)	92.322	28.900	1.045	120.918	28
H ₂ SiO ₃ (c)	78.100	-284.100	1.000	-1188.674	30
H ₂ SiO ₃ (ao)	78.100	-282.700	1.005	-1182.817	31
H ₄ SiO ₄ (c)	96.116	-354.000	1.000	-1481.136	32
H ₄ SiO ₄ (ao)	96.116	-351.000	1.005	-1468.584	33
H ₂ Si ₂ O ₅ (c)	138.185	-499.200	1.000	-2088.653	34
H ₆ Si ₂ O ₇ (c)	174.216	-638.000	1.000	-2669.392	35
SiF(g)	47.084	-4.508	1.889	-18.861	36 37
SiF ⁺ (g)	47.084	163.892	5.345	685.725	55

Compound	Formula	ΔH_f°	Error	ΔH_f°	Reactions
	weight	(kcal/mol)	(kj/mol)	for ΔH_f°	
$\text{SiF}_2(\text{c})$	66.083	-186.289	2.009	-779.431	38
Polymer					
$\text{SiF}_2(\text{g})$	66.083	-140.786	1.122	-589.048	37 39
					40 41
					56
$\text{SiF}_2^+(\text{g})$	66.083	120.172	8.199	502.801	56 60
$\text{SiF}_3(\text{g})$	85.081	-227.269	14.143	-950.892	57
$\text{SiF}_3^+(\text{g})$	85.081	-31.269	10.002	-130.828	58
$\text{SiF}_3^-(\text{g})$	85.081	-304.269	14.284	-1273.060	42
$\text{SiF}_4(\text{g})$	104.080	-385.989	0.191	-1614.976	43 44
					45 46
					47 48
					60 122
					124
$\text{SiF}_4^+(\text{g})$	104.080	-23.689	10.002	-99.113	59
$\text{SiF}_6^-(\text{a})$	142.076	-571.000	1.000	-2389.064	49
$\text{Si}_2\text{F}_4^+(\text{g})$	132.166	-105.997	18.278	-443.493	64
$\text{Si}_2\text{F}_5(\text{g})$	151.164	-366.617	20.834	-1533.927	63
$\text{Si}_2\text{F}_5^-(\text{g})$	151.164	-203.558	12.916	-851.687	61
$\text{Si}_2\text{F}_6(\text{c})$	170.162	-575.958	8.189	-2409.809	51
$\text{Si}_2\text{F}_6(\text{l})$	170.162	-572.058	8.231	-2393.491	53
$\text{Si}_2\text{F}_6(\text{g})$	170.162	-565.558	8.174	-2366.295	50 60
$\text{Si}_3\text{F}_7^+(\text{g})$	217.247	-414.266	18.277	-1733.288	65
$\text{Si}_3\text{F}_8(\text{c})$	236.245	-769.686	15.307	-3220.366	52
$\text{Si}_3\text{F}_8(\text{l})$	236.245	-764.386	15.329	-3198.190	54
$\text{Si}_3\text{F}_8(\text{g})$	236.245	-755.586	15.298	-3161.371	62
$\text{Si}_3\text{F}_8^+(\text{g})$	236.245	-505.586	18.277	-2115.371	66
$\text{SiH}_3\text{F}(\text{g})$	50.108	-105.000	5.000	-439.320	72
$\text{H}_2\text{SiF}_2(\text{g})$	68.099	-194.000	10.000	-811.696	73
$\text{SiHF}_3(\text{g})$	86.089	-283.000	5.000	-1184.072	74
$\text{H}_2\text{SiF}_6(20\text{H}_2\text{O})$	144.092	-575.530	0.512	-2408.016	67
$\text{H}_2\text{SiF}(\text{a1})$	144.092	-575.373	0.473	-2407.362	68 71
In 1.444 HF. 211.729H ₂ O					183
$\text{H}_2\text{SiF}(\text{a2})$	144.092	-574.670	0.471	-2404.420	69 70
In 41.157 HF. 174.657H ₂ O					184
$\text{H}_2\text{SiF}_6(\text{a3})$	144.092	-574.670	0.471	-2404.420	70 83
In 179 HF. 758H ₂ O					184
$\text{H}_2\text{SiF}_6(\text{a4})$	144.092	-575.373	0.473	-2407.362	71 131
H_2SiF_6 , 1.5HF. 600H ₂ O					135 136
					137 182
$\text{SiCl}(\text{g})$	63.539	46.909	10.037	196.266	75
$\text{SiCl}_2(\text{g})$	98.992	-39.913	0.175	-166.998	84
$\text{SiCl}_3^+(\text{g})$	134.445	103.899	7.073	434.713	85 86
$\text{SiCl}_4(\text{c})$	169.898	-167.322	0.162	-700.076	76
$\text{SiCl}_4(\text{l})$	169.898	-165.477	0.161	-692.356	77 78
					79 80
					81 82
					83
$\text{SiCl}_4(\text{g})$	169.898	-158.387	0.190	-662.690	77 85
$\text{SiH}_3\text{Cl}(\text{g})$	66.563	-48.000	5.000	-200.832	87
$\text{SiH}_2\text{Cl}_2(\text{g})$	101.008	-80.000	10.000	-334.720	88
$\text{SiCl}_3\text{H}(\text{l})$	135.453	-121.723	0.319	-509.290	90
$\text{SiCl}_3\text{H}(\text{g})$	135.453	-115.223	0.303	-482.094	86 89
$\text{ClSiF}_3(\text{g})$	120.534	-315.000	5.000	-1317.960	91
$\text{FSiCl}_3(\text{g})$	153.443	-201.000	5.000	-840.984	92
$\text{SiBr}(\text{g})$	107.990	50.000	1.000	209.200	93
$\text{SiBr}_2(\text{g})$	187.894	-11.142	0.792	-46.618	97
$\text{SiBr}_4(\text{l})$	347.702	-105.844	1.280	-442.851	95 96
$\text{SiBr}_4(\text{g})$	347.702	-95.464	1.297	-399.421	94
$\text{SiHBr}_3(\text{l})$	268.806	-76.170	2.000	-318.695	99

Compound	Formula	ΔH_f°	Error	ΔH_f°	Reactions
	weight	(kcal/mol)	(kj/mol)	for ΔH_f°	
SiHBr ₃ (g)	268.806	-67.170	2.236	-281.039	98
SiI ₂ (g)	281.895	19.346	2.462	80.944	103
SiI ₄ (c)	535.704	-48.320	2.197	-202.171	100
SiI ₄ (l)	535.704	-45.320	0.909	-189.619	102
SiI ₄ (g)	535.704	-28.420	2.197	-118.909	101
SiS(g)	60.146	16.926	0.101	70.818	104
SiS ₂ (c)	92.206	-51.604	2.552	-215.912	106
SiS ₂ (l)	92.206	-46.604	2.741	-194.992	105
SiSe(g)	107.046	23.780	0.101	99.496	107
SiSe ₂ (c)	186.006	-7.000	1.000	-29.288	108
SiTe(g)	155.686	51.710	1.000	216.355	109
SiTe ₂ (g)	283.286	53.300	5.000	223.007	110
Si ₂ Te ₃ (c)	438.972	-19.750	5.000	-82.634	111
SiN(g)	42.093	90.890	10.037	380.283	112
Si ₂ N(g)	70.179	92.340	1.995	386.349	113
Si ₃ N ₄ (c)	140.285	-177.606	0.995	-743.105	114 115
Alpha					
(NH ₄) ₂ SiF ₆ (c1)	178.154	-640.940	0.101	-2681.693	116
Hexagonal					
(NH ₄) ₂ SiF ₆ (c2)	178.154	-640.670	0.143	-2680.563	117
Cubic					
(NH ₄) ₂ SiF ₆ (a)	178.154	-634.700	1.000	-2655.585	118
(NH ₄) ₂ SiF ₆ (550H ₂ O)	178.154	-634.005	1.000	-2652.677	119
(NH ₄) ₂ SiF ₆ (800H ₂ O)	178.154	-633.865	1.000	-2652.091	120
(NH ₄) ₂ SiF ₆ (1500H ₂ O)	178.154	-633.600	1.000	-2650.982	121
SiC(c1)	40.097	-17.270	0.422	-72.258	122 123
Beta, cubic					
SiC(c2)	40.097	-17.505	0.392	-73.240	124 125
Alpha, hexagonal					
SiC(g)	40.097	172.225	3.443	720.588	126
SiC ₂ (g)	52.108	147.217	2.854	615.957	127 128
Si ₂ C(g)	68.183	128.524	3.028	537.746	129 130
CH ₃ SiH ₂ ⁺ (g)	45.137	212.223	17.399	887.941	162
CH ₃ SiH ₃ (l)	46.145	-12.164	20.068	-50.894	142
CH ₃ SiH ₃ (g)	46.145	-7.774	20.068	-32.526	160
(CH ₃) ₂ SiH(g)	59.164	5.870	1.724	24.560	158
(CH ₃) ₂ SiH ⁺ (g)	59.164	184.320	10.135	771.195	164
(CH ₃) ₂ SiH ₂ (l)	60.172	-25.077	14.238	-104.922	143
(CH ₃) ₂ SiH ₂ (g)	60.172	-19.977	14.238	-83.584	161
					128 130
					215
(CH ₃) ₃ Si(g)	73.191	-9.321	1.581	-39.001	157 159
					167 170
					174
(CH ₃) ₃ Si ⁺ (g)	73.191	154.379	3.081	645.922	163 165
					167 169
(CH ₃) ₃ SiH(l)	74.199	-43.250	1.653	-180.958	144
(CH ₃) ₃ SiH(g)	74.199	-37.430	1.650	-156.607	157 163
(CH ₃) ₄ Si(l1)	88.226	-62.733	2.078	-262.476	131 145
(CH ₃) ₄ Si(g1)	88.226	-56.481	2.079	-236.314	145 165
					202
(C ₂ H ₅) ₂ SiH ₂ (l2)	88.226	-50.322	3.367	-210.547	132
(C ₂ H ₅) ₂ SiH ₂ (g2)	88.226	-43.142	3.368	-180.505	146
(CH ₃) ₂ Si(CH ₃) ₃ (l)	100.237	-40.873	3.367	-171.012	139
(CH ₃) ₂ Si(CH ₃) ₃ (g)	100.237	-32.573	3.404	-136.285	154
1 : 1-Dimethyl silacyclobutane					
(CH ₃) ₂ SiC ₂ H ₅ (l)	102.253	-66.321	10.466	-277.487	147
(CH ₃) ₃ SiC ₂ H ₅ (g)	102.253	-59.121	10.464	-247.362	166
(CH ₃) ₂ Si(CH ₃) ₄ (l)	114.264	-51.939	1.732	-217.312	140
(CH ₃) ₂ Si(CH ₃) ₄ (g)	114.264	-42.939	1.803	-179.656	155
1 : 1-Dimethyl silacyclopentane					
(C ₂ H ₅) ₃ SiH(l1)	116.280	-56.254	5.022	-235.366	133
(C ₂ H ₅) ₃ SiH(g1)	116.280	-47.554	5.031	-198.965	148

Compound	Formula weight	ΔH_f° (kcal/mol)	Error	ΔH_f° (kj/mol)	Reactions for ΔH_f°
(C ₂ H ₅) ₄ Si(l)	144.334	-72.586	5.022	-303.699	134
(C ₂ H ₅) ₄ Si(g)	144.334	-63.086	5.047	-263.951	149
(CH ₃) ₅ Si ₂ (g)	131.347	-22.534	8.474	-94.283	169 173
(CH ₃) ₅ Si ₂ ⁺ (g)	131.347	120.743	6.619	505.188	168 170
					173
Si ₂ C ₆ H ₁₆ (l)	144.366	-80.882	3.465	-338.409	141
Si ₂ C ₆ H ₁₆ (g)	144.366	-71.082	3.501	-297.406	156
1 : 1.3 : 3 - Tetramethyl, 1,3,-Disilacyclobutane					
(CH ₃) ₆ Si ₂ (l)	146.382	-94.770	2.554	-396.520	135 150
(CH ₃) ₆ Si ₂ (g)	146.382	-85.830	2.556	-359.115	150 159
					167 168
(CH ₃) ₇ Si ₃ (g)	189.503	-28.608	11.600	-119.698	172
(CH ₃) ₇ Si ₃ ⁺ (g)	189.503	94.312	8.007	394.603	171 174
(CH ₃) ₈ Si ₃ (l)	204.538	-123.564	4.455	-516.992	136 151
(CH ₃) ₈ Si ₃ (g)	204.538	-112.567	4.457	-470.980	151 169
					170 171
(CH ₃) ₉ Si ₄ (g)	247.659	-23.500	15.494	-98.325	175
tris-Trimethylsilyl silyl radical					
(CH ₃) ₁₀ Si ₄ (l)	262.694	-146.132	4.994	-611.415	137 152
(CH ₃) ₁₀ Si ₄ (g)	262.694	-133.629	5.006	-559.106	152 173
n-Decamethyl tetrasilane					
					174
(CH ₃) ₁₂ Si ₅ (c)	320.850	-151.321	10.275	-633.128	138
(CH ₃) ₁₂ Si ₅ (g)	320.850	-131.321	11.427	-549.448	153
Tetrakis-trimethylsilyl silane					
CH ₃ Si(OH) ₃ (a)	94.143	-275.355	4.904	-1152.085	176
(CH ₃) ₂ Si(OH) ₂ (a)	92.171	-206.238	2.428	-862.900	177 197
(CH ₃) ₃ SiOH(l)	90.198	-130.206	0.746	-544.780	178
(CH ₃) ₃ SiOH(g)	90.198	-119.306	0.855	-499.175	179
(CH ₃) ₃ SiOC ₂ H ₅ (l)	118.252	-126.131	0.746	-527.730	180
(C ₆ H ₅) ₂ Si(OH) ₂ (a)	216.313	-155.000	5.000	-648.520	181
(CH ₃) ₆ Si ₂ O(l)	162.381	-194.496	1.478	-813.772	182 183
(CH ₃) ₆ Si ₂ O(l)	162.381	-194.496	1.478	-813.772	182 183
					184 204
					205 219
					220
(CH ₃) ₈ Si ₂ O(g)	162.381	-185.596	1.536	-776.534	188
Hexamethyl disiloxane					
(CH ₃) ₅ Si ₃ O ₂ (l)	236.537	-338.072	3.622	-1414.491	185
(CH ₃) ₅ Si ₃ O ₂ (g)	236.537	-328.572	3.657	-1374.743	189
Octamethyl trisiloxane					
(CH ₃) ₁₀ Si ₄ O ₃ (l)	310.692	-471.946	10.177	-1974.624	186
(CH ₃) ₁₀ Si ₄ O ₃ (g)	310.692	-460.446	10.189	-1926.508	190
Decamethyl tetrasiloxane					
(CH ₃) ₁₂ Si ₅ O ₄ (l)	384.848	-624.021	5.530	-2610.905	187
(CH ₃) ₁₂ Si ₅ O ₄ (g)	384.848	-611.321	5.553	-2557.768	191
Dodecamethylpentasiloxane					
CH ₃ SiCl ₃ (l)	149.480	-138.709	4.876	-580.358	193 201
CH ₃ SiCl ₃ (g)	149.480	-131.206	4.874	-548.964	201 203
CH ₃ HSiCl ₂ (l)	115.035	-105.618	5.008	-441.907	196
(CH ₃) ₂ SiCl ⁺ (g)	93.609	126.656	10.029	529.928	207
(CH ₃) ₂ HSiCl(l)	94.617	-79.595	5.008	-333.026	195
(CH ₃) ₂ SiCl ₂ (l)	129.062	-117.049	2.425	-489.734	194 197
					200
(CH ₃) ₂ SiCl ₂ (g)	129.062	-109.511	2.421	-458.193	200 202
					203
(CH ₃) ₃ SiCl(l)	108.644	-91.747	0.741	-383.870	192 199
					204 205
(CH ₃) ₃ SiCl(g)	108.644	-84.544	0.764	-353.733	199 202
					203
(C ₆ H ₅) ₂ SiCl ₂ (l)	253.204	-66.316	5.028	-277.466	198
(CH ₃) ₂ SiBr ⁺ (g)	138.060	143.301	10.030	599.573	210
(CH ₃) ₃ SiBr(l)	153.095	-78.099	0.746	-326.764	208

Compound	Formula	ΔH_f°	Error	ΔH_f°	Reactions
	weight	(kcal/mol)		(kj/mol)	for ΔH_f°
(CH ₃) ₃ SiBr(g)	153.095	-70.299	0.774	-294.129	209
(CH ₃) ₂ SiI ⁺ (g)	185.060	154.846	14.474	647.876	212
(CH ₃) ₃ SiI(g)	200.095	-49.454	10.464	-206.916	211
(CH ₃) ₃ SiSC ₄ H ₉ (l) N-butylthio-trimethylsilane	162.367	-91.011	0.746	-380.788	213
(CH ₃) ₃ SiSC ₄ H ₉ (g)	162.367	-81.311	0.753	-340.203	214
SiCN(g)	54.104	90.582	2.344	378.994	215 216
(CH ₃) ₃ SiNHCH ₃ (l)	103.241	-63.939	0.746	-267.519	217
(CH ₃) ₃ SiNHCH ₃ (g)	103.241	-55.339	0.753	-231.537	223
(CH ₃) ₃ SiN(CH ₃) ₂ (l)	117.268	-66.856	0.746	-279.724	218
(CH ₃) ₃ SiN(CH ₃) ₂ (g)	117.268	-59.256	0.753	-247.925	224
(CH ₃) ₃ SiN(C ₂ H ₅) ₂ (l)	145.322	-87.647	0.800	-366.715	206
(CH ₃) ₆ Si ₂ NH(l)	161.397	-123.573	1.479	-517.029	219 220
(CH ₃) ₆ Si ₂ NH(g)	161.397	-113.673	1.482	-475.608	225
(CH ₃) ₆ Si ₂ NCH ₃ (l)	175.424	-117.429	1.492	-491.324	221
(CH ₃) ₆ Si ₂ NCH ₃ (g)	175.424	-108.129	1.496	-452.412	226
(CH ₃) ₉ Si ₃ N(c)	233.580	-172.317	2.286	-720.973	222
(CH ₃) ₉ Si ₃ N(g)	233.580	-159.317	2.288	-666.581	227

Compounds from other sections

Compound	Formula	ΔH_f°	Error	ΔH_f°
	weight	(kcal/mol)		(kj/mol)
O ₂ (gs)	31.999	0.000	0.000	0.000
H(g)	1.008	52.103	0.000	217.999
H ₂ (gs)	2.016	0.000	0.000	0.000
H ₂ O(l)	18.015	-68.315	0.000	-285.830
H ₂ O(g)	18.015	-57.795	0.000	-241.814
F(g)	18.998	18.880	1.000	78.994
F ⁻ (g)	18.998	-64.700	0.500	-270.705
F ₂ (gs)	37.997	0.000	0.000	0.000
HF(g)	20.006	-65.320	0.072	-273.297
HF(4.0H ₂ O)	20.006	-76.831	0.070	-321.461
HF(25H ₂ O)	20.006	-77.015	0.070	-322.230
HF(30H ₂ O)	20.006	-77.021	0.070	-322.256
HF(80H ₂ O)	20.006	-77.046	0.070	-322.360
Cl(g)	35.453	28.989	0.001	121.290
Cl ₂ (gs)	70.906	0.000	0.000	0.000
HCl(a)	36.461	-39.933	0.001	-167.080
HCl(10H ₂ O)	36.461	-38.537	0.001	-161.239
HCl(50H ₂ O)	36.461	-39.502	0.001	-165.276
HCl(100H ₂ O)	36.461	-39.638	0.001	-165.845
HCl(500H ₂ O)	36.461	-39.793	0.001	-166.494
HCl(600H ₂ O)	36.461	-39.804	0.001	-166.540
HCl(800H ₂ O)	36.461	-39.820	0.001	-166.607
HCl(1500H ₂ O)	36.461	-39.848	0.001	-166.724
Br ₂ (ls)	159.808	0.000	0.000	0.000
HBr(50H ₂ O)	80.912	-28.708	0.001	-120.114
I(g)	126.905	25.517	0.003	106.763
S(cs)	32.060	0.000	0.000	0.000
H ₂ S(g)	34.076	-4.930	0.000	-20.627
Se(cs)	78.960	0.000	0.000	0.000
Te ₂ (g)	255.200	40.200	0.000	168.197
N(g)	14.007	112.970	0.003	472.666
N ₂ (gs)	28.013	0.000	0.000	0.000
NH ₄ ⁺ (a)	18.039	-31.850	0.001	-133.260
NH ₄ Cl(50H ₂ O)	53.492	-71.647	0.002	-299.771
NH ₄ Cl(1500H ₂ O)	53.492	-71.720	0.002	-300.076

Compound	Formula	ΔH_f°	Error	ΔH_f°
	weight	(kcal/mol)		(kJ/mol)
C(cs)	12.011	0.000	0.000	0.000
CO ₂ (g)	44.010	-94.051	0.000	-393.509
CH ₃ (g)	15.035	33.200	0.000	138.909
C ₂ H ₅ (g)	29.062	25.000	0.000	104.600
C ₂ H ₅ OH(50H ₂ O)	46.069	-68.740	0.000	-287.608
CF ₄ (g)	88.005	-223.040	0.000	-933.199
C ₄ H ₉ SH(l)	90.184	-29.720	0.000	-124.348
(C ₂ H ₅) ₂ NH(l)	73.139	-24.780	0.000	-103.680
CH ₃ NH ₃ Cl(50H ₂ O)	67.519	-69.650	0.000	-291.416
(CH ₃) ₂ NH ₂ Cl(50H ₂ O)	81.546	-68.567	0.000	-286.884
(C ₂ H ₅) ₂ NH ₂ Cl(c)	109.600	-85.760	0.000	-358.820
Ga(l)	69.720	1.330	0.000	5.565
Ga ₂ O(g)	155.439	-21.000	0.000	-87.864
Ag(cs)	107.868	0.000	0.000	0.000
AgCl(c)	143.321	-30.370	0.000	-127.068
AgBr(c)	187.772	-23.990	0.000	-100.374
AgI(c)	234.772	-14.780	0.000	-61.840
Ca(g)	40.080	42.600	0.000	178.238
CaF(g)	59.078	-65.000	0.000	-271.960
CaF ₂ (c)	78.077	-291.500	0.000	-1219.636
CaF ₂ (g)	78.077	-186.800	0.000	-781.571
Na(cs)	22.990	0.000	0.000	0.000
NaOH(c)	39.997	-101.990	0.000	-426.726
NaF(c)	41.988	-137.910	0.000	-577.015
NaCl(c)	58.443	-98.230	0.000	-410.994
NaBr(c)	102.894	-86.030	0.000	-359.950
Na ₂ SiO ₃ (c)	122.064	-363.000	0.000	-1518.792

Reactions

	Reaction	ΔH° (kcal/mol)	Error	Ref
1	$\text{Si}(\text{cs})=\text{Si}(\text{l})$	11.585	0.100	71/1
2	$\text{Si}(\text{cs})=\text{Si}(\text{am})$	1.000	0.100	NBS
3	$\text{Si}(\text{cs})=\text{Si}(\text{g})$	107.700	1.000	71/1
	Residual	-0.220		
4	$\text{Si}(\text{g})=\text{Si}^+(\text{g})$	189.450	0.100	49/1
5	$\text{Si}_2(\text{g})=2\text{Si}(\text{g})$	74.000	3.000	71/1
6	$\text{Si}_3(\text{g})=3\text{Si}(\text{g})$	171.000	10.000	71/1
7	$\text{SiO}_2(\text{c1})+\text{H}_2(\text{gs})=\text{SiO}(\text{g})+\text{H}_2\text{O}(\text{g})$	135.200	5.000	63/2
	Residual	-0.227		
8	$\text{Si}(\text{cs})+\text{SiO}_2(\text{c1})=2\text{SiO}(\text{g})$	168.800	5.000	63/2
	Residual	0.353		
9	$2\text{Ga}(\text{l})+\text{SiO}_2(\text{am})=\text{SiO}(\text{g})+\text{Ga}_2\text{O}(\text{g})$	170.800	5.000	62/1
	Residual	3.491		
10	$\text{Si}(\text{cs})+\text{SiO}_2(\text{c2})=2\text{SiO}(\text{g})$	165.960	5.000	62/1
	Residual	-1.984		
11	$\text{SiO}(\text{g})=\text{SiO}^+(\text{g})$	249.000	12.000	55/1
12	$\text{SiO}_2(\text{am})=\text{SiO}_2(\text{c1})$	-2.210	0.100	12/1
	Residual	0.043		
		-2.330	0.100	21/1
	Residual	-0.077		
		0.500	0.200	62/1
13	$\text{SiO}_2(\text{c1})=\text{SiO}_2(\text{c2})$	Residual	-0.003	
		0.600	0.100	71/1
14	$\text{SiO}_2(\text{c1})=\text{SiO}_2(\text{c3})$	1.950	0.300	71/1
15	$\text{SiO}_2(\text{c1})=\text{SiO}_2(\text{l})$	143.000	8.000	71/1
16	$\text{SiO}_2(\text{c3})=\text{SiO}_2(\text{g})$	1.500	0.100	31/1
17	$\text{SiO}_2(\text{a})=\text{SiO}_2(\text{am})$	270.000	12.000	55/1
18	$\text{SiO}_2(\text{g})=\text{SiO}_2^+(\text{g})$	70.000	3.000	71/1
19	$\text{SiH}(\text{g})=\text{Si}(\text{g})+\text{H}(\text{g})$	-7.300	0.300	61/1
20	$\text{SiH}_4(\text{g})=\text{Si}(\text{am})+2\text{H}_2(\text{gs})$	Residual	0.002	
		279.000	9.000	63/3
21	$\text{SiH}_4(\text{g})=\text{SiH}_2^+(\text{g})+\text{H}_2(\text{gs})$	Residual	2.011	
		274.400	9.000	69/3
	Residual	-2.589		
		274.700	9.000	62/2
	Residual	-2.289		
		272.100	9.000	63/3
22	$\text{SiH}_4(\text{g})=\text{SiH}_3^+(\text{g})+\text{H}(\text{g})$	Residual	-6.450	
		272.400	9.000	66/2
	Residual	-6.150		
		283.700	9.000	69/3
	Residual	5.150		
		286.000	9.000	62/2
	Residual	7.450		
23	$\text{Si}_2\text{H}_6(\text{g})=2\text{Si}(\text{am})+3\text{H}_2(\text{gs})$	-17.100	0.300	61/1
24	$\text{Si}_2\text{H}_6(\text{l})=\text{Si}_2\text{H}_6(\text{g})$	4.600	0.500	33/1
25	$\text{Si}_2\text{H}_6(\text{g})=\text{SiH}_2^+(\text{g})+\text{SiH}_4(\text{g})$	275.300	9.000	62/5
	Residual	0.806		
		275.600	9.000	69/3
26	$\text{Si}_2\text{H}_6(\text{g})=\text{SiH}_3^+(\text{g})+\text{SiH}_3(\text{g})$	Residual	1.106	
		273.300	9.000	62/5
	Residual	3.400		
		275.600	9.000	69/3
	Residual	5.700		
		260.800	9.000	66/2
27	$\text{Si}_2\text{H}_6(\text{g})=\text{Si}_2\text{H}_5^+(\text{g})+\text{H}(\text{g})$	Residual	-9.100	
		262.900	9.000	69/3

Reaction	ΔH° (kcal/mol)	Error (kcal/mol)	Ref
28 $\text{Si}_3\text{H}_8(\text{g}) = 3\text{Si}(\text{am}) + 4\text{H}_2(\text{gs})$	-25.900	1.000	64/3
29 $\text{Si}_3\text{H}_8(\text{l}) = \text{Si}_3\text{H}_8(\text{g})$	6.700	0.500	33/1
30 $2\text{H}_2(\text{gs}) + 3\text{O}_2(\text{gs}) + 2\text{Si}(\text{cs}) = 2\text{H}_2\text{SiO}_3(\text{c})$	-568.200	2.000	NBS
31 $\text{H}_2\text{SiO}_3(\text{c}) = \text{H}_2\text{SiO}_3(\text{ao})$	1.400	0.100	NBS
32 $2\text{H}_2(\text{gs}) + 2\text{O}_2(\text{gs}) + \text{Si}(\text{cs}) = \text{H}_4\text{SiO}_4(\text{c})$	-354.000	1.000	NBS
33 $\text{H}_4\text{SiO}_4(\text{c}) = \text{H}_4\text{SiO}_4(\text{ao})$	3.000	0.100	NBS
34 $2\text{H}_2(\text{gs}) + 5\text{O}_2(\text{gs}) + 4\text{Si}(\text{cs}) = 2\text{H}_2\text{Si}_2\text{O}_5(\text{c})$	-998.400	2.000	NBS
35 $6\text{H}_2(\text{gs}) + 7\text{O}_2(\text{gs}) + 4\text{Si}(\text{cs}) = 2\text{H}_6\text{Si}_2\text{O}_7(\text{c})$	-1276.000	2.000	NBS
36 $\text{SiF}(\text{g}) = \text{Si}(\text{g}) + \text{F}(\text{g})$	125.000	10.000	58/1
	Residual	-6.308	
37 $\text{Si}(\text{g}) + \text{SiF}_2(\text{g}) = 2\text{SiF}(\text{g})$	23.500	3.000	71/1
	Residual	-0.350	
38 $\text{SiF}_2(\text{c}) + \text{F}_2(\text{gs}) = \text{SiF}_4(\text{g})$	-199.700	1.700	70/3
39 $\text{CaF}_2(\text{c}) + \text{Si}(\text{g}) = \text{Ca}(\text{g}) + \text{SiF}_2(\text{g})$	86.700	1.500	64/9
	Residual	1.306	
40 $\text{CaF}_2(\text{g}) + \text{Si}(\text{g}) = \text{Ca}(\text{g}) + \text{SiF}_2(\text{g})$	-16.700	1.500	64/9
	Residual	2.606	
41 $2\text{CaF}(\text{g}) + \text{Si}(\text{g}) = 2\text{Ca}(\text{g}) + \text{SiF}_2(\text{g})$	-34.700	0.900	64/9
	Residual	-1.194	
42 $\text{SiF}_3(\text{g}) = \text{SiF}_3^-(\text{g})$	-77.000	2.000	70/2
43 $\text{Si}(\text{cs}) + 2\text{F}_2(\text{gs}) = \text{SiF}_4(\text{g})$	-385.980	0.190	63/2
	Residual	0.009	
44 $\text{SiF}_4(\text{g}) + 2\text{H}_2\text{O}(\text{g}) = \text{SiO}_2(\text{c3}) + 4\text{HF}(\text{g})$	24.530	5.000	63/1
	Residual	1.627	
45 $\text{SiF}_4(\text{g}) + 2\text{H}_2\text{O}(\text{g}) = \text{SiO}_2(\text{am}) + 4\text{HF}(\text{g})$	26.370	5.000	63/1
	Residual	1.814	
46 $\text{SiO}_2(\text{am}) + 2\text{F}_2(\text{gs}) = \text{SiF}_4(\text{g}) + \text{O}_2(\text{gs})$	-170.040	0.250	63/2
	Residual	0.204	
47 $\text{SiO}_2(\text{c1}) + 2\text{F}_2(\text{gs}) = \text{SiF}_4(\text{g}) + \text{O}_2(\text{gs})$	-168.260	0.280	63/2
	Residual	-0.269	
48 $\text{SiF}_4(\text{g}) + 4\text{Na}(\text{cs}) = 4\text{NaF}(\text{c}) + \text{Si}(\text{cs})$	-174.100	10.000	60/2
	Residual	-8.449	
49 $\text{Si}(\text{cs}) + 3\text{F}_2(\text{gs}) = \text{SiF}_6^{--}(\text{a})$	-571.000	1.000	NBS
50 $2\text{Si}(\text{cs}) + 3\text{F}_2(\text{gs}) = \text{Si}_2\text{F}_6(\text{g})$	-565.000	10.000	64/6
	Residual	0.558	
51 $\text{Si}_2\text{F}_6(\text{c}) = \text{Si}_2\text{F}_6(\text{g})$	10.400	0.500	65/1
52 $\text{Si}_3\text{F}_8(\text{c}) = \text{Si}_3\text{F}_8(\text{g})$	14.100	0.500	65/1
53 $\text{Si}_2\text{F}_6(\text{c}) = \text{Si}_2\text{F}_6(\text{l})$	3.900	0.800	65/1
54 $\text{Si}_3\text{F}_8(\text{c}) = \text{Si}_3\text{F}_8(\text{l})$	5.300	0.800	65/1
55 $\text{SiF}(\text{g}) = \text{SiF}^+(\text{g})$	168.400	5.000	68/1
56 $\text{SiF}_2(\text{g}) = \text{SiF}_2^+(\text{g})$	260.400	6.900	68/1
	Residual	-0.558	
57 $\text{SiF}_3(\text{g}) = \text{SiF}_3^+(\text{g})$	196.000	20.000	68/1
58 $\text{SiF}_4(\text{g}) = \text{SiF}_4^+(\text{g}) + \text{F}(\text{g})$	373.600	6.900	68/1
59 $\text{SiF}_4(\text{g}) = \text{SiF}_4^+(\text{g})$	362.300	6.900	68/1
60 $\text{Si}_2\text{F}_6(\text{g}) = \text{SiF}_2^+(\text{g}) + \text{SiF}_4(\text{g})$	300.300	6.900	68/1
	Residual	0.558	
61 $\text{Si}_2\text{F}_6(\text{g}) = \text{Si}_2\text{F}_5^+(\text{g}) + \text{F}^-(\text{g})$	297.300	6.900	68/1
62 $\text{Si}_3\text{F}_8(\text{g}) = \text{SiF}_2^+(\text{g}) + \text{Si}_2\text{F}_6(\text{g})$	310.200	6.900	68/1
63 $\text{Si}_3\text{F}_8(\text{g}) = \text{SiF}_3^+(\text{g}) + \text{Si}_2\text{F}_5(\text{g})$	357.700	6.900	68/1
64 $\text{Si}_3\text{F}_8(\text{g}) = \text{Si}_2\text{F}_4^+(\text{g}) + \text{SiF}_4(\text{g})$	263.600	6.900	68/1
65 $\text{Si}_3\text{F}_8(\text{g}) = \text{Si}_3\text{F}_7^+(\text{g}) + \text{F}(\text{g})$	360.200	6.900	68/1
66 $\text{Si}_3\text{F}_8(\text{g}) = \text{Si}_3\text{F}_8^-(\text{g})$	250.000	6.900	68/1
67 $\text{SiO}_2(\text{c1}) + 6\text{HF}(4.0\text{H}_2\text{O})$ $= \text{H}_2\text{SiF}_6(20\text{H}_2\text{O}) + 2\text{H}_2\text{O}(\text{l})$	-33.290	0.080	51/1
	Residual	-0.113	
	-33.000	0.100	48/1
	Residual	0.177	
68 $\text{Si}(\text{cs}) + \text{O}_2(\text{gs}) + 6\text{HF}(30\text{H}_2\text{O})$ $= \text{H}_2\text{SiF}_6(\text{a1}) + 2\text{H}_2\text{O}(\text{l})$	-249.890	0.250	64/2
	Residual	-0.012	
69 $\text{Si}(\text{cs}) + \text{O}_2(\text{gs}) + 6\text{HF}(4.0\text{H}_2\text{O})$ $= \text{H}_2\text{SiF}_6(\text{a2}) + 2\text{H}_2\text{O}(\text{l})$	-250.300	0.250	64/2
	Residual	0.015	

	Reaction	ΔH° (kcal/mol)	Error (kcal/mol)	Ref
70	$H_2SiF_6(a3)=H_2SiF_6(a2)$	0.000	0.000	64/4
71	$H_2SiF_6(a1)=H_2SiF_6(a4)$	0.000	0.000	72/1
72	$2Si(cs)+3H_2(gs)+F_2(gs)=2SiH_3F(g)$	-210.000	30.000	71/1
73	$Si(cs)+H_2(gs)+F_2(gs)=H_2SiF_2(g)$	-194.000	15.000	71/1
74	$2Si(cs)+H_2(gs)+3F_2(gs)=2SiHF_3(g)$	-566.000	30.000	71/1
75	$SiCl(g)=Si(g)+Cl(g)$	90.000	15.000	64/8
76	$SiCl_4(c)=SiCl_4(l)$	1.845	0.020	22/1
77	$SiCl_4(l)=SiCl_4(g)$	7.090	0.100	71/1
78	$Si(cs)+2Cl_2(gs)=SiCl_4(l)$	-165.490	0.160	69/1
		Residual	-0.013	
79	$SiCl_4(l)+4Ag(cs)=Si(cs)+4AgCl(c)$	42.700	2.000	64/5
		Residual	-1.297	
80	$SiCl_4(l)+6NaOH(c)=Na_2SiO_3(c)+4NaCl(c)+3H_2O(l)$	-192.310	9.000	61/2
		Residual	-8.862	
81	$SiCl_4(l)+2H_2O(l)=SiO_2(a)+4HCl(100H_2O)$	-79.750	5.000	66/1
		Residual	-6.061	
82	$SiCl_4(l)+2H_2O(l)=SiO_2(a)+4HCl(500H_2O)$	-76.600	5.000	71/1
		Residual	-2.291	
83	$SiCl_4(l)+6HF(4.0H_2O)=H_2SiF_6(a3)+4HCl(10H_2O)$	-98.610	5.000	64/4
		Residual	3.745	
84	$SiCl_4(g)+Si(cs)=2SiCl_2(g)$	78.350	0.500	71/1
		Residual	-0.210	
		Residual	78.490	0.500
		Residual	-0.070	71/1
		Residual	78.840	0.500
		Residual	0.280	
85	$SiCl_4(g)=SiCl_3^+(g)+Cl(g)$	287.800	9.000	62/2
		Residual	-3.475	
86	$SiCl_3H(g)=SiCl_3^+(g)+H(g)$	274.700	9.000	62/2
		Residual	3.475	
87	$3H_2(gs)+Cl_2(gs)+2Si(cs)=2SiH_3Cl(g)$	-96.000	30.000	71/1
88	$H_2(gs)+Cl_2(gs)+Si(cs)=SiH_2Cl_2(g)$	-80.000	15.000	71/1
89	$SiCl_3H(g)+5NaOH(c)=Na_2SiO_3(c)+3NaCl(c)+2H_2O(l)+H_2(gs)$	-169.150	0.300	61/2
		Residual	-0.003	
90	$SiCl_3H(l)=SiCl_3H(g)$	6.500	0.100	61/2
91	$Cl_2(gs)+3F_2(gs)+2Si(cs)=2ClSiF_3(g)$	-630.000	30.000	71/1
92	$3Cl_2(gs)+F_2(gs)+2Si(cs)=2FSiCl_3(g)$	-402.000	30.000	71/1
93	$2Si(cs)+Br_2(l)=2SiBr(g)$	100.000	2.000	NBS
94	$SiBr_4(l)=SiBr_4(g)$	10.380	0.200	47/1
95	$SiBr_4(l)+4Ag(cs)=Si(cs)+4AgBr(c)$	13.300	1.560	64/5
		Residual	3.416	
96	$SiBr_4(l)+6NaOH(c)=Na_2SiO_3(c)+4NaBr(c)+3H_2O(l)$	-199.200	2.000	62/4
		Residual	-4.919	
97	$Si(cs)+SiBr_4(g)=2SiBr_2(g)$	71.500	1.000	67/4
		Residual	-1.680	
		Residual	79.900	2.000
		Residual	6.720	66/3
		Residual	9.000	1.000
98	$SiHBr_3(l)=SiHBr_3(g)$			62/4
99	$SiHBr_3(l)+5NaOH(c)=Na_2SiO_3(c)+3NaBr(c)+2H_2O(l)+H_2(gs)$	-171.600	2.000	62/4
100	$SiI_4(c)=SiI_4(l)$	3.000	2.000	
	Estimated value			
101	$SiI_4(l)=SiI_4(g)$	16.900	2.000	
	Estimated value			
102	$SiI_4(l)+4Ag(cs)=Si(cs)+4AgI(c)$	-13.800	0.850	64/5
103	$SiI_4(g)=SiI_2(g)+2I(g)$	98.800	1.000	67/4
104	$Si(cs)+S(cs)=SiS(g)$	16.926	0.100	71/1
105	$SiS_2(c)=SiS_2(l)$	5.000	1.000	71/1
106	$SiS_2(c)+6HF(4.0H_2O)=H_2SiF_6(20H_2O)+2H_2S(g)$	-72.800	2.200	54/1

	Reaction	ΔH° (kcal/mol)	Error	Ref
107	$\text{Si}(\text{cs}) + \text{Se}(\text{cs}) = \text{SiSe}(\text{g})$	23.780	0.100	NBS
108	$\text{Si}(\text{cs}) + 2\text{Se}(\text{cs}) = \text{SiSe}_2(\text{c})$	-7.000	1.000	NBS
109	$2\text{Si}(\text{cs}) + \text{Te}_2(\text{g}) = 2\text{SiTe}(\text{g})$	61.600	6.000	67/1
	Residual	-1.620		
		63.400	2.000	67/1
	Residual	0.180		
110	$\text{Si}(\text{cs}) + \text{Te}_2(\text{g}) = \text{SiTe}_2(\text{g})$	13.100	4.000	67/1
111	$2\text{Si}_2\text{Te}_3(\text{c}) = 4\text{Si}(\text{cs}) + 3\text{Te}_2(\text{g})$	160.600	8.000	67/1
	Residual	0.500		
		159.600	8.000	67/1
	Residual	-0.500		
112	$\text{SiN}(\text{g}) = \text{Si}(\text{g}) + \text{N}(\text{g})$	130.000	15.000	71/1
113	$2\text{Si}_2\text{N}(\text{g}) = 4\text{Si}(\text{g}) + \text{N}_2(\text{gs})$	247.000	2.000	67/2
114	$\text{Si}_3\text{N}_4(\text{c}) = 3\text{Si}(\text{cs}) + 2\text{N}_2(\text{gs})$	177.600	1.000	71/1
	Residual	-0.006		
115	$\text{Si}_3\text{N}_4(\text{c}) = 3\text{Si}(\text{l}) + 2\text{N}_2(\text{gs})$	213.000	7.000	71/1
	Residual	0.637		
116	$\text{N}_2(\text{gs}) + 4\text{H}_2(\text{gs}) + \text{Si}(\text{cs}) + 3\text{F}_2(\text{gs})$ $= (\text{NH}_4)_2\text{SiF}_6(\text{c1})$	-640.940	0.100	NBS
117	$(\text{NH}_4)_2\text{SiF}_6(\text{c1}) = (\text{NH}_4)_2\text{SiF}_6(\text{c2})$	0.270	0.100	NBS
118	$(\text{NH}_4)_2\text{SiF}_6(\text{a}) = 2\text{NH}_4^+(\text{a}) + \text{SiF}_6^{-(\text{a})}$	0.000	0.000	
119	$(\text{NH}_4)_2\text{SiF}_6(550\text{H}_2\text{O}) = (\text{NH}_4)_2\text{SiF}_6(\text{a})$	-0.695	0.001	NBS
120	$(\text{NH}_4)_2\text{SiF}_6(800\text{H}_2\text{O}) = (\text{NH}_4)_2\text{SiF}_6(\text{a})$	-0.835	0.001	NBS
121	$(\text{NH}_4)_2\text{SiF}_6(1500\text{H}_2\text{O}) = (\text{NH}_4)_2\text{SiF}_6(\text{a})$	-1.100	0.001	NBS
122	$\text{SiC}(\text{c1}) + 4\text{F}_2(\text{gs}) = \text{SiF}_4(\text{g}) + \text{CF}_4(\text{g})$	-591.790	0.380	70/1
	Residual	-0.032		
123	$\text{SiC}(\text{c1}) = \text{C}(\text{cs}) + \text{Si}(\text{g})$	125.000	3.000	58/2
	Residual	-0.190		
124	$\text{SiC}(\text{c2}) + 4\text{F}_2(\text{gs}) = \text{SiF}_4(\text{g}) + \text{CF}_4(\text{g})$	-591.530	0.340	70/1
	Residual	-0.006		
125	$\text{SiC}(\text{c2}) = \text{C}(\text{cs}) + \text{Si}(\text{g})$	126.000	3.000	61/3
	Residual	0.575		
126	$\text{C}(\text{cs}) + \text{Si}(\text{g}) = \text{SiC}(\text{g})$	65.940	4.000	71/1
	Residual	1.635		
		62.670	4.000	71/1
	Residual	-1.635		
127	$2\text{C}(\text{cs}) + \text{Si}(\text{g}) = \text{SiC}_2(\text{g})$	39.000	4.000	71/1
	Residual	-0.297		
		38.300	4.000	71/1
	Residual	-0.997		
128	$2\text{SiC}(\text{c1}) = \text{Si}(\text{l}) + \text{SiC}_2(\text{g})$	194.800	4.000	71/1
	Residual	1.457		
129	$\text{C}(\text{cs}) + 2\text{Si}(\text{g}) = \text{Si}_2\text{C}(\text{g})$	-88.000	4.000	71/1
	Residual	-0.685		
		-86.800	4.000	71/1
	Residual	0.515		
130	$\text{SiC}(\text{c1}) + \text{Si}(\text{l}) = \text{Si}_2\text{C}(\text{g})$	134.400	4.000	71/1
	Residual	0.191		
131	$(\text{CH}_3)_4\text{Si}(\text{II}) + 8\text{O}_2(\text{gs}) + 6\text{HF}(80\text{H}_2\text{O})$ $= \text{H}_2\text{SiF}_6(\text{a4}) + 4\text{CO}_2(\text{g}) + 8\text{H}_2\text{O}(\text{l})$	-971.400	2.400	72/1
	Residual	1.689		
132	$(\text{C}_2\text{H}_5)_2\text{SiH}_2(\text{l2}) + 8\text{O}_2(\text{gs}) + 6\text{HF}(80\text{H}_2\text{O})$ $= \text{H}_2\text{SiF}_6(\text{a4}) + 4\text{CO}_2(\text{g}) + 8\text{H}_2\text{O}(\text{l})$	-985.500	3.200	72/1
133	$(\text{C}_2\text{H}_5)_3\text{SiH}(\text{l1}) + 11\text{O}_2(\text{gs}) + 6\text{HF}(80\text{H}_2\text{O})$ $= \text{H}_2\text{SiF}_6(\text{a4}) + 6\text{CO}_2(\text{g}) + 10\text{H}_2\text{O}(\text{l})$	-1304.300	3.700	72/1
134	$(\text{C}_2\text{H}_5)_4\text{Si}(\text{l}) + 14\text{O}_2(\text{gs}) + 6\text{HF}(80\text{H}_2\text{O})$ $= \text{H}_2\text{SiF}_6(\text{a4}) + 8\text{CO}_2(\text{g}) + 12\text{H}_2\text{O}(\text{l})$	-1612.700	3.500	72/1
135	$2(\text{CH}_3)_6\text{Si}_2(\text{l}) + 25\text{O}_2(\text{gs}) + 24\text{HF}(80\text{H}_2\text{O})$ $= 4\text{H}_2\text{SiF}_6(\text{a4}) + 12\text{CO}_2(\text{g}) + 26\text{H}_2\text{O}(\text{l})$	-3167.600	3.600	72/1
	Residual	0.052		
136	$(\text{CH}_3)_8\text{Si}_3(\text{l}) + 17\text{O}_2(\text{gs}) + 18\text{HF}(80\text{H}_2\text{O})$ $= 3\text{H}_2\text{SiF}_6(\text{a4}) + 8\text{CO}_2(\text{g}) + 18\text{H}_2\text{O}(\text{l})$	-2199.500	3.900	72/1
	Residual	-1.693		

	Reaction	ΔH° (kcal/mol)	Error (kcal/mol)	Ref
137	$2(\text{CH}_3)_{10}\text{Si}_4(\text{l}) + 43\text{O}_2(\text{gs}) + 48\text{HF}(80\text{H}_2\text{O}) = 8\text{H}_2\text{SiF}_6(\text{a4}) + 20\text{CO}_2(\text{g}) + 46\text{H}_2\text{O}(\text{l})$	-5635.400 Residual	10.600 0.629	72/1
138	$(\text{CH}_3)_{12}\text{Si}_5(\text{c}) + 26\text{O}_2(\text{gs}) + 30\text{HF}(80\text{H}_2\text{O}) = 5\text{H}_2\text{SiF}_6(\text{a4}) + 12\text{CO}_2(\text{g}) + 28\text{H}_2\text{O}(\text{l})$	-3455.600	7.400	72/1
139	$(\text{CH}_3)_2\text{Si}(\text{CH}_2)_3(\text{l}) + 9\text{O}_2(\text{gs}) + 6\text{HF}(80\text{H}_2\text{O}) = \text{H}_2\text{SiF}_6(\text{a4}) + 5\text{CO}_2(\text{g}) + 8\text{H}_2\text{O}(\text{l})$	-1089.000	2.600	72/1
140	$2(\text{CH}_3)_2\text{Si}(\text{CH}_2)_4(\text{l}) + 21\text{O}_2(\text{gs}) + 12\text{HF}(80\text{H}_2\text{O}) = 2\text{H}_2\text{SiF}_6(\text{a4}) + 12\text{CO}_2(\text{g}) + 18\text{H}_2\text{O}(\text{l})$	-2480.600	3.000	72/1
141	$\text{Si}_2\text{C}_6\text{H}_{16}(\text{l}) + 12\text{O}_2(\text{gs}) + 12\text{HF}(80\text{H}_2\text{O}) = 2\text{H}_2\text{SiF}_6(\text{a4}) + 6\text{CO}_2(\text{g}) + 12\text{H}_2\text{O}(\text{l})$	-1529.400	3.000	72/1
142	$\text{CH}_3\text{SiH}_3(\text{l}) = \text{CH}_3\text{SiH}_3(\text{g})$	4.390	0.100	53/1
143	$(\text{CH}_3)_2\text{SiH}_2(\text{l}) = (\text{CH}_3)_2\text{SiH}_2(\text{g})$	5.100	0.100	53/1
144	$(\text{CH}_3)_3\text{SiH}(\text{l}) = (\text{CH}_3)_3\text{SiH}(\text{g})$	5.820	0.100	53/1
145	$(\text{CH}_3)_4\text{Si}(\text{l1}) = (\text{CH}_3)_4\text{Si}(\text{g1})$	6.250	0.100	53/1
		Residual	-0.003	
146	$(\text{C}_2\text{H}_5)_2\text{SiH}_2(\text{l2}) = (\text{C}_2\text{H}_5)_2\text{SiH}_2(\text{g2})$	7.180	0.100	53/1
147	$(\text{CH}_3)_3\text{SiC}_2\text{H}_5(\text{l}) = (\text{CH}_3)_3\text{SiC}_2\text{H}_5(\text{g})$	7.200	0.200	46/1
148	$(\text{C}_2\text{H}_5)_3\text{SiH}(\text{l1}) = (\text{C}_2\text{H}_5)_3\text{SiH}(\text{g1})$	8.700	0.300	72/1
149	$(\text{C}_2\text{H}_5)_4\text{Si}(\text{l}) = (\text{C}_2\text{H}_5)_4\text{Si}(\text{g})$	9.500	0.500	46/1
150	$(\text{CH}_3)_6\text{Si}_2(\text{l}) = (\text{CH}_3)_6\text{Si}_2(\text{g})$	8.940	0.100	59/1
		11.000	0.200	72/3
151	$(\text{CH}_3)_8\text{Si}_3(\text{l}) = (\text{CH}_3)_8\text{Si}_3(\text{g})$	Residual	0.003	
152	$(\text{CH}_3)_{10}\text{Si}_4(\text{l}) = (\text{CH}_3)_{10}\text{Si}_4(\text{g})$	12.500	0.400	72/3
		Residual	-0.002	
153	$(\text{CH}_3)_{12}\text{Si}_5(\text{c}) = (\text{CH}_3)_{12}\text{Si}_5(\text{g})$	20.000	5.000	72/3
154	$(\text{CH}_3)_2\text{Si}(\text{CH}_2)_3(\text{l}) = (\text{CH}_3)_2\text{Si}(\text{CH}_2)_3(\text{g})$	8.300	0.500	72/1
155	$(\text{CH}_3)_2\text{Si}(\text{CH}_2)_4(\text{l}) = (\text{CH}_3)_2\text{Si}(\text{CH}_2)_4(\text{g})$	9.000	0.500	72/1
156	$\text{Si}_2\text{C}_6\text{H}_{16}(\text{l}) = \text{Si}_2\text{C}_6\text{H}_{16}(\text{g})$	9.800	0.500	72/1
157	$(\text{CH}_3)_3\text{SiH}(\text{g}) = (\text{CH}_3)_3\text{Si}(\text{g}) + \text{H}(\text{g})$	80.200	0.500	71/3
		Residual	-0.012	
158	$(\text{CH}_3)_3\text{SiH}(\text{g}) = (\text{CH}_3)_2\text{SiH}(\text{g}) + \text{CH}_3(\text{g})$	76.500	0.500	71/3
159	$(\text{CH}_3)_6\text{Si}_2(\text{g}) = 2(\text{CH}_3)_3\text{Si}(\text{g})$	67.000	2.000	68/3
		Residual	-0.188	
160	$\text{CH}_3\text{SiH}_3(\text{g}) = \text{CH}_3\text{SiH}_2^+(\text{g}) + \text{H}(\text{g})$	272.100	9.000	70/5
161	$(\text{CH}_3)_2\text{SiH}_2(\text{g}) = (\text{CH}_3)_2\text{SiH}^+(\text{g}) + \text{H}(\text{g})$	256.400	9.000	70/5
162	$(\text{CH}_3)_2\text{SiH}_2(\text{g}) = \text{CH}_3\text{SiH}_2^+(\text{g}) + \text{CH}_3(\text{g})$	265.400	9.000	70/5
163	$(\text{CH}_3)_3\text{SiH}(\text{g}) = (\text{CH}_3)_3\text{Si}^+(\text{g}) + \text{H}(\text{g})$	242.600	9.000	70/5
		Residual	-1.312	
		247.200	9.000	66/5
		Residual	3.288	
		244.400	9.000	67/5
		Residual	0.488	
		246.100	9.000	72/2
		Residual	2.188	
		251.600	9.000	70/5
		Residual	-3.350	
		258.300	9.000	68/2
		Residual	3.350	
		236.400	9.000	70/5
		Residual	-7.659	
		242.800	9.000	66/5
		Residual	-1.259	
		239.800	9.000	67/5
		Residual	-4.259	
		242.800	9.000	69/4
		Residual	-1.259	
		238.500	9.000	66/5
		231.300	9.000	66/5
		Residual	0.412	
		230.600	9.000	67/5
		Residual	-0.288	

Reaction		ΔH° (kcal/mol)	Error (kcal/mol)	Ref
168	$(CH_3)_6Si_2(g) = (CH_3)_5Si_2^+(g) + CH_3(g)$	232.900	9.000	67/6
	Residual	2.012		
	235.700	9.000	69/4	
	Residual	4.812		
	237.100	9.000	72/3	
	Residual	-2.673		
169	$(CH_3)_8Si_3(g) = (CH_3)_3Si^+(g) + (CH_3)_5Si_2(g)$	246.800	9.000	72/3
	Residual	2.388		
	247.700	9.000	72/2	
	Residual	3.288		
170	$(CH_3)_8Si_3(g) = (CH_3)_5Si_2^+(g) + (CH_3)_3Si(g)$	229.500	9.000	72/3
	Residual	5.512		
171	$(CH_3)_8Si_3(g) = (CH_3)_7Si_3^+(g) + CH_3(g)$	238.500	9.000	72/3
	Residual	-1.579		
172	$(CH_3)_{10}Si_4(g) = (CH_3)_3Si^+(g) + (CH_3)_7Si_3(g)$	259.400	9.000	72/3
173	$(CH_3)_{10}Si_4(g) = (CH_3)_5Si_2^+(g) + (CH_3)_5Si_2(g)$	229.000	9.000	72/3
	Residual	-2.838		
174	$(CH_3)_{10}Si_4(g) = (CH_3)_7Si_3^+(g) + (CH_3)_3Si(g)$	220.200	9.000	72/3
	Residual	1.579		
175	$(CH_3)_{12}Si_5(g) = (CH_3)_3Si^+(g) + (CH_3)_9Si_4(g)$	262.200	9.000	72/3
176	$CH_3SiCl_3(l) + 3H_2O(l) = CH_3Si(OH)_3(a) + 3HCl(a)$	-51.500	0.500	53/3
177	$2C(cs) + Si(cs) + 4H_2(gs) + O_2(gs)$ $= (CH_3)_2Si(OH)_2(a)$	-205.000	5.000	66/1
	Residual	1.238		
178	$2(CH_3)_3SiOH(l) = (CH_3)_6Si_2O(l) + H_2O(l)$	-2.400	0.200	67/7
179	$(CH_3)_3SiOH(l) = (CH_3)_3SiOH(g)$	10.900	0.400	53/2
180	$2(CH_3)_3SiOC_2H_5(l) + H_2O(l)$ $= (CH_3)_6Si_2O(l) + 2C_2H_5OH(50H_2O)$	-11.400	0.200	72/4
181	$12C(cs) + Si(cs) + 6H_2(gs) + O_2(gs)$ $= (C_6H_5)_2Si(OH)_2(a)$	-155.000	5.000	66/1
182	$(CH_3)_6Si_2O(l) + 12O_2(gs) + 12HF(80H_2O)$ $= 2H_2SiF_6(a4) + 6CO_2(g) + 13H_2O(l)$	-1484.800	4.200	72/1
	Residual	-0.699		
183	$(CH_3)_6Si_2O(l) + 12O_2(gs) + 12HF(25H_2O)$ $= 2H_2SiF_6(a1) + 6CO_2(g) + 13H_2O(l)$	-1484.510	1.490	64/2
	Residual	-0.037		
184	$(CH_3)_6Si_2O(l) + 12O_2(gs) + 12HF(4.0H_2O)$ $= 2H_2SiF_6(a2) + 6CO_2(g) + 13H_2O(l)$	-1485.460	1.620	64/2
	Residual	-0.186		
185	$(CH_3)_8Si_2O_2(l) + 16O_2(gs) + 18HF(80H_2O)$ $= 3H_2SiF_6(a4) + 8CO_2(g) + 18H_2O(l)$	-1983.300	2.400	72/1
186	$(CH_3)_{10}Si_4O_3(l) + 20O_2(gs) + 24HF(80H_2O)$ $= 4H_2SiF_6(a4) + 10CO_2(g) + 23H_2O(l)$	-2492.200	5.000	72/1
187	$2(CH_3)_{12}Si_5O_4(l) + 48O_2(gs) + 60HF(80H_2O)$ $= 10H_2SiF_6(a4) + 24CO_2(g) + 56H_2O(l)$	-5965.800	14.600	72/1
188	$(CH_3)_6Si_2O(l) = (CH_3)_6Si_2O(g)$	8.900	0.400	47/1
189	$(CH_3)_8Si_3O_2(l) = (CH_3)_8Si_3O_2(g)$	9.500	0.500	72/1
190	$(CH_3)_{10}Si_4O_3(l) = (CH_3)_{10}Si_4O_3(g)$	11.500	0.500	72/1
191	$(CH_3)_{12}Si_5O_4(l) = (CH_3)_{12}Si_5O_4(g)$	12.700	0.500	72/1
192	$(CH_3)_3SiCl(l) + 6O_2(gs)$ $= SiO_2(a) + 3CO_2(g) + 4H_2O(l) + HCl(600H_2O)$	-720.900	5.000	70/6
	Residual	-0.186		
193	$CH_3SiCl_3(l) + 2O_2(gs)$ $= SiO_2(a) + CO_2(g) + 3HCl(600H_2O)$	-283.800	5.000	70/6
	Residual	8.198		
194	$(CH_3)_2SiCl_2(l) + 4O_2(gs)$ $= SiO_2(a) + 2CO_2(g) + 2H_2O(l) + 2HCl(600H_2O)$	-503.400	5.000	68/4
	Residual	1.135		
195	$2(CH_3)_2HSiCl(l) + 9O_2(gs)$ $= 2SiO_2(a) + 4CO_2(g) + 6H_2O(l) + 2HCl(600H_2O)$	-1141.000	5.000	68/4
196	$2CH_3HSiCl_2(l) + 5O_2(gs)$ $= 2SiO_2(a) + 2CO_2(g) + 2H_2O(l) + 4HCl(600H_2O)$	-707.200	5.000	68/4

	Reaction	ΔH° (kcal/mol)	Error (kcal/mol)	Ref
197	$(CH_3)_2SiCl_2(l) + 2H_2O(l) = (CH_3)_2Si(OH)_2(a) + 2HCl(500H_2O)$	-32.090 Residual -32.500 Residual	0.200 0.055 0.500 -0.355	66/1 53/3
198	$(C_6H_5)_2SiCl_2(l) + 2H_2O(l) = (C_6H_5)_2Si(OH)_2(a) + 2HCl(500H_2O)$	-31.640 7.210 Residual	0.500 0.200 0.007	66/1 60/1
199	$(CH_3)_3SiCl(l) = (CH_3)_3SiCl(g)$	7.540 Residual	0.200 0.002	60/1
200	$(CH_3)_2SiCl_2(l) = (CH_3)_2SiCl_2(g)$	7.500 Residual	0.200 -0.003	60/1 60/1
201	$CH_3SiCl_3(l) = CH_3SiCl_3(g)$	3.600 Residual	2.000 0.503	48/2
202	$2(CH_3)_3SiCl(g) = (CH_3)_2SiCl_2(g) + (CH_3)_4Si(g)$	3.600 Residual	2.000 0.328	48/2
203	$2(CH_3)_2SiCl_2(g) = CH_3SiCl_3(g) + (CH_3)_3SiCl(g)$	Residual		
204	$2(CH_3)_3SiCl(l) + H_2O(l) = (CH_3)_6Si_2O(l) + 2HCl(50H_2O)$	-21.600 Residual	0.200 0.091	67/7
205	$2(CH_3)_3SiCl(l) + H_2O(l) = (CH_3)_6Si_2O(l) + 2HCl(800H_2O)$	-22.360 Residual	0.120 -0.033	66/6
206	$(CH_3)_3SiCl(l) + 2(C_2H_5)_2NH(l) = (CH_3)_3SiN(C_2H_5)_2(l) + (C_2H_5)_2NH_2Cl(c)$	-32.100 244.400	0.300 9.000	62/6 68/2
207	$(CH_3)_3SiCl(g) = (CH_3)_2SiCl^+(g) + CH_3(g)$	246.800	9.000	68/2
208	$2(CH_3)_3SiBr(l) + H_2O(l) = (CH_3)_6Si_2O(l) + 2HBr(50H_2O)$	-27.400 7.800 246.800	0.200 0.200 9.000	67/7 67/7 68/2
209	$(CH_3)_3SiBr(l) = (CH_3)_3SiBr(g)$	232.900	9.000	67/5
210	$(CH_3)_3SiBr(g) = (CH_3)_2SiBr^+(g) + CH_3(g)$	Residual	3.550	
211	$(CH_3)_3SiI(g) = (CH_3)_3Si^+(g) + I(g)$	225.800 Residual	9.000 -3.550	72/2
212	$(CH_3)_3SiI(g) = (CH_3)_2SiI^+(g) + CH_3(g)$	237.500	9.000	68/2
213	$2(CH_3)_3SiSC_4H_9(l) + H_2O(l) = (CH_3)_6Si_2O(l) + 2C_4H_9SH(l)$	-3.600 9.700 221.000	0.200 0.100 8.000	67/7 67/7 70/4
214	$(CH_3)_3SiSC_4H_9(l) = (CH_3)_3SiSC_4H_9(g)$	Residual	5.296	
215	$2SiC(c1) + N_2(gs) = 2SiCN(g)$	-36.000 Residual	4.000 -1.324	70/4
216	$2Si(g) + N_2(gs) + 2C(cs) = 2SiCN(g)$			
217	$2(CH_3)_3SiNHCH_3(l) + H_2O(l) + 2HCl(50H_2O) = (CH_3)_6Si_2O(l) + 2CH_3NH_3Cl(50H_2O)$	-58.600 -50.600 -34.500	0.200 0.200 0.200	67/7 67/7 67/7
218	$2(CH_3)_3SiN(CH_3)_2(l) + H_2O(l) + 2HCl(50H_2O) = (CH_3)_6Si_2O(l) + 2(CH_3)_2NH_2Cl(50H_2O)$	Residual	0.253	
219	$(CH_3)_6Si_2NH(l) + H_2O(l) + HCl(50H_2O) = (CH_3)_6Si_2O(l) + NH_4Cl(50H_2O)$	-38.900 -34.490 -38.900	0.200 0.040 0.200	67/7 66/6 67/7
220	$(CH_3)_6Si_2NH(l) + H_2O(l) + HCl(1500H_2O) = (CH_3)_6Si_2O(l) + NH_4Cl(1500H_2O)$	Residual	-0.010	
221	$(CH_3)_6Si_2NCH_3(l) + H_2O(l) + HCl(50H_2O) = (CH_3)_6Si_2O(l) + CH_3NH_3Cl(50H_2O)$	-98.200 8.600 7.600 7.900 9.300 13.000	1.000 0.100 0.100 0.100 0.100 0.100	67/7 67/7 67/7 67/7 67/7 67/7
222	$2(CH_3)_9Si_3N(c) + 3H_2O(l) + 2HCl(50H_2O) = 3(CH_3)_6Si_2O(l) + 2NH_4Cl(50H_2O)$			
223	$(CH_3)_3SiNHCH_3(l) = (CH_3)_3SiNHCH_3(g)$			
224	$(CH_3)_3SiN(CH_3)_2(l) = (CH_3)_3SiN(CH_3)_2(g)$			
225	$(CH_3)_6Si_2NH(l) = (CH_3)_6Si_2NH(g)$			
226	$(CH_3)_6Si_2NCH_3(l) = (CH_3)_6Si_2NCH_3(g)$			
227	$(CH_3)_9Si_3N(c) = (CH_3)_9Si_3N(g)$			

References

- 12/1 Mullert, *Z. anorg. Chem.*, 75, 198 (1912).
21/1 Wietzel, *Z. anorg. Chem.*, 116, 71 (1921).
22/1 Latimer, *J. Amer. Chem. Soc.*, 44, 90 (1922).
31/1 Thiesson and Koerner, *Z. anorg. Chem.*, 197, 307 (1931).
33/1 Stock, *Hydrides of Boron and Silicon*, Cornell University Press, Ithaca, N.Y., (1933).
46/1 Whitmore, Somner, DiGiorgio, Strong, Van Strien, Bailey, Pietrusza and Kerr, *J. Amer. Chem. Soc.*, 68, 475 (1946).
47/1 Hildebrand, *J. Chem. Phys.*, 15, 727 (1947).
47/2 Stull, *Ind. Eng. Chem.*, 39, 517 (1947).
48/1 Torgeson and Sahama, *J. Amer. Chem. Soc.*, 70, 2156 (1948).
48/2 Zemany and Price, *J. Amer. Chem. Soc.*, 70, 4222 (1948).
49/1 Moore, *Nat. Bur. Stand. Circ.* 467.
51/1 King, *J. Amer. Chem. Soc.*, 73, 656 (1951).
53/1 Tannenbaum, Kaye and Lewenz, *J. Amer. Chem. Soc.*, 75, 3753 (1953).
53/2 Grubb and Osthoff, *J. Amer. Chem. Soc.*, 75, 2230 (1953).
53/3 Andrianov and Pavlov, *Dokl. Akad. Nauk. SSSR.*, 88, 811 (1953).
54/1 Rocquet and Ancey-Moret, *Bull. Soc. Chim. France*, 1038 (1954).
55/1 Porter, Chupka and Inghram, *J. Chem. Phys.*, 23, 216 (1955).
58/1 Johns and Barrow, *Proc. Phys. Soc. London*, 71, 166 (1958).
58/2 Drowart, DeMaria, and Inghram, *J. Chem. Phys.*, 29, 1015 (1958).
59/1 Suza and Seki, *Bull. Chem. Soc. Japan.*, 32, 1088 (1959).
60/1 Popov, *Plasticheskie Massy*, 8, 26 (1960); *Chem. Abs.*, 55, 11006e (1961).
60/2 Vorob'ev, Kolesov and Skuratov, *Russ. J. Inorg. Chem.*, 5, 681 (1960).
61/1 Gunn and Green, *J. Phys. Chem.*, 65, 779 (1961).
61/2 Wolf, *Z. anorg. Chem.*, 313, 228 (1961).
61/3 Davis, Anthrop and Searcy, *J. Chem. Phys.*, 34, 659 (1961).
62/1 Cochran and Foster, *J. Phys. Chem.*, 66, 380 (1962).
62/2 Steele, Nichols and Stone, *J. Amer. Chem. Soc.*, 84, 4441 (1962).
62/3 Margrave, Kana'an and Pease, *J. Phys. Chem.*, 66, 1200 (1962).
62/4 Wolf, Stahn and Schonherr, *Z. anorg. Chem.*, 319, 168 (1962).
62/5 Steele and Stone, *J. Amer. Chem. Soc.*, 84, 3599 (1962).
62/6 Claydon and Mortimer, *J. Chem. Soc.*, 3212 (1962).
63/1 Feder, Hubbard, Wise and Margrave, *J. Phys. Chem.*, 67, 1149 (1963).
63/2 Wise, Margrave, Feder and Hubbard, *J. Phys. Chem.*, 67, 815 (1963).
63/3 Saalfeld and Svec, *Inorg. Chem.*, 2, 46 (1963).
64/1 Balk and Dong, *J. Phys. Chem.*, 68, 960 (1964).
64/2 Good, Licina, DePrater and McCullough, *J. Phys. Chem.*, 68, 579 (1964).
64/3 Gunn and Green, *J. Phys. Chem.*, 68, 946 (1964).
64/4 Beezer and Mortimer, *J. Chem. Soc.*, 2727 (1964).
64/5 Schafer and Heine, *Z. anorg. Chem.*, 332, 25 (1964).
64/6 Kana'an and Margrave, *Inorg. Chem.*, 3, 1037 (1964).
64/7 Verhaegen, Stafford, and Drowart, *J. Chem. Phys.*, 40, 1622 (1964).
64/8 Verma, *Can. J. Phys.*, 42, 2345 (1964).
64/9 Ehlert and Margrave, *J. Chem. Phys.*, 41, 1066 (1964).
65/1 Timms, Kent, Ehlert, and Margrave, *J. Amer. Chem. Soc.*, 87, 2824 (1965).
65/2 Hess, Lampe and Sommer, *J. Amer. Chem. Soc.*, 87, 5327 (1965).
66/1 Ring, O'Neal, Kadhim and Jappe, *J. Organometal. Chem.*, 5, 124 (1966).
66/2 Saalfeld and Svec, *J. Phys. Chem.*, 70, 1753 (1966).
66/3 Wolf and Herbst, *Z. anorg. Chem.*, 347, 113 (1966).
66/4 Davidson and Stephenson, *Chem. Comm.*, 746 (1966).
66/5 Connor, Finney, Leigh, Haszeldine, Robinson, Sedgewick and Simmons, *Chem. Comm.*, 178 (1966).
66/6 Beezer and Mortimer, *J. Chem. Soc. A*, 514 (1966).
67/1 Exsteen, Drowart, Vander Auwera-Mahieu and Callaerts, *J. Phys. Chem.*, 71, 4130 (1967).
67/2 Zmbov and Margrave, *J. Amer. Chem. Soc.*, 89, 2492 (1967).
67/3 Wolf, *Z. Chem.*, 7, 283 (1967).
67/4 Schafer, Bruderreck and Morcher, *Z. Anorg. Chem.*, 352, 122 (1967).
67/5 Band, Davidson, Lambert and Stevenson, *Chem. Comm.*, 723 (1967).
67/6 Davidson and Stephenson, *J. Organometal. Chem.*, 7, p.24 (1967).
67/7 Baldwin, Lappert, Pedley and Treverton, *J. Chem. Soc. A*, 1980 (1967).

- 68/1 McDonald, Williams, Thompson, and Margrave, *Mass Spectrometry in Inorganic Chemistry*. Monograph No. 72, p. 261. American Chemical Society, Washington, D.C. (1968).
- 68/2 Band, Davidson and Lambert, *J. Chem. Soc. A*, 2068 (1968).
- 68/3 Davidson and Stephenson, *J. Chem. Soc. A*, 282 (1968).
- 68/4 Hajiev and Agarunov, *J. Organometal. Chem.*, 11, 415 (1968).
- 69/1 Gross, Hayman and Mwroka, *Trans. Faraday Soc.*, 65, 2856 (1969).
- 69/2 Ryklis, Bolgar and Fesenko, *Porosh. Met.*, 9, 92 (1969).
- 69/3 Potzinger and Lampe, *J. Phys. Chem.*, 73, 3912 (1969).
- 69/4 Lappert, Simpson and Spalding, *J. Organometal. Chem.*, 17, p.1 (1969).
- 69/5 Tel'noi, Kol'yakova, Rabinovich and Vyazankin, *Dokl. Akad. Nauk. SSSR*, 185, 374 (1969).
- 70/1 Greenburg, Natke and Hubbard, *J. Chem. Thermodyn.*, 2, 193 (1970).
- 70/2 Thynne and MacNeil, *Inorg. Chem.*, 9, 1946 (1970).
- 70/3 Adams, Sharp, Wilson, and Margrave, *J. Chem. Thermodyn.*, 2, 439 (1970).
- 70/4 Muenow and Margrave, *J. Phys. Chem.*, 74, 2577 (1970).
- 70/5 Potzinger and Lampe, *J. Phys. Chem.*, 74, 719 (1970).
- 70/6 Hajiev and Agarunov, *J. Organometal. Chem.*, 22, 305 (1970).
- 71/1 JANAF Thermochemical Tables, Office of Standard Reference Data, National Bureau of Standards, Washington, D.C. 20234, June (1971).
- 71/2 Gaidis, Briggs and Shannon, *J. Phys. Chem.*, 75, 974 (1971).
- 71/3 Davidson and Lambert, *J. Chem. Soc. A*, 882 (1971).
- 72/1 Iseard, Pedley and Treverton, (unpublished results).
- 72/2 Spalding, (unpublished results).
- 72/3 Iseard, Pedley and Krisnamurthy, (unpublished results).

