AD-773 954

RE-ENTRY PHYSICS

1 25

ŝ.

1 - " - SAL (+ "

ALL A LA LA PARTA

John P. Appleton, et al

- ----

Massachusetts Institute of Technology

Prepared for: Office of Naval Research

December 1973

DISTRIBUTED BY:



U. S. DEPARTMENT OF COMMERCE 5285 Port Royal Road, Springfield Va. 22151 AD773954

ł

RE-ENTRY PHYSICS: FINAL REPORT

Cffice of Naval Research Contract No. N00014-67-A-0204-0040, NR 051-141

Reference: ONR: 438: RDC: mg

NR 061-141

Submitted by

Fluid Mechanics Laboratory Department of Mechanical Engineering Massachusetts Institute of Technology Cambridge, Mass. 02139

This research was supported by the Advanced Research Projects Agency of the Department of Defense and monitored by the Office of Naval Research under Contract No. N00014-67-A-0204-0040, NR 061-141

This document has been approved for public release and sale; its distribution is unlimited.

and the second second second second

December 1973

Reproduced by NATIONAL TECHNICAL INFORMATION SERVICE U S Department of Commerce Springfield VA 22151

REPORT DOCIMENTATION PACE	READ INSTRUCTIONS
REPORT NUMBER	BEFORE COMPLETING FORM
Final Report	2A 773951
	11 CO 10/70
RE-ENTRY PHYSICS: FINAL REPORT	11/69 - 12//2
	S. PERFORMING ORG. REPORT NUMBER
AUTHOR(a)	B. CONTRACT OR GRANT NUMBER(A)
Prof. John P. Appleton	N-0014-6/-A-0204-0640
Prof.James A. Fay	
PERFORMING ORGANIZATION NAME AND ADOPESS	10. PROGRAM ELEMENT, PROJECT, TASK APEA & WORK UNIT NUMBERS
Fluid Mechanics Laboratory, Department of	
of Technology, Cambridge, MA 02139	NR 061-141
CONTROLLING OFFICE NAVE AND AD JRESS	13. REPURT DATE
Office of Naval Research (Code 438)	December 1973
arlington, Virginia	13. NUMBER OF PAGES
MONITORING AGENCY NAME & ADDRESSII different from Controlling Office	13. SECURITY CLASS. (of this report)
- monte oning the dot while a Auguresatir different from Condotting Utile)	INCLASSIFIED
	156. DECLASSIFICATION DOW GRADING
ON STATENENT (of this Report)	
Distribution of this document is unlimited	
Distribution of this document is unlimited DISTRIBUTION STATEMENT (of the abstract entered in Block 20, 11 different to	rom Report)
Distribution of this document is unlimited DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different is	rom Ref.ort)
Distribution of this document is unlimited DISTRIBUTION STATEMENT (of the abetrect entered in Block 20, 11 different is	rom Refort)
Distribution of this document is unlimited Distribution STATEMENT (of the abetrect entered in Block 20, if different to	rom R+Cort)
Distribution of this document is unlimited DISTRIBUTION STATEMENT (of the obstract entered in Block 20, if different for SUPPLEMENTARY NOTES	rom Ref.orf)
Distribution of this document is unlimited DISTRIBUTION STATEMENT (of the obstreet entered in Block 20, 11 different & SUPPLEMENTARY NOTES	rom Rscort)
Distribution of this document is unlimited DISTRIBUTION STATEMENT (of the abetrect entered in Block 20, if different to SUPPLEMENTARY NOTES	rom R+f.orf)
Distribution of this document is unlimited DISTRIBUTION STATEMENT (of the abetrect entered in Block 20, 11 different & SUPPLEMENTARY NOTES	rom Rs(ort)
Distribution of this document is unlimited DISTRIBUTION STATEMENT (of the abetrect entered in Black 20, if different to SUPPLEMENTARY NOTES KEY WORDS (Continue on reverse side if necessary and identify by block sumLe	rom R+f.orf)
Distribution of this document is unlimited DISTRIBUTION STATEMENT (of the abetrect entered in Block 20, 11 different for SUPPLEMENTARY NOTES KEY WORDS (Continue on reverse elde if necessary and identify by block sumle inetics Phase-Space Theory	rom Ref.orf)
Distribution of this document is unlimited Distribution STATEMENT (of the abetrect entered in Block 20, if different to SUPPLEMENTARY NOTES KEY WORDS (Continue on reverse eide if necessary and identify by block sumle inetics Phase-Space Theory issociation Monte Carlo Trajectorie	roæ: K+¢ort)
Distribution of this document is unlimited DISTRIBUTION STATEMENT (of the abetrect entered in Block 20, 11 different to SUPPLEMENTARY NOTES KEY WORDS (Continue on reverse elde if necessary and identify by block number inetics Phase-Space Theory issociation Monte Carlo Trajectorie ecombination Stratified Flows	rom K+(ort)
Distribution of this document is unlimited Distribution STATEMENT (of the abetreed entered in Black 20, if different is SUPPLEMENTARY NOTES KEY WORDS (Continue on reverse elde if necessery and identify by block sumile inetics Phase-Space Theory dissociation Monte Carlo Trajectorie ecombination Stratified Flows on-Molecule Reactions Momentunless Wakes	rom R+f.orf)
Distribution of this document is unlimited Distribution STATEMENT (of the obstreet entered in Block 20, 11 different for SUPPLEMENTARY NOTES KEY WORDS (Continue on reverse elde if necessary and identify by block sumle inetics Phase-Space Theory dissociation Monte Carlo Trajectorie ecombination Stratified Flows on-Molecule Reactions Momentumless Wakes ABSTRACT (Continue on reverse elde If necessary and identify by block number	rom <i>Rs</i> (ort)
Distribution of this document is unlimited Distribution STATEMENT (of the abstract entered in Block 20, if different is SUPPLEMENTARY NOTES KEY WORDS (Continue on reverse elde if necessary and identify by block number inetics Phase-Space Theory dissociation Monte Carlo Trajectorie ecombination Stratified Flows on-Molecule Reactions Momentumless Wakes ABSTRACT (Continue on reverse elde if mecessary and identify by block number The modified phase space theory of reaction r	rom Ketori) rom Ketori) r) s rates and Monte Carlo trajec-
Distribution of this document is unlimited Distribution STATEMENT (of the obstract entered in Block 20, 11 different t SUPPLEMENTARY NOTES KEY WORDS (Continue on reverse elde if necessary and identify by block sumle inetics Phase-Space Theory dissociation Monte Carlo Trajectorie ecombination Stratified Flows on-Molecule Reactions Momentumless Wakes ABSTRACT (Continue on reverse elde If necessary and identify by block number The modified phase space theory of reaction r bry calculations we applied to the problem of thr	rom Refort) rom Refort) s s s cates and Monte Carlo trajec- ree-body recombination and which includes
Distribution of this document is unlimited Distribution STATEMENT (of the obstreed entered in Block 20, if different t SUPPLEMENTARY NOTES SUPPLEMENTARY NOTES KEY WORDS (Continue on reverse side if necessary and identify by block scance Ginetics Phase-Space Theory Dissociation Monte Carlo Trajectorie ecombination Stratified Flows On-Molecule Reactions Momentumless Wakes ABSTRACT (Continue on reverse side if necessary and identify by block manber The modified phase space theory of reaction r bry calculations we applied to the problem of the Issociation of diatomic molecules. Reactions stu	rom Refort) rom Refort) s s cates and Monte Carlo trajec- cee-body recombination and idied include:
Distribution of this document is unlimited Distribution STATEMENT (of the obstreed entered in Black 20, if different is SUPPLEMENTARY NOTES KEY WORDS (Continue on reverse elds if necessary and identify by block number dinetics Phase-Space Theory Dissociation Monte Carlo Trajectorie ecombination Stratified Flows On-Molecule Reactions Momentumless Wakes ABSTRACT (Continue on reverse elds if necessary and identify by block number The modified phase space theory of reaction r bry calculations we applied to the problem of the Issociation of diatomic molecules. Reactions stu N ₂ + A ₁ + N + N + Ar	rom Racord) rom Racord) s s s cates and Monte Carlo trajec- cee-body recombination and idied include:
Distribution of this document is unlimited Distribution STATEMENT (of the abstract entered in Black 20, if different is SUPPLEMENTARY NOTES KEY WORDS (Continue on reverse eide if necessary and identify by block number dinetics Phase-Space Theory dissociation Monte Carlo Trajectorie ecombination Stratified Flows On-Molecule Reactions Momentumless Wakes ABSTRACT (Continue on reverse eide if necessary and identify by block number The modified phase space theory of reaction r bry calculations we applied to the problem of thr issociation of diatomic molecules. Reactions stu N ₂ + A ₁ + N + N + Ar NC + A = + N + Of + Ar	rom Reform) From Reform) S S Trates and Monte Carlo trajec- rece-body recombination and Idied include: I by NAL TECHNICAL MATION SERVICE
Distribution of this document is unlimited Distribution STATEMENT (of the abstract entered in Block 20, 11 different t SUPPLEMENTARY NOTES KEY WORDS (Continue on reverse eide if necessary and identify by block science inetics Phase-Space Theory dissociation Monte Carlo Trajectorie ecombination Stratified Flows on-Molecule Reactions Momentumless Wakes ABSTRACT (Continue on reverse eide II necessary and identify by block science The modified phase space theory of reaction r bry calculations we applied to the problem of thr issociation of diatomic molecules. Reactions stu N ₂ + A ₁ + N + N + Ar NATIC HC + Ar + H + C2 + Ar H, + H + H + H + H + H	rom Refort) rom Refort) s s s cates and Monte Carlo trajsc- ree-body recombination and udied include: t by NAL TECHNICAL MATION SERVICE partment of Commerce ingfield VA 22151
Distribution of this document is unlimited Distribution STATEMENT (of the abstract entered in Block 20, 11 different t SUPPLEMENTARY NOTES KEY WORDS (Continue on reverse side if necessary and identify by block mumber inetics Phase-Space Theory dissociation Monte Carlo Trajectorie ecombination Stratified Flows on-Molecule Reactions Momentumless Wakes ABSTRACT (Continue on reverse side if necessary and identify by block mumber The modified phase space theory of reaction r bry calculations we applied to the problem of thr issociation of diatomic molecules. Reactions stu- N ₂ + A _r + N + N + Ar N ₂ + A _r + N + N + Ar N ₂ + A _r + N + N + Ar NaTiC HC + Ar + H + CL + Ar H ₂ + H + H + H + H - Sp	rom Kaconi) rom Kaconi) s s s cates and Monte Carlo trajec- ree-body recombination and udied include: t by NAL TECHNICAL MATION SERVICE partment of Commerce ringfield VA 22151
Distribution of this document is unlimited Distribution STATEMENT (of the abstract entered in Block 20, 11 different & SUPPLEMENTARY NOTES KEY WORDS (Continue on reverse elds if necessary and identify by block simile inetics Phase-Space Theory dissociation Monte Carlo Trajectorie ecombination Stratified Flows on-Molecule Reactions Momentumless Wakes ABSTRACT (Continue on reverse side if necessary and identify by block number The modified phase space theory of reaction r bry calculations we applied to the problem of thr issociation of diatomic molecules. Reactions stu N ₂ + A ₁ + N + N + Ar N ₂ + A ₁ + N + N + Ar N ₂ + A ₁ + N + N + Ar HC + Ar + H + CL + Ar H ₂ + H + H + H + H + H FORM 1473 EDITION OF 1 NOV 65 IS OBSOLETE	rom K+(ort) From K+(ort) S S Cates and Monte Carlo trajec- ree-body recombination and Idied include: I by NAL TECHNICAL MATION SERVICE partment of Commerce ringfield VA 22151 INCLASSIFIED

ŝ

÷

VIANAN'S

• •

UNCLASSIFIED

LUURITY CLASSIFICATION OF THIS PAGE (When Date Entered)

and in the ion molecule reactions

 $F + (Ar, N_2, CO) \stackrel{\neq}{\leftarrow} F + e + (Ar, N_2, CO)$ Br + Ar $\stackrel{\neq}{\leftarrow}$ Br + e + Ar

On the basis of comparisons made between the phase space theory predictions of the rate coefficients and experimental measurements it is concluded that the phase space theory is capable of yielding good a priore estimates of threebody rate coefficients over a wide temperature range.

Exploratory experiments were carried out in a stratified-flow wind tunnel in an attempt to measure the growth of momentumless wake in a stratified flow. It was concluded that the small wind tunnel was not a sufficiently refined facility for making the desired measurements.

UNCLASSIFIED

. ia/

SECURITY CLASSIFICATION OF THE PAGE(When Dats Salered)

ABSTRACT

The modified phase space theory of reaction rates and Monte Carlo trajectory calculations we applied to the problem of threebody recombination and dissociation of diatomic molecules. Reactions studied include:

 $N_{2} + A_{r} \stackrel{\neq}{\leftarrow} H + N + Ar$ $HC\ell + Ar \stackrel{\neq}{\leftarrow} H + C\ell + Ar$ $H_{2} + H \stackrel{\neq}{\leftarrow} H + H + H$

and in the ion-molecule reactions

 F^{-} + (Ar, N₂, CO) $\stackrel{2}{\leftarrow}$ F + e⁻ + (Ar, N₂, CO) Br⁻ + Ar $\stackrel{2}{\leftarrow}$ B1 + e⁻ + Ar

On the basis of comparisons made between the phase space theory predictions of the rate coefficients and experimental measurements it is concluded that the phase space theory is capable of yielding good a priore estimates of three-body rate coefficients over a wide temperature range.

Exploratory experiments were carried out in a stratifiedflow wind tunnel in an attempt to measure the growth of momentumless wake in a stratified flow. It was concluded that the small wind tunnel was not a sufficiently refined facility for making the desired measurements.

ie

SUMMARY OF RESEARCH CHEMICAL KINETICS

Monte Carlo Trajectory Calculations of the Three-Body Recombination and Dissociation of Diatomic Molecules^{(1)*}

The modified phase-space theory of reaction rates was applied to the problem of the three-body recombination and dissociation of diatomic molecules. The results illustrate the important influence of the weak a tractive minimum in the third-body interaction potential and the effect of barrier penetration for recombination at low temperatures. The system N_2 + Ar was used as a typical illustrative example, and good agreement was obtained between the theoretical predictions and the experimental measurements of the reaction rate coefficients over the temperature range 200-12,000°K. The recrossing factor and the nonequilibrium factor were obtained from Monte Carlo trajectory calculations for states near the dissociation limit. The trajectories were sampled within the reaction zone with a weight proportional to the equilibrium reaction rate, and numerically integrated in both timewise directions to determine the complete histories of the collisions. A simple, separable function for the equilibrium transition rate $R(\varepsilon_i, \varepsilon_f)$ from initial energy states ε_i to final energy states ε_f , which could be characterized by three parameters, was obtained to fit the numer-

*Numbers in parenthesis refer to publications listed and page 7

-1-

ical data and cas used to solve the steady-state master equation. Histributions of the trajectories with respect to energies and impact parameters are presented, and several typical trajectories are shown to illustrate the important features of the collisions. The contribution to the over-all reaction rate from the "complex mechanism" was also obtained.

Monte Carlo Trajectory Calculations of the Dissociation of HCL in Ar⁽²⁾

The modified pahse-space theory of reaction rates was applied to the dissociation of HCL in a heat bath of argon atoms. Excellent agreement was obtained between the theoretical predictions and the shock-tube measurements of the dissociation rate coefficient over the temperature range 2500-5000°K. The recrossing correction factor and nonequilibrium correction factor were obtained from Monte Carlo trajectory calculations for states near the dissociation limit. The trajectories were sampled within the reaction zone, with a weight proportional to the equilibrium reaction rate, and numerically integrated in both timewise directions to determine the complete histories of the collisions. A simple separable function for the equilibrium causition rate $R(\varepsilon_i, \varepsilon_f)$ from initial states ε_i to final states ε_f was obtained to fit the numerical data with sufficient accuracy and was used to solve the steady-state master equation. Important features of collisions of highly asymmetric diatomic molecules

Preceding Page Blank

-3-

are discussed, and several typical reactive trajectories are shown to illustrate the improtance of rotational motion in such collisions.

Thermal Dissociation and Recombination of Hydrogen According to the Reactions: $H_2 + H \stackrel{2}{\leftarrow} H + H + H \stackrel{(3)}{\leftarrow}$

Three-body recombination and dissociation rate coefficients (k_r and k_d respectively) for the reactions $H_2 + H \stackrel{2}{\leftarrow} H + H + H$ have been evaluated for temperatures ranging between 300 and 10,000°K on the basis of the modified phase-space theory of reaction rates and Monte Carlo trajectory calculations. The semiempirical Porter-Karplus surface for H_3 was used in the calculations. Good agreement was obtained between the theoretical estimates of k_d and the bulk of the dissociation rate measurements. However, in contradistinction to some of the recent experimental investigations, neither the extremely steep temperature decay (αT^{-6}) of k at high temperatures nor the existence of a pronounced maximum in k, at moderate temperatures are supported by this theoretical work. Possible reasons for these differences are suggested. A simple ratio was deduced which relates $k_{\mu}^{(H)}$ and the corresponding rate coefficients of the isotopes of hydrogen. The result is in good agreement with available experimental measurements and enables one to predict the latter rate coefficients without additional trajectory calculations. An example is given to illustrate the capability of the present Monte Carlo method in obtaining other useful information about atomic and molecular collision processes.

Research initiated under ONR Contract No. N00014-67-0204-0040, NR 061-141 and completed under NSF Research Grant GP-32522X

-4-

Phase-Space Theory of Electron Detachment in Slow Atomic Collisions;

A classical phase-space theory for the excitation of electrons in slow atomic collisions is presented. An equation which governs the energy transfer getween the electron and the heavy particles is derived for slow collisions (adiabatic approximation). This equation is analogous to the first law of thermodynamics for an open system. The theory contains only one adjustable parameter and gives excellent correlation for existing experimental measurements of detachment rate coefficients of F^- + (Ar, Z_2 , CO) and Br^- + Ar. Possible modifications to the present theory are summarized. It is anticipated that the theory should be applicable to other electronic rate processes such as ionization and charge transfer.

¹ Research initiated under ONR Contract No. N00014-67-0204-0040, NR 061-141 and completed under NSF Research Great GP-32522X

-5-

MOMENTUMLESS WANE GROWTH IN STRATIFIED FLOW

Experiments were conducted in a stratified-flow .ind tunnel to determine the growth of momentumless wakes in a stratified flow. Zero momentum wakes were formed behind a flat disc supplied at its center with an axial jet, the momentum of the latter bean a sjusted to balance the calculated drag of the disc. Wake growth and ervitational collapse were detected visually by smoke injection and by sampling with a hydrocarbon analyzer after introducing ethane into the jet. Experiments were conducted at disc Reynolds numbers exceeding 200, which was a sufficient condition to ensure turbulent growth.

Unile wake growth and subsequent collapse were visually evident when smoke was introduced into the wake, the iso-concentration lines of ethane showed no marked gravitational collapse corresponding to that observed with the smoke tracer, although turbulent wake growth was clearly evident. Because a long sampling time of about a minute was required to determine each concentration reading, it was concluded that some unsteadiness in vertical flow velocity component was smearing out what would otherwise be steeper vertical concentration gradients, causing significant differences between the near instantaneous smoke trace and the time-averaged ethane concentrations.

It was concluded that the steady flow technique for examining momentumless wake growth and collapse in a stratified flow would encounter significanc difficulties because of flow unsteadiness when operating close to the critical Froude number. The small wind tunnel used in these experiments was not a refined facility and could not be used for quantitative measurements of the type desired.

-6-

PUBLICATIONS

- V.H. Shui, "Monte Carlo Trajectory Calculations of Three-Body Recombination and Dissociation of Diatomic Molecules", J. Chem. Phys. <u>57</u>. ;704-1717 (1972)
- V.H. Shui, J.P. Appleton, and J.C. Keck, "Monte Carlo Trajectory Calculations of the Dissociation of HCL in Ar", J. Chem. Phys. 56, 4266-4274 (1972)
- 3. V.H. Sui, "Thermal Dissociation and Recombination of Hydrogen According to the Reactions $H_2 + H + H + H$, " J. Chem Phys. 58, 4868-4879 (1973)
- V.II. Shui, and J.C. Keck, "Phase-Space Theory of Electron Detachment in Slow Atomic Collisions," M.I.T. Fluid Mechanics Laboratory Report 73-4, May 1973. Accepted for publication in J. Chem. Phys.

-7-