# UNCLASSIFIED

# AD NUMBER

### AD843857

## NEW LIMITATION CHANGE

TO

Approved for public release, distribution unlimited

### FROM

Distribution authorized to U.S. Gov't. agencies and their contractors; Foreign Government Information; JUL 1968. Other requests shall be referred to the Army Biological Laboratory, Attn: Technical Releases Branch [TID], Fort Detrick, MD 21701.

# AUTHORITY

AMXFD, per ltr dtd 9 Feb 1972

THIS PAGE IS UNCLASSIFIED



TRANSLATION NO. 1005

DATE: July 1968

(JU) -

i li i ca a c

**.** . . . . .

### DDC AVAILABILITY NOTICE

This document is subject to special export controls and each transmittal to foreign governments or foreign nationals may be made only with prior approval of Commanding Officer, Fort Detrick, ATTN: SMUFD-AE-T, Frederick, Md. 21701.

> DEPARTMENT OF THE ARMY Fort Detrick Frederick, Maryland

T-159- Annals of Histochemistry I, 199-212, 1956

CONSIDERATIONS REGARDING THE REDUCTION OF TETRAZOLIUM SALTS (A REVIEW)

-1-

by J. Verne and R. Wegmann.

Introduction.

Tetrazolium salts play a preponderant part during the transfer of hydrogen in the chain of enzymatic reactions which lead to the reduction of a substance by oxidation of another. Their use is becoming general, but, simultaneously, there is an expansion of the category of substances capable of reducing tetrazolium salts. We thought it worthwhile to re-examine the question and to draw some useful conclusions.

#-1005

The most frequent use of tetrazolium (TZ) derivatives is in connection with the detection of tissual dehydrogenases. They werve, in the same way as methylene blue, as acceptors of hydrogen and function as redox indicators. The reduction of methylene blue yields a colorless substance, the leuco-blue, the speed of whose formation measures the rate of dehydrogenasic activity. That reduction is reversible. Furthermore, it requires; very strict anaerobic conditions. The TZ's, which are colorless and soluble in water, bring about, through reduction, a formation of colored derivatives, the formazans, which are not soluble in water. This process is irreversible and does not require the absence of oxygen.

Therefore TZ salts present an underiable advantage over the use of methylene blue, for they make possible a precise localization of the point of formation of the formazons (FZ), both in the tissue and in the cells. The clearness and intensity of the reaction depend in particular on the variety of the TZ's which are used, on the specific activity of the tissue, but also on a number of factors which we shall examine later, such as pH, isotonicity, presence of cations, a well-plugged medium, duration of incubation, not to mention the mode of manipulation of the tissue itself. All those conditions, connected with the choice of the substrate, combine to delimit the nature of the substances which bring about a reduction of TZ salts.

1. Different tetrazolium salts.

There are several tetrazolium salts which all possess characteristica enabling them, to a various degree, to achieve the histochemical detection, or to serve in biochemical dosages. The first to be utilized was the 2-3-5 triphenyltetrazolium or TTZ (Kuhn and Jerchel, 1941) (28). Its general constitution and its reduction to FZ is represented by fig. 1.

 $N = N + -C_{6} + 3$  + 2 + 4  $N = N - C_{6} + 3$ +2+++  $r = c_1 - c_2 + 3$ (red color) Formozan (color: red) Triphenyl salt (colorless) tetrazolium or TTZ

Figure 1

There are, furthermore:

3

- 2-3 diphenyl 5 methyl tetrazolium (40) chloride or DTZ.

- 2-2' (p-diphenylene)-bis-(3-5 diphenyl-tetrazolium) chloride, which is Neotetrazolium or NTZ.

-- 3:3'-dismisole bis 4:4' (3:5 diphenyl-tetrazolium) chloride, or tetrazolium Blue (43) or BTZ. It is therefore the dimethoxy derivate of neotetrazolium.

- 2B neotetrazolium (49) phosphete or PTZ.

2.5 diphenyl 3 (4-styrylphenyl) tetrazolium chloride or M and B 1767 (43), or STZ.

FTZ yields a red formazan., BTZ a blue FZ, and so dies NTZ. Pearce (43) has noted that BTZ yields FZ's of a different color, according to the compound: blue to purple for the sulfhydrides, red for the lipids, blue for sugar reducing agents. The reduction of TZ salts is achieved progressively (Jerchel, 1949) (26) and forms a red\_semi-reduced compound, then a reduced blue compound (Kun, 1951) (29).

According to the studies made by Shelton and Schneider in 1952 (56) regarding the differences in the activity of TTZ, BTZ, NTZ and VTZ, and, according to the comparative study concerning TTZ and NTZ made by Cascarano and Zweifach in 1955 (10), TTZ is inferior to NTZ when it comes to obtain cytological details. It is spontaneously reduced, pales under the impact of light, or with the passing of time. It will not be reduced by the blood vessels, the skebetal muscles and the peripherical nerves (10). For Shelton and Schneider (56), BTZ appears inferior to NTZ because of the long period of incubation and of the irregularity of the precipitation of the formozans' crystals. VTZ has a major drawback: its formozans' crystals are long and fine and destroy cell stuctures (56).

The use of a TZ derivative rather than of another derivative has been prompted by those differences of reactivity. Sometimes, it was also difficult to obtain the various kinds of TZ. Their use must be considered in function of the problem that must be studied and  $\frac{1}{2}$  to  $\frac{1}{2}$  to  $\frac{1}{2}$  be stated that one is absolutely superior to the

-2-

other.

TTZ is thus used by Black and his cides (4,5,6,7,04), by Brodie and Gots (9), by Bodine, Lu and West (8), by Coscarano and Zweifach (19), by Fried and Zweifach (20), by Gomori (21), by Gonse and Yotsuganagi (22), by Jensen, Sacks and Beldauski (25), by Jerchel (26), by Kuhn and Jerchel (28), by Kun (29), by Kun and Abood (30), by Lillie(31), by Lison (32), by Mattson, Jensen and Dutcher (35), by McMillan, Klatzo and Duff (37), by Everson Pearse (43, 44), by Roberts (48, 49), by Shelton and Schneider (56), by Smith (58), by Steigleder (60).

BTZ has been used much less, in particular by Cooper (11), Pearse (43), by Rutenburg, Gofstein and Seligman (52), by Findlay (16).

NTZ occupies an intermediary position and has been used especially by Bajusz and Szirmai (2), by Cascarano and Zweifach (11), byForaker and Denham (17), by Foraker, Denham and Mitchel (18), by Fried and Zweifach (20), by Martin, Cooper, Chaudhuri and Green (34), by Mustakallio (38, 39), by Mustakallio and Jannes (40), by Pearse (43, 44), by Rennels and Ruskin (46), by Shelton and Schneider (56) and by Villareal and Burgos (63), by Findlay (16).

Some other salts have been used only rarely, the ITZ by Defendi and Paason (12) and by Pearson and Defendi (45), by Seligman, Gofstein and Rutenburg (54); DTZ by by Roberts (49), PTZ by Roberts (49). VTZ by Shelton and Schneider (56) and by Roberts (49), STZ by Pearse (43) and TZ purple (PPTZ) by Ortmann (42) and by Steigleder (61).

Table 1 provides a summary of the results obtained by some authors who have used several TZ salts and compared the results.

Authyr	Ref.	TTZ	BTZ	MTZ	YTz	PT2	בדע
Cascarano Findlay	(10) (16)	2	- 2	33	-	-	
Pearse	(43)		2	3	-	~	~
Roberts	(49)	1	2	3	2	2	0
Shelton	(56)	2	3	3	3		

TABLE 1

Mean intensity of reactions by various TZ salts.

(o : no reaction; 1 : feeble reduction; 2: mean reduction; 3:strong reaction).

i i i

-3-

2. Nature of the compounds reduced by TZ.

อ

If TZ salts serve as activity tests of a whole series of compounds characterized by their reducing power, the nature of those substances can be very variable. a) Endogenous reductases.

-4-

The primary condition for bringing them into evidence is the use of fresh cuts, which have not been frozen, nor dried, nor, especially, fixed in a fixator of any nature whatsoever (Shelton and Schneider (56)). The simple section of cuts with a rezor may demage them (Cascarano and Zweifach (10)), and so may also do the contact between them and the water which collects the cuts (Seligman and Rutenburg ((55)). Contact with heavy metals (Barnet and Seligman (3)) or the presence of metabolical inhibitors (Black, Kleiner and Speer (5)) destroys or prevents their action. Reduction only occurs within pH limits comprized within pH 6 and 7. It is therefore quite improbable that in those conditions it should not be the case of one or several enzymatic systems. The activity of this endogenous reductase can be intensified in the vessels by addition of mannose, which appears to act rather as a co-factor rather than in a role of substrate (Reid and Zeifach) (20).

This endogenous redactose has been sought in the kidneys (Cascarano and Zweifach) (10), (Black and Speer) (6); in the suprarenal (Black and Speer) (6); on the Cilisted (Fauré-Fremiet and Gauchery) (15); in \_\_\_\_\_\_ sea-urchin eggs (Gonse and Yotsuganagi) (22); on atheromatose scales of the rabbit's sorts (Forsker, Denham and Mitchell)18); in the mesenteric capillaries and in the carotide (Fried and Zweifach) (20); in the heart muscle (Cooper) (11).

Most of those authors note an intense activity essentially localized in the lipidical enclaves. It is intense in the glomerulaes and in the reticulated (?) of the mouse (Black and Speer) (6) and in the lipidical intra-atheromatose droplets (McMillan and colleagues) (37). Cooper does not detect any activity in the ventricle or the suricle (11), before and after birth, in the rat.

The precise significance of the reductasic activity is not exactly known as yet. Rutenburg, Gofstein and Seligman (52), as well as Foraker, Denham and Mitchel (18) admit that it can be superimposed to the succinodeshydrogenasical activity. Nevertheless those latter authors (18) consider the endogenous activity to be twice more intense than that of succinodeshydrogenase, that being at the level of the ovaries.

b) Specific reductases.

They are much easier to study, for their detection may be achieved on frozen tissue. In order to search for them, one adds a specified substrate to their inwell cubation medium. On the other hand, it requires systoppered medium. Succinodehydrogenase. (SD)

In the kidney of rats who had been rendered hypertensive a dose of DOCA and by a salted fare, no change of the reaction, which is normally intense, occurs (Erankö and Niemi) (14). Infection with Mycobacterium tuberculosis provokes in the guineapig a marked fall of renal SD. The adjunction in vitro of a tissue factor extracted from the kidney of a normal ox and with a structure of a nucleotide associated to a sulfhydrile, restores normal activity (Marten, Cooper, Chaudhuri and Green) (34).

Cooper (11), on pre- and post-natal rat hearts, notes an increase of the SD activity, which seems to him to perallel the increase in the number of mitochondries. In the ovaries, a strong SD activity is observed in the folliculous cells and in the stroma cells. It is very marked in the lutenic cells and has a small intensity in the hile vessels. Foraker, Denham and Mitchel concl. de that there is, eventually, a correlation between the strong SD activity and the hormone production; Ortmann arrives at a similar conclusion (42) in his study on the placenta. The cytotrophoblaste, which is supposed to be an important area of hormone production, is very rich in SD. It is the same case with the works of Meyer, McShan and Erway (36), on the subject of the ovary and the lutenic cells.

Mustakallio (38) has made an interesting observation on the spithelium of the gall bladder and in the liver of a pregnant mouse. The SD activity, strongly marked in the non gestating mouse, diminishes in very strong proportions under the impact of the pregnancy, and seems to be only slightly influenced by fast or by ingestion of food. This author attributes this fall in the SD rate to an increase in the ostrogens rate.

In the malignant hepatic tumors, induced by p-dimethylazobenzene, Pearson and Defendi (45) do not note any constant differences of SD activity between the normal cells and the tumoral ones. It is easy to understand the importance of the study of SD in the neoplasical processes (Rutenburg, Gofstein and Seligman) (52). . While the specificity of the succinodehydrogenesical activity can hardly be

-5-

doubted, certain authors like Fried and Zweifach (20), do not admit the presence of a specific enzyme SD in the vessels (endothelium and smooth muscle), as a result of the positive reaction obtained also with other substrates. We think that this restriction cannot be generalized for all tissues and organs. In support of this hypothesis, one can mention the works of Steigleder on the subject of the derm and the epiderm (60, 61), where there is no parallelism between the activity SD and the activity of other substrates in the pilous folicie and in the collar of the seb-ceousglands.

A more important problem concerns the activity or the absence of reduction intermediaries in the chain of hydrogen transport. Kellin and Hartree (27) think that there is a BAL-sensitive factor, probably a hematine, which serves as an intermediary between the succinodehydrogenase-cytochrome b agent and Cytochrome. If there is no other factor associated with this system, one would have the right to admit a specific evidence of the existence of SD at the exact spot of the reduction of TZ. The question is not solved, as yet. Defendi and Pearson (12) rightly that methylene blue is directly reduced by succinodehydrogenase, without the intermediary of the BAL-sensitive factor.

The study of SD presents at least the advantage of a quantitative analysis perfected by Defendi and Pearson ( :). Villareal and Burgos (63), finally, in a comparative histochemical and Diochemical study of SD in the gastric mucuous of the rat and of the frog, succeed in determining most clearly the relations between the Krebs cycle and the main cells. At their level the SD reaction is the most intense and it can be activated lwith acetyl-beta-methylcholine, or inhibited with mercurial diuretics. BAL diminishes the hyperactive reaction which is useless from the point of view of experiments.

Altogether, the use of TZ's in the study of the SD activity opens here an interesting field for research and puts this reaction among the most interesting enzymatic histochemical reactions.

#### \$-giycerophosphodehydrase (GPD).

It has been studied by Steigleder in the skin (60) and manifests itself actively in the pilous follicle and in the collect of sebaceous glands.

#### lactodehydrogenase (LD).

Also analysed by Steigleder (60), it is also active in the pilous follicle to the same degree as SD or GPD. On the other hand, it is almost absent in the collect of the sebaceous glands. According to the works of Strominger and Lowry (62), on the brain, the LD activity is parallel to the aldolasic activity. In order to become manifest, it requires the association of DPN (diphosphonucleotide).<sup>†</sup> The absense of an addition of DNP may explain the results obtained by Steigleder.

- 6 -

malico- and glutomodehydragenases (MG and GD).

In principle, these enzymatic systems can also be studied, since the research carried out by Strominger and Lowry (62), which was histochemical and qualitative, at the brain level, has shown clearly marked differences of activity between these various enzymes, probably in connection with metabolical differences. Furthermore, these enzymes are poorly thermolabile.

+7-

#### desulfhydrase or desulfurase (DS)

ŧ

n

Its detection presents numerous problems, which are far from being solved. For, if one adds some cystein or some gluthation reduced to an incubation medium, one can wonder whether it is the cystein which directly reduces tetrazolium salt or if it is the desulfurase which acts on the cystein. <u>In vitro</u>, cystein reduces TTZ, PTZ and NTZ, in growing order (Roberts) (49). Cystein reduces none of the TZ salts. The addition of cyanide to cystine provokes, in growing order, a reduction of TTZ, of BTZ and VTZ and of NTZ. The glutathion ; associated with the cyanide reduces TTZ slightly and reduces strongly those other TZ salts (Roberts) (49).

All those reactions have been carried out by Roberts (49) with pH 74. Those results do not tally with those obtained by Fried and Zweifach (20), who obtained no reduction for the reduced glutathion, but, on the contrary, an inhibition, in the mesovaric vessels and in the carotide. Smith (58), when pH is above pH 9, obtains an increased reduction by cystein. One cannot, therefore, take into consideration all those reductions obtained with pH below 9; in any case, the doubt regarding the presence of desulfurase remains, because of the spontaneous raduction by sulfhydrile groups in vitro. Barrnett (sic) and Seligman (3) similarly found a reduction of the TZ blue with disulfide groups reduced in vivo with cyenide. Wid Fit is impossible to dissociate the activity proper to exogenous sulfhydriles from the enzymatic reduction with DS, there is nevertheless an indirect way to evaluate the DS activity. It consists in adding cyanide to the incubation medium, that cyanide having a pH above 9. The cyanide reduces the disulfide functions which are present and forms sulfhydriles, which teduce the TZ salts. This DS activity cannot be evidenced except on fresh material, or at least material which has not been included with paraffin. The results reported by Everson Pearse (43), nevertheless, relate solely organs fixed with formol or with a formol-sublimate, where the desulfurases have been destroyed. Nevertheless, according to observations made by Pearse (43), the strongly alcaline medium extracts, among other matters, proteins and muco-proteins. It does not seem that, up to now, it has been possible to study directly tissular desulfurases, in spite of their obvious interest. Lillie (31) admits, however, that the addition of cystein makes it possible

to study the desulfhydrasis.

Direct reduction by means of reducing agents.

As we have just seen, substances with sulfhydrile functions, when added to sections in incubation with TZ, reduce it directly. If it is true that the presence of these sulfhydriles stabilizes <u>in vitro</u> the reduction of NTZ (Findlay) (16), the addition of  $\pm$  single cyanide can already provoke a reduction of TZ. The association of cyanide and of cystein considerably increases the quantity of formazan which is produced (findlay) (16).

In fact, the reduction of the TZs is achieved not on fresh sections (cuts), but on cuts of organs fixed in various fixators, such as acetone, neutral formol, or preferably trichloracetic acid in 80% alcohol (Findlay) (16). Then the groups -- SH and SS -- present: in the tissue are characterized. Everson Pearse recommends formolated fixators (43). The cuts are cut during the freezing (16, 43) or are even used after iclusion of paraffin (43). While Pearse recommends a pH of 12.8, Findlay recommends a pH 10, having noticed a very weak reactivity of cystein to pHs above 11. On the contrary, glucids react favorably to pHs above 11. On the contrary, glucids react favorably to pH's above 11 (Smith) (58); Findlay (16).

The presence of sulfhydrile groups has been put in evidence by Findlay (16), by Pearse (43) and by Steigleder (60), mostly in the epiderm, in the proteins(?) and in microproteins, as well as in the hairs of the phaneres (Rogers) (50). Bodine, Lu and West (8) also admit a TZ reduction by sulfhydrile groups, which apparently is produced in the cells in mitosis, and in Penicillium chrysogonium, according to Fred (sic) and Knight (19). In the meristeme of plants this reduction by means of sulfhydrile groups is especially marked (Roberts) (49).

An addition of cyanide makes possible the formation of sulfhydrile groups on the basis of disulfuric groups, at the rate of two groups -- SH per group --SS -- (Findlay (16). An inhibition of sulfhydrile groups is achieved by means of iodocerate. A better scission is achieved by means of iodacetate is achieved by means of thioglycolate (Bernett and Seligman (31) and Findlay (16). A blocking of all groups -- SH thus firstly requires a blocking of the groups -- free SHs, followed by a scission of the SS groups which are also blocked.

Beside the preceding compounds, other reducing substances may intervene, such as ascorbic acid. For Defendi and Pearson, <u>in vitro</u>, (12), ascorbic acid reduces triphenyltetrazolium, with pH at 9.1. On the contrary, Mattson notes no reduction with this same pH\*(35). Kuhn and Jerchel (28) had already proven that this substance cannot be made responsible for the reduction of the TZ salts, for it does not occur above pH 9. If one excludes the possibility of endogenous reduction

-8-

with ascorbic ucid, one can achieve its exopenous reduction. Steigleder (b0) has applied it to the study of keratine and notes a reduction in the keratinized zones.

- 4 -

Can glucids bring about a reduction of TZ sults? According to Jensen and his colleagues (25), reducing sugars do not reduce TZs below pH 11. Pearse (43) obtained, with pm 12.5, the reduction of a whole series of reducing glucids, in particular with glucopyranose monomers. Polymers react less favorably, as in the case of glycogen. Findlay obtains a reduction of glucose at pH 10.2, while Smith (58) does not notice any reaction below pH 11. It must be noted that in the test of Findlay (16) reduction was achieved in the presence of cyanide, whose reducing power on the TZs is known. Glucogen has yielded doubtful results. The reduction at the level of the vessels studied by Fried and Zweifach (20) leads those authors to admit a cofactor role on the part of glucids, and more precisely on the part of mannose. Glucose seems to act less well, other hexoses rather little and the diascharids not at all. They do not admit, however, a direct reduction by those glucids. The pH of the middle part of the incubation is only of pH 7.3, and cannot promote a reduction.

The reduction of other substances has been considered. Pearse (43) gets in vitro a reduction of the DOPA. Black and Speer (6) reduce it with cortisone, with F  $\sim$  Compound and with DOCA.

According to all those data, it will be noted that the FZs can derive a whole group of reducing substances, but it is the addition of those substances to the cuts which will condition the response to be obtained. On organs in which the enzyms have been destroyed by fixation, the number of reducing compounds must become rather small and will be limited by the pH. If the pH is inferior to 11 but above 9, one can study mainly the sulfhydrile and disulfide groups. With pH in the neighborhood of 13, one will rather get reducing sugars. The analysis table presented by Pearse (43) must be completed with an indication of the pH.

3 - RELATION WITH THE LIPIDS

Lipids have a particular affinity with the derivatives from the reduction of TZs. They never contain FZ crystals and always take a pink or red coloration, with TTZ as well as with BTZ and NTZ. The grains of lipofuscines on the cuts included with paraffin take those colorings (Pearse) (43). In the screnals, the coloration occurs in the lippidic enclaves in the zones which secret steroid hormones. The coloration is restricted to those enclaves and serves as their cortico-surrenal functioning index. For Edwards and Ball (13), the phospholipids have a role of cement and of vector between the various carriers of hydrogen of the succino-oxydasical complex.

Is this a case of a special affinity of the FZs for the lipids? (Gonse and Yotsu-

ganagi) (22). Or the case of a diffusion of the FZs formed by other substances towards the lipids? Steigleder notes a close parallelism between NADI and the FZs. Yet it seems that, under the impact of aldosterone and DOCA in a suuenalectomized animal, it is no longer the enclaves which are reacting, but the lipoproteic base (Bajusz and Szirmai) (2).

The problem thus remains as to whether the enclaves directly reduce the salts of TZ.

4- RELATIONS WITH MITOCHONDRIES.

The specific succinohydrogenasical activity is very high in the granular fraction of the testicle obtained by Greif (23), where it is associated with the hyaluronidasical activity. It is parallel to the increase in the number of mitochondries in the gall bladder (Mustekallio) (38) as well as in the heart (Cooper) (11). The reduction did not occur at their level, but in their vicinity. The coloration of the mitochondries appears to be only secundary. When the FZ crystals occur, it is always outside the mitochondries. At that moment the coloration of the mitochondries disappears. In the presence of lipids no grain coloration occurs. Only the fat enclaves are tainted (Gonse and Yotsuganagi) (22). Harman (24) confirms this enzymatic activity.

If there actually is an SD activity at their level, it does not explain in which a there elements one must seek the remaining 65% of SD activity.

5- PHYSIOLOGICAL ROLE.

Surrenalectomy diminishes in all organs the reaction to NTZ (Bejusz and Szirmai) (2). DOCA inhibits the SD activity of the glomerular (gland), without touching other zones (Zweifach, Black and Speer) (64). Cortisone inhibits the activity of the glomerular (gland) (Black and Speer) (6). On the contrary, on an animal carrying a tumor, the glucorticoids inhibit and DOCA stimulates the glomerular (gland) (Black and Speer) (6). While the electrocortine does not modify the distribution of the FZ in the animals serving as control samples, it brings back to normal the aspect of the organs of surrenalectomized animals (Bajusz and Szirmai) (2).

6- TECHNICAL FACTORS.

The thickness of the cuts plays an important part. Too thin cuts may give no reduction. The latter may be spontaneous on too thick cuts. Defendi and Pearson (12) find a  $f \not\vdash$  thickness to be sufficient for a quantitative analysis of the SJ. Inversely, Cascarano and Zweifach make 1 mm cuts before obtaining a reaction. On the average, a thickness of 20 to 30  $\mu$  seems to be right.

An incubation duration of 30 to 60° intensifies the reaction of the surrenal (gland). In the kidney, on the contrary, it brings about an aberrant modification of the crystals and a variation in their shape (Cascarano and Zweifach) (10). An average duration varies between 20 and 60 minutes. It also depends on the salt which is employed and on the aim which is to be achieved. For a quantitative dosage 5 minutes are sufficient (Defendi and Pearson) (12). For SD between 30 and 120 minutes are necessary, for the sulfhydrile groups between 40 minutes at  $60^{\circ}$  C (Pearse) (43) and 4 hours at  $37^{\circ}$ C (Findlay) (16).

For the study of enzymatic systems, tensio-active substances must be rejected, for they suppress the coloration of the FZ. On the contrary, for Findlay (16), an addition of Tween 40 increases the reactivity.

An addition of inhibitors to the incubation media prevents the formation of FZ. Cyanide, azide, malonate inhibit the reduction adypocytes, but not in the vessels (Fried and Zweifach) (20). In the study of endogenous reductases, a simple agitation during the incubation (Gonse and Yotsuganagi) (22), intempestive aeration (Fried and Zweifach) (20), darkness (22) may prevent the reduction. for the study of the -SH functions, the iodacetamide, the iodacetate play the part of inhibitors. Pr. ferably, the use of ether for the killing must be avoided (Malaty and Bourne) (33).

The accivators are the glucids (see above), light (22) and, in principle but not in fact, the DPN. (20, the Al, Mg ions, the carbongtes (Rutenburg, Wolman and Seligman)(53), the Ca and Al ions (Keilin and Hartree) (27), very stric anaerobic conditions (53). A well stoppered medium, to which ions of Cl, Ca, PO<sub>4</sub> and Na have been added, provides better results (Fried and Zweifach) (20). That is how Cascarano and Zweifach had noted the presence of extracellular depositions in the absence of electrolytes of the incubation medium.

#### 7- LIVING COLORATION.

Tetrazolium salts have also been used for that purpose (35 and 47).

#### CONCLUSION

This brief review of the reduction of TTZ salts shows that numerous factors intervene in the course of their application to the study of various substrates and that it is not always easy to compare the results because of the diversity of the techniques which have been used.

#### SUMMARY

Tetrazolium salts are reduced by various substances. On fresh, non-frozen cuts it is possible to put in evidence the endogenous reduction. On cuts obtained by congelation or frigodessication, this reductasis is destroyed. The addition of

-11-

a substrate, such as succinic acid, brings into evidence the succinohydrogenasis. Other substrates may be used, such as substances with sulfhydrile functions: then sulfhydroses occur.

-12-

On fixed cuts, included or not with paraffin, at pH 10, one detects, in principle, the sulfhydrile functions. With pH 12.8 reducing glucides are essentially characterized. Those are endogenous substrates. Other substances may also be present and they are considered in their relations with the lipids. Finally, 72 salts may serve as life colorants.

(HISTOLOGY LABORATORY, FACULTY OF MEDECINE, PARIS.)

#### BIBLIOGRAPHY

 Antopol (W), Glaubach (S.) and Goldman (L.). - Effects of a new tetrazolim derivative on tissue, bacteria and onion root tips. Public Health Reports, 1948, 63: 1231-1238.

2. Bajusz (E.) and Szirmai (E.) - Wirkung des neuesten Nebennierenrindenhormons "Electrocortin-Aldesteron" auf die histochemische Reaktion von Succino-Dehydrogenase. Acta histochemica, 1955, 2: 1-15.

3. Barnett (R.\_J.) and Seligman(A.M.) -Histochemical demonstration of sulfhydryl and disulfide groups of protein. J. Nat. Cancer Inst., 1954, 14: 769-804.

4. Black (M.M.) and Xleiner (I.S.). - The use of TTC for the study of respiration of tissue slices. Science, 1949, 110:660-661.

5. Black (M.M.), Kleiner (I.S.) and Speer (F.D.). - Effects of enzyme inhibitors on in vitro dehydrogenese activity of cancer tissue slices. Proc. Soc. Exp. Biol. Med., 1951, 77:611-615.

6. Black (MM) and Speer (F.D.) - Effects of glucorticoids and mineralocorticoids on dehydrogenase acrivity of adrenal slices. A.M.A. Arch. Path. 1954, 58:433-442.
7. Black (M.M.), Zweifach (B.W.) and Speer (F.D.).- Tetrazolium salts, a new tool in general and experimental pathology. Am. J. Clin. Path., 1953, 23: 332-339.
8. Bodine (J.H.), Lu (K.H.) and West (W.L.). - Reduction of triphenyltetrazolium chloride by minerally active and blocked embryonic cells. Biol. Bull., 1952, 102: 16-21.

9. Brodie (A.F.) and Gots (J.S.) - Effects of an isolated dehydrogenase enzyme and flavoprotein on the reduct on of triphenyltetrazolium chloride. Science, 1951, 114:40-41.

10. Cascarano (j) and Zweifach (B.W.). - Comparative histochemical and quantitative study of adrenal and kidney tissue by tetrazolium technique. J. Histochem. Cytochem., 1955, 3:369-381. 11. Cooper (W.G.). - Succinic dehydrogenase activity in the pre-natal and postnatal rat heart. Anat. Rec., 1955, 123:103-123.

-13-

12. Defendi (V.) and Pearson (B.). - Quantitative estimation of succinic dehydrogenase activity in a single microscopic tissue section. J. Histochem. Cytochem., 1955, 3:61-69.

13. Edward (S.W.) and Ball (E.G.). - The action of phospholipases on succinate oxidase and cytichrome oxidase. J. Biol. Chem., 1954, 209: 619-633.

14. Eranko (O) and Niemi (M.). - Histological and Histochemical observations on the kidney and the liver of rats made hypertensive with desoxycortisosterone and sodium chloride. Acta Path. et Micr. Scand., 1955, 36: 293-303.

16. Findlay (G.H.).- The value of some tetrazolium salts as histochemical reagents for sulfhydryl groups. J. Histochem. Cytochem., 1955, 3: 331-338.

17. Foraker (A.G.) and Denham (S.W.). - Neotetrazolium in determination of succine dehydrogenase activity in the overy. Proc. Soc. Exp. Biol. Med., 1952, 80: 123-134.

 Fred (R.B.) and Knight..... The reduction of 2, 3, 5-triphenyltetrazolium chloride by Pe..... chrysogenium. Science, 1949, 109: 169-170.
 Fried (G.H.) and Zweifach (B.W.). - Neotetrazolium studies of blood vessels.
 Anat. Rec., 1955, 121.....07.

21. Gomori (G.). - Microscopic.....mistry. Principles and Practice. University of Chicago Press, 1952, ......50-153.

22. Gonse (P.H.) and Yotsuganagi (Y.). - Note on intracellular reduction of triphenyltetrazolium in the ..... of sea-urchin. Exper. Cell. Res., 1955, 8: 500-505.

23. Greif (R.L.). - Distribution of .....ic dehydrogenase and hyaluronidase in adult rat testis. Proc. Soc. Exp. Biol. et Med., 1954, 85: 674-677.

24. Harman (J.W.). - Studies on ..... ia: I. The association of cyclophorase with mitochondria. Exp. Cell. Res., 1950, 1: 382-402.

25. Jensen (C.O.), Sacks (W.) and Baldauski. - The reduction of triphenultetrazolium chloride by dehydrogenase of corn embryos. Science, 1951, 113: 65-66.

26. Jerchel (D).-Stufenweise Hyderiumg..... Tetrazoliumsalzen (Tetrazolium Salts) Ang. Chem., 1949, 61: 452.

27. Keilin (D) and Hartree (E.F.) - Activity of the succinic dehydrogenase-cytochrome system in different tissue preparation. Biochem., 1949, 44: 205-218. 23. Kuhn (R.) and Jerchel (D.) - Reduction of Tetrazolium salts through bacteries, (geerende Hefe und kimende Sam.....)....Disch. Chem. Gesellsch., 1941, 74: 949-952.
29. Kun (Z.). - Mechanism of enzymat.c reduction of triphenylterazolium chloride.

Proc. Soc. Exp. Biol et Med., 1951, 77: 195-197.

30. Kun (E) and Abood (L.G.). - Chlorimetric estimation of succinic dehydragenase by triphenyl tetrazolium chloride. Science, 1949, 109: 144-146.

31. Lillie (R.D.). - Histopsthologic technic and Practical Histochemistry. The Blackiston Company, Inc. New York, 1954, 233-236.

32. Lison (L.). - Animal histo'chemistry and cytochemistry. Principles and Methods. Gauthier-Villars, Editeur (=Publishers), Paris, 1953, pages 493-496.

33. Malaty (H.A.) and Bourne (G.H.). - Histochemistry of succinic dehydrogenase. Nature, 1953, 171: 295-297.

34. Martin (S.P.), Cooper (C.D.), Chaudhuri (S.N. and Green (R.). - A tissue factor influencing succinic dehydrogenese activity in tuberculous guinea pigs. J. Exper. Med., 1955, 1011 630-646.

35. Mattson (A.M.), Jensen (C.O.) and Dutcher (R.A.). - Triphenyl tetrazolium chloride as a dye for vital tissues. Science, 1947, 106: 294-295.

36. Mayer (R.K.), McShan (W.H.) and Erway (W.F.). - The succinic dehydrogenase activity of ovarian and lutein tissue. Endocrinology, 1945, 37: 431-436.
37. McMillan (G.C.), Klatzo (L.) and Duff (G.L.). - Histochemical staining reac-

tions of tissues and organs of cholesterol for rabbits. Lab Invest., 1954, 3: 451-468.

38. Mustakallio (K.K.). - Succinic dehydrogenase inhibition in gall bladder epithelium and in liver cells or pregnant mouse. Science, 1954, 119: 881-882.
39. Mustakallio (K.K.). - Histochemical alterations in succinic dehydrogenase activity of guinea pig tissues following administration of diphterie toxin. Exp. Cell. Res., 1954, 7: 592-594.

40. Mustakallio (K.K.) and Jannes (L.). - Intracellular distribution of dehydrogenase activity in C. Diphteriae and in C. Pseudodiphteriticum. Exp. Cell. Res., 1954, 7: 595-596.

41. Novikoff (A.N.). - Histochemical and cytochemical staining methods. In: Analytical Cytology. Methods for studying cellular form and function. The Blackiston Division McGraw-Hill Company, Inc. New York, 1955, p. 242-247.

42. Oriman (R.). - Histochemische Untersuchungen an menschlicher Placenta mit
besonderer Bericksichtigung der Kerakugeln (Kerneinschlusse) under der Plasmalipoidenschlusse. Ztschr. f. Anat. et Entraicklungsgesch., 1955, 119: 28-54.
43. Pearse (A. G. E. ). - Application of the alcaline tetrazolium reaction to the study of reducing groups in tissue section<sup>6</sup>. J. Path. Bact., 1954, 67:129-136. 44. Pearse (A.G.E.). - Histochemistry. Theoretical and Applied. J. and A. Churchill Ltd, London, 1953, p. 297-303 and p. 469-471.

-15-

45. Pearson (B.) and Defendi (V.). - Histochemical demonstration of succinic dehydrogenase in thin tissue sections by means of 2-(p-iodophenyl)-3-(p-nitrophenyl0--5-phenyl tetrazolium chloride under aerobic conditions. J. Histochem. Cytochem., 1954, 2: 248-257.

46. Rennels (E.G.) and Ruskin (A.). - Histochemical changes in succinic dehydragenase activity in rat kidney following administration of mercurial diuretics. Soc. Exper. Biol. Med., 1954, 85Y 309-314.

47. Ried (W.).- Formazane und Tetrazoliumsalze, ihre synthesen und ihre Bedeutung als Reduktionsindikatoren und Vitalfarbstoffe. Angew. Chem., 1952, 64: 391-396.
48. Roberts (L.W.).- Survey of factors responsible for reduction of 2, 3, 5triphenyltetrazolium chloride in plant meristems. Science, 1950. 113: 692-693.
49. Rogers (L.W.). - <u>In vitro</u> reduction of <u>setrazolium</u> indicators by sulfhydryl compounds. Stain. Technol., 1954, 29: 63-67.

50. Rogers (G.E.). - The localization of dehydrogenase activity and sulfhydryl groups in wool and hair follicles by the use of tetrazolium salts. J. Microso. Sc., 1953, 94: 253-268.

51. Rotschild (H.A.) and Barron (E.S.G.). - The oxidation of betaine aldehyde by betaine aldehyde dehydragenase. J. Biol. Chem., 1954, 209: 511-523.

52. Rutenburg (A.M.), Gofstein (R.) and Seligman (A.M.). - Preparation of a new tetrazolium salt which yields a blue pigment on reduction and in use in the demonstration of enzymes in normal and neoplastic tissues. Cancer Res., 1950, 10: 113-121.

53. Rutenburg (A.M.), Wolman (M.) and Seligman (A.). - Comparative distribution of succinic dehydragenase in six mammals and modification in the histochemical technic. J. Histochem. Cytochem., 1953, 1: 66-81.

54. Seligman (A.M.), Gofstein (F.) and Rutenburg (A.M.). - Preparation of a radioactive iodotetrazolium salt and its distribution in mice. Cancer. Res., 1949, 9: 366.

55. Seligman (A.M.) and Rutenburg (A.M.). - The histochemical demonstration of succinic dehydrogenase. Science, 1951, 113: 317-320.

56. Shelton (E.) and Schneider (W.C.). - On the usefulness of tetrazolium salts as histochemical indicators of dehydrogenese activity. Anat. Sec., 1952, 112:61-81.

57. Schneider (W.C.) and Potter (V.R.). - Succinic dehydrogenese and cytochrome oxidase, J. Biol. Chem., 1943, 149: 217-227.

58. Smith (E.E.). - Tetrazolium salt. Science, 1951, 113: 751-754.

59. Sonneblick (B.P.), Antopol (W.) and Goldman (L.). - Influence of tetrazolium salt on the growth and cytology of root tips. Trans. N.Y. Acad. Sci., 1950, 12: 161-163.

60. Steigleder (K.). - Reduzierende Substanzen in der normalen Meuschenhaut und in der normalen und verbreiterten Haut der Ratte. Arch. Dermat. et Syphil., 1955, 199: 304-400.

61. Steigleder (G.K.). - Zur Funktion der Acanthose. Arch. Dermat. et Syphil., 1955, 200: 377-395.

62. Strominger (J.L.) and Lowry (O.H.). - The quantitative histochemistry of brain.
IV. Lactic, malic and glutanic dehydrogenases. J. Biol. Chem., 1955, 213: 633-646.
63. Villareal (r) and Burgos (M.H.). - A correlated biochemical and histochemical study of succinic dehydrogenase activity in the gastric mucosa of the rat and frog. J. Cell. et Comp. Physiol., 1955, 45: 327-330.

64. Zweifach (B.W.), Black (M.M.) and Schorr (S.). - Evaluation of tetrazolium as a histochemical index of adrenal cortical activity. Proc. Soc. Exper. Biol. et Med., 1951, 76: 446-454.