



AFRL-SA-WP-SR-2017-0027

Relevancy of Serum Calcium in Predicting Blood Product Transfusion in Trauma



**Lt Col Rance Bryan, MOANG, NC; Kathryn Lindsay, RN;
Lt Col Bradley Putty, MOANG, MC; Maj Jason Grimm,
USAF, NC; Carl Freeman, MD**

Air Force Expeditionary Medical Skills Institute, C-STARS St. Louis,
St. Louis University



August 2017

**DISTRIBUTION STATEMENT A. Approved
for public release. Distribution is unlimited.**

STINFO COPY

**Air Force Research Laboratory
711th Human Performance Wing
U.S. Air Force School of Aerospace Medicine
International Expeditionary Education &
Training Department
2510 Fifth St., Bldg. 840
Wright-Patterson AFB, OH 45433-7913**

NOTICE AND SIGNATURE PAGE

Using Government drawings, specifications, or other data included in this document for any purpose other than Government procurement does not in any way obligate the U.S. Government. The fact that the Government formulated or supplied the drawings, specifications, or other data does not license the holder or any other person or corporation or convey any rights or permission to manufacture, use, or sell any patented invention that may relate to them.

Qualified requestors may obtain copies of this report from the Defense Technical Information Center (DTIC) (<http://www.dtic.mil>).

AFRL-SA-WP-SR-2017-0027 HAS BEEN REVIEWED AND IS APPROVED FOR PUBLICATION IN ACCORDANCE WITH ASSIGNED DISTRIBUTION STATEMENT.

//SIGNATURE//

COL BARBARA A. JONES
Chair, Int'l Exped Educ & Training Dept

//SIGNATURE//

DR. RICHARD A. HERSACK
Chair, Aeromedical Research Department

This report is published in the interest of scientific and technical information exchange, and its publication does not constitute the Government's approval or disapproval of its ideas or findings.

REPORT DOCUMENTATION PAGE				<i>Form Approved</i> <i>OMB No. 0704-0188</i>	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE (DD-MM-YYYY) 10 Aug 2017		2. REPORT TYPE Special Report		3. DATES COVERED (From – To) September 2014 – March 2016	
4. TITLE AND SUBTITLE Relevancy of Serum Calcium in Predicting Blood Product Transfusion in Trauma				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER FA8650-14-2-6-C02	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Lt Col Rance Bryan, Kathryn Lindsay, Lt Col Bradley Putty, Maj Jason Grimm, Dr. Carl Freeman				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) USAF School of Aerospace Medicine International Expeditionary Education & Training Dept/ETS 2510 Fifth St., Bldg. 840 Wright-Patterson AFB, OH 45433-7913				8. PERFORMING ORGANIZATION REPORT NUMBER AFRL-SA-WP-SR-2017-0027	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSORING/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT DISTRIBUTION STATEMENT A. Approved for public release. Distribution is unlimited.					
13. SUPPLEMENTARY NOTES Cleared, 88PA, Case # 2017-5643, 9 Nov 2017.					
14. ABSTRACT Uncontrolled hemorrhage is the leading cause of preventable mortality in trauma. Rapid identification of patients requiring blood transfusion during assessment and resuscitation is necessary for mortality reduction. Most pre-hospital or field medical criteria used to predict blood product needs in trauma patients rely on a combination of physiological, anatomic, and mechanism-of-injury components. Calcium (Ca) activates protein kinase C and is required for coagulation factors to bind properly in the clotting cascade. Ca has presented a novel laboratory test to predict early blood transfusion. The purpose of this study was to identify a relationship between serum Ca levels and the need for early transfusion. In this retrospective analysis, serum Ca <7.9 mg/dL was strongly associated with the need for blood product transfusion, as well as transfusion of more than four units within 4 hours, even after controlling for other clinical variables. This effect was age specific for the subject group aged 40 years and below. Patients with normal blood pressure could give medical teams a false sense of patient stability. Younger patients have a compensating capacity with blood loss and will not show signs of shock until they have exhausted this mechanism.					
15. SUBJECT TERMS Trauma resuscitation, blunt trauma, penetrating trauma, hypotension, anemia, calcium, transfusion, coagulopathy					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT SAR	18. NUMBER OF PAGES 11	19a. NAME OF RESPONSIBLE PERSON Lt Col Rance A. Bryan
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U			19b. TELEPHONE NUMBER (include area code)

This page intentionally left blank.

TABLE OF CONTENTS

Section	Page
LIST OF TABLES	ii
1.0 SUMMARY	1
2.0 INTRODUCTION	1
3.0 BACKGROUND	1
4.0 METHODS	2
5.0 RESULTS	2
6.0 DISCUSSION	4
7.0 CONCLUSIONS.....	4
8.0 REFERENCES	4
LIST OF ABBREVIATIONS AND ACRONYMS	5

LIST OF TABLES

	Page
Table 1. Early Blood Transfusion Following Trauma.....	3
Table 2. Early Large Volume Transfusion Following Trauma.....	Error! Bookmark not defined.
Table 3. Predictors of Early Blood Transfusion Following Trauma	Error! Bookmark not defined.

1.0 SUMMARY

In this retrospective analysis, serum calcium <7.9 mg/dL was strongly associated with the need for blood product transfusion, as well as transfusion of more than four units within 4 hours, even after controlling for other clinical variables. This effect was age specific for the subject group aged 40 years and below. Younger patients have a compensating capacity with blood loss and will not show signs of shock until they have exhausted this mechanism.

2.0 INTRODUCTION

Uncontrolled hemorrhage is the leading cause of preventable mortality in trauma [1]. Rapid identification of patients requiring blood transfusion during assessment and resuscitation is necessary for mortality reduction [2]. Most pre-hospital or field medical criteria used to predict blood product needs in trauma patients rely on a combination of physiological, anatomic, and mechanism-of-injury components. Calcium (Ca) activates protein kinase C and is required for coagulation factors to bind properly in the clotting cascade. Ca has presented a novel laboratory test to predict early blood transfusion. The purpose of this study was to identify a relationship between serum calcium (sCa) levels and the need for early transfusion.

3.0 BACKGROUND

In the current conflicts, the most common causes of death in combat casualties have been head injury followed by truncal exsanguinations, and major hemorrhage remains the most frequent avoidable cause of early mortality in combat casualties [3]. Massively bleeding casualties die fast. Even in the civilian setting, those who survive at least 15 minutes into advanced trauma care go on to die of bleeding and half are dead within 2 hours. The most important advance in trauma transfusion medicine in the last decade has been the recognition of the acute coagulopathy of trauma and the need to supply coagulation factors (as plasma) and platelets in near-equivalence with red cells and do so quickly [4]. However, although large numbers of units of packed red blood cells (pRBCs) can be provided immediately upon arrival, equivalent proportions of platelets and plasma typically require a minimum of 20 minutes lead time to be thawed or otherwise processed and to arrive from the blood bank. In the military setting, where fresh whole blood may be available as a “walking blood bank,” appropriate donors need to be identified, prepped, and drawn. Advanced recognition of the need for blood transfusion from knowing the sCa levels can change outcomes and save lives.

Most pre-hospital, field medical criteria used to predict outcome and the need for interventions in trauma patients rely on a combination of physiological, anatomic, and mechanism-of-injury components that, in the experienced trauma clinician, are summed up as good clinical judgment. However, this approach still fails to identify a number of bleeding patients with severe injuries [3]. Civilian paramedic judgment has been assessed objectively and shown to provide little support in the diagnosis of major trauma, mainly due to differences across emergency medical and trauma care systems, individual experience levels, and incompatible methods of data collection. Patients with normal blood pressure could give medical teams a false sense of patient stability [5]. Despite the use of improving physiologic monitoring systems and better training to identify cardiorespiratory instability, in the civilian trauma system, rates of over-triage and under-triage remain high [3,4]. Likewise, once casualties reach advanced care,

the number of individuals who receive one to four units of uncrossmatched group O pRBCs in the trauma center is surprisingly high, apparently based on the trauma team's assessment of the need for immediate transfusion. However, these casualties neither receive nor apparently require any other blood products in the first 24 hours and have good outcomes, raising the issue of whether they needed the blood—or the risk of transfusion—in the first place.

4.0 METHODS

An Institutional Review Board-approved retrospective review of adult subjects 18-65 years of age and presenting as level 1 trauma activations to an urban level 1 trauma center over a 48-month period was conducted. Deaths in the emergency department were excluded, as were subjects with labs drawn more than 60 minutes following arrival. Patient demographics, laboratory tests performed within 1 hour of arrival (including hemoglobin (HGB), international normalized ratio (INR), base excess (BE), and sCa), transfusion (including administration time), and outcome data were analyzed, using Student's t-test and chi-square analysis. Early transfusion was defined as blood product transfusion within 4 hours of arrival. Significant early transfusion was defined as at least four units of pRBCs transfused within 4 hours. Factors with $p < 0.1$ on univariate analysis were entered into logistic regression analysis, with $p < 0.05$ considered significant.

5.0 RESULTS

After exclusions, 488 subjects met criteria for analysis. Median age was 28.5 years (range: 18-65). Approximately half of the subjects presented with a penetrating mechanism of injury ($n=253$, 51.8%), and 207 (42.4%) subjects required red blood cell transfusions. There were no significant differences in age, but subjects receiving early transfusion presented with lower Revised Trauma Score (RTS), increased INR, decreased BE, and lower Ca (Table 1). On multiple logistic regression, early (<4 hours) transfusion was associated with systolic blood pressure (SBP) <110 mmHg (odds ratio (OR) 4.0, confidence interval (CI) 1.5-11.3, $p < 0.01$) and hypocalcemia ($\text{Ca} < 7.9$: OR 3.4, CI 1.2-9.6, $p = 0.02$) (Table 2). Furthermore, transfusion of >4 units within 4 hours remained associated with hypocalcemia (OR 3.0, CI 1.4-6.0, $p = 0.001$) (Table 3).

In subjects 40 years old or less who had $\text{sCa} < 8.0$ and $\text{INR} > 1.1$ ($N=84$), 77 (92%) received blood products. Fifty out of these 77 subjects (65%) had SBP >109 mmHg at time of arrival (TOA). Nineteen out of the total 84 subjects (23%) received massive blood product transfusion from TOA to 1 hour, and of these, 9 had SBP >109 mmHg (47%) at TOA.

Table 1. Early Blood Transfusion Following Trauma

	pRBCs within 4 h (n=99)	No Early Transfusion (n=369)	p-value
Mean Age (yr)	34 (23,48)	31 (24, 44)	0.13
Mean RTS	6.4 (3, 7.8)	7.8 (6, 8)	0.007
Mean GCS	13 (3, 15)	15 (9.5, 15)	<0.001
Mean SBP (mmHg)	108 (82, 133)	134 (116, 146)	<0.001
Mean HGB (g/dL)	11.6 (9.9, 13)	13 (12, 14)	<0.001
Mean INR	1.1 (1, 1.8)	1 (1, 1)	<0.001
Mean BE (mmol/L)	-9 (-13.3, -4.3)	-5 (-8.8, -3)	0.005
Mean Ca (mg/dL)	7.2 (6.7, 8)	8 (7.9, 8.8)	<0.001

Note: >1 unit pRBCs in first 4 h.

GCS = Glasgow Coma Scale.

Table 2. Predictors of Early Blood Transfusion Following Trauma

Factor	Univariate OR (95% CI)	p-value	Adjusted OR (95% CI)	p-value
Hypotension ^a	6.35 (4.0, 10.2)	<0.001	4.07 (1.46, 11.35)	0.007
Anemia ^b	4.2 (2.7, 6.6)	<0.001	1.97 (1.46, 11.35)	0.218
INR > 1.0	4.8 (3.0, 7.6)	<0.001	1.05 (0.35, 3.12)	0.938
BE < -6	4.4 (1.9, 10.3)	0.001	2.49 (0.90, 6.88)	0.079
Hypocalcemia ^c	6.8 (4.2, 10.9)	<0.001	3.38 (1.18, 9.63)	0.022

^aHypotension defined as SBP <110 mmHg.

^bAnemia: HGB <12 g/dL.

^cHypocalcemia: sCa <7.9.

Table 3. Early Large Volume Transfusion Following Trauma

	≥4 Units pRBCs within 4 h (n=66)	<4 units pRBCs within 4 h (n=402)	p-value
Mean Age (yr)	34 (22, 47)	31 (24, 44)	0.12
Mean RTS	5 (2.9, 7.8)	7.8 (6, 8)	0.007
Mean GCS	10.5 (3, 15)	15 (8, 15)	<0.001
Mean SBP (mmHg)	98 (85, 121)	134 (116, 146)	<0.001
Mean HGB (g/dL)	11 (8.5, 12.4)	13 (11.9, 14)	<0.001
Mean INR	1.3 (1, 2)	1 (1, 1)	<0.001
Mean BE (mmol/L)	-12.9 (-17.6, -6.1)	-5 (-9, -3)	0.007
Mean Ca (mg/dL)	7.2 (6.7, 8)	8 (7.9, 8.7)	<0.001

Note: >4 units pRBCs in first 4 h.

6.0 DISCUSSION

The results of this study will increase the knowledge of early indicators of blood product needs and conservation of resources through decreased mortality and will enhance accurate blood product delivery.

7.0 CONCLUSIONS

In this retrospective analysis, $sCa < 7.9$ was strongly associated with the need for red blood cell transfusion, as well as transfusion of more than four units within 4 hours, even after controlling for other clinical variables. These results could be a significant tool to aid in the prioritization for evacuation to a higher level of care before deterioration. Further prospective analysis, for validation as well as the role of Ca replacement in hemostasis, is warranted.

8.0 REFERENCES

1. Maegele M, Gu ZT, Huang QB, Yang H. Updated concepts on the pathophysiology and the clinical management of trauma hemorrhage and coagulopathy. *Chin J Traumatol.* 2017; 20(3):125-132.
2. Vandromme MJ, Griffin RL, McGwin G Jr, Weinberg JA, Rue LW 3rd, Kerby JD. Prospective identification of patients at risk for massive transfusion: an imprecise endeavor. *Am Surg.* 2011; 77(2):155-161.
3. Callcut RA, Cotton BA, Muskat P, Fox EE, Wade CE, et al. Defining when to initiate massive transfusion: a validation study of individual massive transfusion triggers in PROMMTT patients. *J Trauma Acute Care Surg.* 2013; 74(1):59-65, 67-68, discussion 66-67.
4. Dente CJ, Shaz BH, Nicholas JM, Harris RS, Wyrzykowski AD, et al. Early predictors of massive transfusion in patients sustaining torso gunshot wounds in a civilian level I trauma center. *J Trauma.* 2010; 68(2):298-304.
5. Thorson CM, Van Haren RM, Ryan ML, Pereira R, Olloqui J, et al. Admission hematocrit and transfusion requirements after trauma. *J Am Coll Surg.* 2013; 216(1):65-73.

LIST OF ABBREVIATIONS AND ACRONYMS

BE	base excess
Ca	calcium
CI	confidence interval
GCS	Glasgow Coma Scale
HGB	hemoglobin
INR	international normalized ratio
OR	odds ratio
pRBC	packed red blood cells
RTS	Revised Trauma Score
SBP	systolic blood pressure
sCa	serum calcium
TOA	time of arrival