



**US Army Corps
of Engineers®**
Engineer Research and
Development Center

ERDC
INNOVATIVE SOLUTIONS
for a safer, better world

Climate Change Vulnerability of Army Installations Attributable to Listed and At-Risk Species

Matthew G. Hohmann, David K. Delaney, and Wade A. Wall

July 2017



Golden blazing star
(*Nuttallia chrysantha*)

Kirtland's snake
(*Clonophis kirtlandii*)



Northern long-eared bat
(*Myotis septentrionalis*)

Representative species from Fort Carson, CO and Camp Atterbury, IN, which have the highest estimated climate change exposure among installations estimated to have high climate change vulnerability

The U.S. Army Engineer Research and Development Center (ERDC) solves the nation's toughest engineering and environmental challenges. ERDC develops innovative solutions in civil and military engineering, geospatial sciences, water resources, and environmental sciences for the Army, the Department of Defense, civilian agencies, and our nation's public good. Find out more at www.erdclibrary.usace.army.mil.

To search for other technical reports published by ERDC, visit the ERDC online library at <http://acwc.sdp.sirsi.net/client/default>.

Climate Change Vulnerability of Army Installations Attributable to Listed and At-Risk Species

Matthew G. Hohmann, David K. Delaney, and Wade A. Wall

*U.S. Army Engineer Research and Development Center (ERDC)
Construction Engineering Research Laboratory (CERL)
2902 Newmark Dr.
Champaign, IL 61826*

Final Report

Approved for public release; distribution is unlimited.

Prepared for Office of the Assistant Secretary of the Army for Acquisition,
Logistics, and Technology, Washington, DC 20310-0103

Under Program Element 622720A896, Work Unit F94KF0,
“Maneuver Area Capacity”

Abstract

Climate change is an important emerging concern for the U.S. Department of Defense (DoD) and the Army. Key among the issues is climate change-driven increases in the number of species listed under the U.S. Endangered Species Act (ESA) and stress to Federally listed species already listed, because both pose additional management requirements and challenges, as well as potential restrictions on training land use. This work developed an approach for characterizing this component of installation climate change vulnerability that integrates multiple factors related to exposure, sensitivity, adaptive capacity, and number of listed and at-risk species. The approach was applied to Army installations in the Continental United States that have Integrated Training Area Management (ITAM) programs. Additionally, the assessment was used to rank installations across the Army based on the aggregate vulnerabilities of species. The approach to vulnerability assessment demonstrated here is suitable for evaluating whether climate change-driven impacts to listed and at-risk species is likely to affect installation resilience.

DISCLAIMER: The contents of this report are not to be used for advertising, publication, or promotional purposes. Citation of trade names does not constitute an official endorsement or approval of the use of such commercial products. All product names and trademarks cited are the property of their respective owners. The findings of this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

DESTROY THIS REPORT WHEN NO LONGER NEEDED. DO NOT RETURN IT TO THE ORIGINATOR.

Contents

Abstract	ii
Tables and Figure.....	iv
Preface.....	v
1 Introduction.....	1
1.1 Background.....	1
1.2 Objectives.....	4
1.3 Approach	4
1.4 Scope.....	4
2 Methods	5
2.1 Climate change vulnerability assessment.....	5
2.2 Direct and indirect climate change exposure	7
2.3 Climate change sensitivity.....	8
2.4 Adaptive capacity.....	9
2.5 Listed and at-risk species.....	11
2.6 Installation ranks	12
2.7 Calculating climate change vulnerability scores for installations	12
2.8 Examining differences in installation climate change vulnerability factors and scores across USFWS regions	13
3 Results and Discussion.....	15
3.1 Direct and indirect exposure	15
3.2 Sensitivity	20
3.3 Adaptive capacity.....	21
3.4 Listed and at-risk species.....	23
3.5 Installation ranks	24
3.6 Threatened and endangered species related climate change vulnerabilities of installations	24
4 Conclusions and Recommendations	26
4.1 Conclusions.....	26
4.2 Recommendations.....	26
Appendix A: Installation Subfactor Values	27
Appendix B: Lists of Threatened, Endangered, and At-Risk Species by Installation	31
Appendix C: Installation Vulnerabilities Calculated Without the Installation Rank Factor	44
Acronyms and Abbreviations.....	50
References	46
Report Documentation Page (SF 298)	52

Tables and Figure

Tables

1	Factors and subfactors used to assess climate change vulnerability.....	6
2	Probability of Federal listing by taxonomic group used to weight species at-risk. Numbers in parentheses represent number of species used to develop the model	12
3	Weights multiplied by the five factors to calculate climate change vulnerabilities of Army ITAM installations	13
4	USFWS regions and associated states.....	14
5	Weighted scores for five factors used to estimate climate change vulnerability of 43 CONUS Army ITAM installations. Installations are ordered high-to-low based on vulnerability.....	15
6	Mean scores and ranges (in parentheses) for five climate change factors and the overall vulnerability of 43 CONUS Army ITAM installations within seven USFWS regions. Means designated by the same letter within columns do not differ at $P = 0.05$ by Tukey's Honestly Significant Difference in an across-region comparison.....	17

Figure

1	Conceptual diagram of combinations of climate change exposure, sensitivity, and adaptive capacity that identify seven classes of vulnerability having different implications for prioritization and strategic planning	18
---	--	----

Preface

This study was conducted for Office of the Assistant Secretary of the Army for Acquisition, Logistics, and Technology, ASA(ALT) under program element 622720A896, “Army Environmental Quality Technology”; Work Unit F94KF0, “Maneuver Area Capacity.” The technical monitor was Steve Sekscienski, OACSIM-ISE.

This work was conducted by the Ecological Processes Branch (CNN), Installations Division (CN), Engineer Research and Development Center, Construction Engineering Research Laboratory (ERDC-CERL). Dr. Chris Rewerts was Chief, CEERD-CNN, and Michelle J. Hanson was Chief, CEERD-CN. The associated Technical Director was Alan Anderson, CEERD-CZT. The Deputy Director of ERDC-CERL was Dr. Kirankumar Topudurti and the Director was Dr. Ilker Adiguzel.

COL Bryan S. Green was Commander of ERDC, and Dr. David W. Pittman was the Director.

THIS PAGE INTENTIONALLY LEFT BLANK

1 Introduction

1.1 Background

Federal land management agencies are faced with the challenge of managing numerous threatened, endangered, and at-risk species (i.e., taxa assessed by NatureServe as critically imperiled [G1/T1] or imperiled [G2/T2]) (NatureServe 2011), a challenge that will be exacerbated by climate change (Dawson et al. 2011, Ellenwood et al. 2012, Shaw et al. 2021, Urban 2015). The U.S. Department of Defense (DoD), which is the fifth largest land management agency in the United States, manages over 12 million hectares of land on more than 425 military installations (Stein et al. 2008). Although this represents less than 5% of the land managed by the four larger agencies combined, 23% of all U.S. Endangered Species Act (ESA) status (endangered, threatened, candidate, or proposed species) and 15% of all at-risk species occur on DoD lands (Stein et al. 2008).

Additionally, more of these DoD ESA status (270; ACSIM 2010) and at-risk species (220; NatureServe 2011) occur on Army lands than occur on all other DoD services combined (Stein et al. 2008). Given the recent U.S. Fish and Wildlife Service (USFWS) lawsuit settlement requiring review of 757 species proposed for listing under the ESA by 2018 (ESA Section 4 Deadline Litigation, Case Number 2165, U.S. District Court for the District of Columbia), a substantive number (ca. 230) of additional listed species are anticipated to occur on/near Army and Army National Guard installations (Sperry et al. 2016). The currently large and increasing Army conservation responsibility related to Federally listed and at-risk species highlights the importance of proactively conducting climate change vulnerability assessments.

Executive Order (EO) 13514, *Federal Leadership in Environmental, Energy and Economic Performance* (White House 2009) required Federal agencies to address climate change risks and vulnerabilities in short- and long-term planning. DoD first acknowledged the potential impacts of climate change on its facilities, infrastructure, military capabilities, and training and testing activities within the 2010 Quadrennial Defense Review (DoD 2010). Subsequently, DoD issued Department of Defense Instruction (DODI) 4715.03 (DoD 2011) pertaining to its Natural Resources

Conservation Program, which requires DoD installations to: (1) address climate change in their Integrated Natural Resources Management Plans (INRMPs), (2) use the best available science to assess potential climate change impacts, and (3) use adaptive strategies to address those impacts. In 2013, this Instruction was followed by Department of Defense Manual (DODM) 4715.03, the *Integrated Natural Resources Management Plan Implementation Manual* (DoD 2013), which outlined procedures for preparing, reviewing, updating, and implementing INRMPs in accordance with Department of Defense Instruction (DODI) 4715.03.

In response to EO 13514, DoD also developed a Climate Change Adaptation Roadmap (CCAR) in 2012 (DoD 2012). The CCAR fulfilled a requirement to include an adaptation planning document as an appendix to the Department's annual Strategic Sustainability Performance Plan. The CCAR established broad climate change adaptation goals to: (1) develop a coordinating committee to address climate change, (2) use the best available science to inform decisions, (3) integrate climate change information into existing processes, and (4) encourage partnerships with other agencies (DoD 2012). Given that impacts to DoD are expected to vary by region, assessing vulnerability to climate change is a large component of the CCAR framework. The CCAR included a detailed table highlighting specific climate change phenomena, potential impacts, and potential mission vulnerabilities. Among the listed potential impacts are stress to protected species and an increase in the number of species at risk.

In 2013, EO 13653, *Preparing the United States for the Impacts of Climate Change* (White House 2013), charged DoD and other Federal agencies to:

complete an inventory and assessment of proposed and completed changes to their land- and water-related policies, programs, and regulations necessary to make the Nation's watersheds, natural resources, and ecosystems, and the communities and economies that depend on them, more resilient in the face of a changing climate.

It also charged DoD and other Federal agencies to "develop and provide authoritative, easily accessible, usable, and timely data, information, and decision-support tools on climate preparedness and resilience." EO 13653 also mandated regular updates to and implementation of the agency adaptation plans required under EO 13514.

In 2014, DoD updated its CCAR in response to requirements set out in EO 13653 (DoD 2014a). In alignment with EO 13653, the updated CCAR established three broad adaptation goals: (1) to identify and assess the effect of climate change on the Department, (2) to integrate climate change considerations across the Department and manage associated risks, (3) to collaborate with internal and external stakeholders on climate change challenges. Each of these goals was evaluated in relation to DoD's plans and operations, training and testing, built and natural infrastructure, and acquisition and supply chain. Additionally, a summary of the potential impacts to DoD's mission was provided within Annex 2 of the 2014 CCAR. Similar to the 2012 roadmap, the 2014 CCAR explicitly identified concerns about climate change-related stress to currently listed threatened and endangered species on and adjacent to DoD installations. Increased numbers of listed and at-risk species and associated management requirements and challenges were also identified as concerns.

The Army needs a means of identifying the impact of future climate change on installations, to generate information about long-term sustainability that can be used to make diverse decisions, such as training mission assignments, base realignment and closure (BRAC), conservation funding investments, and alternative mitigation strategy selection. Wilhoit et al. (2015) proposed a simple method of ranking installations based on potential impacts of climate change-related listed and at-risk species management on training land use. However, their approach did not include any specific information about climate change vulnerability, but instead used the number of species on installations, the conservation status of these species (i.e., Federally listed, proposed for listing, and at-risk), listing probabilities of different taxonomic groups, and installation area. Effectively, their ranking metric calculated a status and probability of listing-adjusted estimate of at-risk and listed species density.

In BRAC studies, the Army ranks installations based on their characteristics. For example, since installation training lands provide the Army with critical mission capabilities, installations with the largest training lands are considered high value. Anything that erodes the size of the available training lands or the use of those lands, negatively impacts the installation's value and is of interest. Climate change is one phenomenon that can erode multiple installation characteristics and thus the installation's resulting value to the Army.

This effort adopted an expanded approach for characterizing the anticipated relative increase in climate change-related conservation responsibilities among installations, one that specifically incorporates climate change vulnerability.

1.2 Objectives

The objectives of this effort were to:

1. Develop an approach for assessing installation climate change vulnerabilities due to listed and at-risk species
2. Implement this climate change vulnerability assessment for Continental United States (CONUS) Army installations included within the Integrated Training Area Management (ITAM) program
3. Use the estimated vulnerabilities to rank installations across the Army
4. Explore the variable contribution of the five factors used to estimate climate change vulnerability
5. Make recommendations about mitigation strategies.

1.3 Approach

The objectives of this work were accomplished in five primary tasks:

1. The listed and at-risk species occurring on each Army installation were identified.
2. Spatial and aspatial datasets for five factors used to assess the climate change vulnerability of species were developed.
3. The developed assessment approach was implemented.
4. Climate change vulnerabilities across installations and regions were compared.
5. Recommendations about suitable mitigation strategies based on the magnitudes of the evaluated factors were summarized.

1.4 Scope

The climate change vulnerability assessments within this effort focused on CONUS installations and species that have been either Federally listed as threatened or endangered, or that have been identified to be at-risk (NatureServe 2014). Assessments were made using climate change projection data for the year 2050.

2 Methods

Climate change vulnerability is generally described as a function of exposure, sensitivity, and adaptive capacity (Schneider et al. 2007, Williams et al. 2008). Exposure is characterized by the magnitude and rate of climate change a region is expected to experience. Different portions of the United States are projected to vary in their direct and indirect exposure to climate change. Sensitivity to climate change is characterized by the degree to which a species or region is anticipated to be affected by climate variables (e.g., drought, extreme temperatures). Regions where substantial annual or interannual climate variability occurs are anticipated to be less susceptible to climate change. Adaptive capacity refers to the ability of species to adapt to climate change and to the availability of mitigation options (e.g., conservation partners and natural landscapes) that might lessen climate change impacts in a region.

2.1 Climate change vulnerability assessment

Multiple approaches for estimating species' and regional climate change vulnerabilities have been proposed and adopted (e.g., USEPA National Center for Environmental Assessment framework, USEPA 2009; NatureServe's Climate Change Vulnerability Index, Young et al. 2011, 2015; Bagne et al. 2011). This effort set out to develop an approach for quantifying the degree to which Army installations may be affected by climate change as a result of increased vulnerability of listed and at-risk species, which can restrict training and testing land access due to requirements of the ESA. Specifically, this work sought an approach that was objective, uncomplicated, transparent, and applicable across the Continental United States, and that could be estimated using available spatial datasets. It is also important to examine multiple components of risk in combination when assessing the potential impacts of climate change on species of concern (see Hinkel 2010, Dickinson et al. 2014). The approach ultimately adopted was based on a simple weighted sum of five factors that characterized installations': (1) direct and indirect climate change exposure, (2) likely sensitivity to climate change exposure, (3) regional adaptive capacity, (4) potential vulnerability to listed and at-risk species, and (5) importance to the Army's overall training mission (Table 1).

Each of these five factors were summarized using one or more standardized subfactors, weighted by a multiplier, and then summed to arrive at an overall index of installation vulnerability. Table 1 lists descriptions of these five factors. The index can be used to assess the relative vulnerability of installations, identify the most important factors affecting vulnerability, identify the conservation partnering and adaptation strategies likely available, and promote coordination and consistency in adaptation planning and management across Army.

Table 1. Factors and subfactors used to assess climate change vulnerability.

Factors	Subfactors	Description
1. Direct and Indirect Exposure	1.1 Temperature Change	Mean predicted change in annual temperature by 2050, calculated across installation landscapes
	1.2 Moisture Change	Mean predicted net change in moisture based on the Hamon AET:PET (actual evapotranspiration potential evapotranspiration) Moisture Metric, calculated across installation landscapes
	1.3 Sea Level Rise	Predicted increase in sea level and consequent influence of storm surges calculated across installation landscapes
2. Sensitivity	2.1. Historical Thermal Variation	Mean seasonal temperature variation (difference between the highest mean monthly maximum temperature and lowest mean monthly minimum temperature from 1951-2006) calculated across installation landscapes. Species exposed to low seasonal temperature variation are expected to be more vulnerable than species exposed to high seasonal temperature variation
	2.2. Historical Precipitation	Mean annual precipitation from 1951-2006 calculated across installation landscapes. Species in arid climates are expected to be more vulnerable to reductions in precipitation than species in mesic climates
3. Adaptive Capacity	3.1 Anthropogenic Barriers	Anthropogenically altered landscapes (e.g., urban or agricultural areas) may hinder the dispersal of species, estimated as percent developed landcover within a 50 km buffer on installations
	3.2 Protected Lands	Federal lands offer listed and at-risk species the most comprehensive protections and represent potential conservation partners, estimated as percent Federal lands within a 50 km buffer on installations
4. Listed and At-Risk Species	4.1 Listed and At-risk Species	Index of the current and potential listed and at-risk species conservation burden
	4.2 Density of Listed and At-risk Species	The index described in 4.1 divided by installation area
5. Installation Rank		ITAM rank reflecting installation importance to the Army's training and testing missions

2.2 Direct and indirect climate change exposure

Direct exposure of installations and their associated listed and at-risk species to future climate change was evaluated in terms of two subfactors: change in average annual temperature and moisture availability projected for the year 2050 (Table 1). Climate data were sourced from the ClimateWizard team (www.climatewizard.org) and NatureServe (www.natureserve.org/conservation-tools/climate-change-vulnerability-index). Climate and all other spatial data were projected to Alber's Equal Area Conic. Climate data specific to Army installations were extracted from these spatial layers based on installation boundary information sourced from the U.S. Geological Survey (USGS) National Map (http://nationalmap.gov/small_scale/mld/fedlanp.html) and using ArcGIS 10.2.2 (ESRI 2016). A similar process was applied for all other spatially explicit subfactors (Sections 2.3 and 2.4).

The average temperature changes projected across installation assessment areas (Table 1, Subfactor 1.1) ranged from 1.95 to 3.21 °C (mean = 2.67 °C) for Joint Base Lewis-McCord and Camp Dodge Joint Maneuver Training Center, respectively (Appendix A). These values were then rescaled (i.e., standardized). A score range procedure (Eastman et al. 1993, Malczewski 2000) was applied:

$$x_i = (R_i - R_{\min}) / (R_{\max} - R_{\min}) \quad (2-1)$$

where:

R_i represents the observed values

R_{\min} and R_{\max} are the range of observed values

x_i are the standardized, dimensionless values on a scale of 0 to 1, with higher values representing greater relevance for climate change vulnerability.

Projected changes in moisture availability (Table 1, Subfactor 1.2) within installation assessment areas were characterized using the Hamon AET:PET moisture metric (Hamon 1961), as prepared by the ClimateWizard team. This metric integrates precipitation and temperature through a ratio of actual evapotranspiration (AET) to potential evapotranspiration (PET), with consideration of total daylight hours and saturated vapor pressure. Average projected changes in Hamon AET:PET moisture metric across installation assessment areas ranged from -0.010 to -0.120% (mean

= -0.068%), for Yuma Proving Ground, AZ and Fort Sill, OK, respectively (Appendix A). Values for this subfactor were standardized with a score range procedure. Resulting values ranged from zero to one, with higher values having greater relevance.

Species and installations may also be impacted by climate change indirectly. Indirect exposure to climate change was assessed via a sea level change subfactor (Table 1, 1.3) that used coastal vulnerability index data acquired from the USGS Coastal Change Hazard Portal (<http://marine.usgs.gov/coastalchangehazardsportal>). The index ranks relative susceptibility of U.S. coasts to sea level rise within four categories (low, moderate, high, and very high) using information on geomorphology, regional coastal slope, tide range, wave height, relative sea level rise, and shoreline erosion and accretion rates (Thieler and Hammar-Klose 2000). The percentages of the installation assessment areas represented by any of the three highest index categories (i.e., moderate, high, and very high) were estimated within ArcGIS 10.2.2. Only two installations were found to be indirectly impacted by sea level change, with Fort Eustis, VA (14.14%) more negatively impacted than Aberdeen Proving Ground, MD (5.5%) (Appendix A). Values for this subfactor were then standardized with a score range procedure. Resulting values ranged from zero to one, with higher values having greater relevance.

2.3 Climate change sensitivity

Populations of listed and at-risk species in arid regions are thought to be more vulnerable to projected reductions in precipitation than those residing in mesic regions (e.g., Vale and Brito 2015). Furthermore, populations of listed and at-risk species that have historically experienced little temperature variation may be less able to tolerate projected increases in temperature (Tomanek 2008). Thus, climate change sensitivity has been assessed by evaluating historical precipitation and seasonal temperature variation (e.g., Young et al. 2015). In this effort, sensitivity of installations and their associated listed and at-risk species to future climate change was evaluated in terms of two subfactors: historical precipitation (Table 1, 2.2) and seasonal temperature variation (Table 1, 2.1). Mean annual precipitation and seasonal temperature variation (i.e., difference between the highest mean monthly maximum and lowest mean monthly minimum) from

1951-2006 were estimated within installation assessment areas using climate data sourced from the Climate Wizard (www.climatewizard.org) and NatureServe (www.natureserve.org/conservation-tools/climate-change-vulnerability-index).

Historical seasonal temperature variation ranged from 15.3 to 43.4 °C (mean = 35.2 °C) for Joint Base Lewis-McCord, WA and Fort McCoy, WI, respectively (Appendix A). Historical precipitation ranged from 106.5-1469.6 mm (mean = 846.5 mm), for Yuma Proving Ground, AZ and Fort Polk, LA, respectively (Appendix A). The average annual precipitation and seasonal temperature variation calculated across installation assessment areas were then standardized using the score range procedure and then subtracted from one. This transformation caused installations in locations with low historical annual precipitation and temperature variation to have high subfactor scores. Resulting values ranged from zero to one, with higher values having greater relevance.

2.4 Adaptive capacity

Adaptive capacity is the ability of a species or system to adapt to changes in climate. Specific information about the evolutionary processes and mechanisms that influence different species' adaptive capacity (e.g., fecundity, mating system, spatial genetic diversity, phenotypic plasticity, etc.) is often not available and is likely hard to synthesize across species. Consequently, in this effort, two subfactors that are likely to affect the adaptive capacity of all species represented on installations were characterized: anthropogenic barriers to dispersal and availability of protected lands that can potentially act as secure stepping stones during range migration.

Anthropogenically altered landscapes (e.g., urban or agricultural areas) reduce and fragment available habitat, which leads to smaller and more isolated populations. They may also hinder the dispersal of species. Dispersal in the face of climate change is important in multiple ways. For example, if climate change causes spatial shifts in vegetation communities, listed and at-risk species that rely on these communities for habitat would need to track this spatial shift, otherwise available habitat would disappear and the species would likely be locally extirpated. Dispersal is also critical in maintaining metapopulation structure. Anthropogenic barriers (Table 1, Subfactor 3.1) to dispersal were characterized by calculating the percentage of

land within 50 km buffers of Federal lands represented by developed pasture/hay and cultivated crop cover types within the 2011 National Land Cover Dataset (Homer et al. 2015). The percent of installation buffers comprised of developed and other intensively used landcover types ranged from 0.2-54.2% (mean = 20.5%), for Dugway Proving Ground, UT and Camp Atterbury, IN, respectively (Appendix A). Values for this subfactor were standardized with a score range procedure. Resulting values ranged from zero to one, with higher values having greater relevance.

Protected lands (Table 1, Subfactor 3.2), unlike private lands, are unlikely to be anthropogenically altered in the future due to socio-economic drivers. Protected lands thus provide a coarse metric of the environmental stability of the landbase that species may be able to use for climate change adaptation (e.g., stepping stones of colonization during range shifts), even if their overlap with biodiversity priorities falls short (Jenkins et al. 2015), or they are externally impinged upon (Wilson et al. 2014, Martinuzzi et al. 2015).

Of protected lands, Federal landholdings offer listed and at-risk species the most comprehensive protections and also represent potential conservation partners for Army. In this effort, the influence of protected lands on adaptive capacity was estimated as the percentage of Federal lands within a 50 km buffer on installations. Spatial data for Federal lands were sourced from the USGS National Map. This dataset includes lands owned or administered by the Federal Government, including the Bureau of Land Management, Bureau of Reclamation, Forest Service, DoD, USFWS, National Park Service, Tennessee Valley Authority, and other agencies. The percent of installation buffers comprised of Federal lands ranged from 0.0-99.1% (mean = 19.1%). Several installations have no Federal lands within their buffers (e.g., Combat Training Center [CTC] Fort Custer Training Support [TS], MI; Fort Drum, NY; and Fort Knox, KY), while Camp Navajo, AZ is almost entirely surrounded by Federal lands (Appendix A). Before combining with the anthropogenic barrier subfactor (see Section 2.7), values for this subfactor were standardized with a score range procedure and then subtracted from one. This transformation caused installations with low percentages of Federal lands nearby to have high subfactor scores. Resulting values ranged from zero to one, with higher values having greater relevance.

2.5 Listed and at-risk species

The anticipated increase in restrictions on Army training land use due to climate change-related impacts on listed and at-risk species (Table 1, 4.1) will likely be a function of the number of species occurring on individual installations. A spreadsheet of listed and at-risk species for CONUS Army installations was compiled based on the most recent at-risk (NatureServe 2014) and listed (ACSIM 2010) species summaries (Appendix B). For species that are not Federally listed, or Candidates for listing, only Global Conservation Status Ranked species at risk of G1 (critically imperiled) and G2 (imperiled) were included. Currently listed species impose known impacts to installations, but at-risk species generally impose less of an impact due to fewer restrictions on training land use and greater flexibility of species management.

Despite installation efforts to proactively manage at-risk species, some are ultimately listed. Sperry et al. (2016) developed a logistic regression model of the probability that species proposed for listing under ESA will in fact be listed. Variables included in the model were the species taxonomic group and the percentage of the species' range lost to intensive human land use (e.g., urbanized or agricultural). This work used the former to weight at-risk species on installations by multiplying the number of species in each taxonomic group by the probabilities listed in Table 2. For at-risk species, an additional weighting, in which the value obtained in the former step was multiplied by 0.25, was applied. This was done to approximate the uncertainty in future listing of at-risk species that have not yet been petitioned for listing under the ESA. The derived numbers for each taxonomic group were then summed and added to the number of listed species on each installation. For example, Fort Hunter Liggett has six listed species and 21 at-risk plant species, which results in a value of 9.3 ($6 + [21 \times 0.63 \times 0.25]$). Resulting values ranged from 0.00 to 16.63 (mean = 3.92). Two installations (e.g., Camp Joseph T. Robinson, AK, and Fort William Henry Harrison, MT) had no listed or at-risk species, while Joint Base Lewis-McCord, WA had the most (Appendix A).

Because the magnitude of the impact that listed and at-risk species are likely to have on installations is also expected to vary as a function of available training area, these values were divided by installation area (km²) to generate a density-based index of listed and at-risk species for each installation

(Table 1, Subfactor 4.2). Resulting values ranged from 0.00 to 0.36 species/km² (mean = 0.25) (Appendix A). The values estimated for both subfactors were standardized with the score range procedure before being combined in the overall index of installation vulnerability (see Section 2.7).

Table 2. Probability of Federal listing by taxonomic group used to weight species at-risk. Numbers in parentheses represent number of species used to develop the model.

Taxonomic Group	Probability Listing
Arthropod (24)	0.29
Birds (18)	0.72
Fish (14)	0.64
Mammal (20)	0.65
Plants (35)	0.63
Reptile (10)	0.60
Source: Sperry et al. (2016).	

2.6 Installation ranks

This factor used the rankings of installations calculated by the Army ITAM Program in 2009. ITAM rankings were calculated based on a variety of factors including training throughput, installation acreage, and soil properties. Of the variables included in the rankings, throughput was more heavily weighted (twice that of other factors). Ranks vary from 1 to 6, with lower values indicating higher relevance for the Army's training and testing missions (Appendix A). The ITAM ranks were transformed using the score range procedure and then subtracted from one. Resulting values ranged from zero to one and had a positive relationship with importance.

2.7 Calculating climate change vulnerability scores for installations

To identify the relative risk that installations are potentially exposed to as a consequence of emerging climate change impacts, the standardized values of the two subfactors under each of the five factors were added (except installation rank, which does not have any subfactors), multiplied by a weight (Table 3) characterizing relative importance, and then summed for each installation (Kirkwood 1997). Note that equal weights (0.15), indicating equal importance, were assigned to all factors except the listed and at-risk species factor, which was assigned a larger weight (0.40) because of its specific emphasis within this vulnerability assessment.

The standardizing transformations applied to the various subfactors described in previous sections ensured that climate change vulnerability was not unduly influenced by the disparate values of the different subfactors. Installation climate change vulnerability scores generated by this process should be interpreted in a relative, rather than an absolute context. Climate change vulnerability was also estimated for installations without including the installation rank factor as it likely encompasses many variables considered by the BRAC process and this redundancy may not be desirable when this is the intended use. Appendix C presents these scores.

Table 3. Weights multiplied by the five factors to calculate climate change vulnerabilities of Army ITAM installations.

Factor	Weighting
Exposure	0.15
Sensitivity	0.15
Adaptive capacity	0.15
Listed and at-risk species	0.40
Installation rank	0.15

2.8 Examining differences in installation climate change vulnerability factors and scores across USFWS regions

Many listed and at-risk species can have surprisingly large geographic ranges (e.g., the Indiana bat), and consequently occur on multiple Army installations. In this case, there are potential benefits to developing and coordinating climate change management initiatives at a regional scale instead of on an installation-by-installation basis. For example, the magnitude of climate change exposure on one installation may cast doubt on the long-term viability of a local population of a given species, while exposure might be minimal on other installations within its range. Given this scenario, it is possible that cross-installation, conservation banking strategies would be welcomed by the USFWS. This type of strategy would likely require significant coordination with multiple USFWS Ecological Services Field Offices and their regional office(s).

Regional differences in the factors that affect climate change vulnerability are also anticipated due to spatial variation in biogeography, elevation, land-use patterns, etc. Thus across regions, installations and their associated listed and at-risk species will be subject to varying climate change vul-

nerabilities. To identify emerging regional challenges and potential opportunities, mean scores for each of the five factors and overall climate change vulnerability were compared for installations within the eight USFWS regions (Table 4).

Table 4. USFWS regions and associated states.

Region	States
1 Pacific	Hawaii, Idaho, Oregon, Washington, American Samoa, Commonwealth of the Northern Mariana Islands, Guam and the Pacific Trust Territories
2 Southwest	Arizona, New Mexico, Oklahoma, and Texas
3 Midwest	Illinois, Indiana, Iowa, Ohio, Michigan, Minnesota, Missouri, and Wisconsin
4 Southeast	Alabama, Arkansas, Florida, Georgia, Kentucky, Louisiana, Mississippi, North Carolina, South Carolina, Tennessee, Puerto Rico, and the U.S. Virgin Islands
5 Northeast	Connecticut, Delaware, District of Columbia, Maine, Maryland, Massachusetts, New Hampshire, New Jersey, New York, Pennsylvania, Rhode Island, Vermont, Virginia, and West Virginia
6 Mountain-Prairie	Colorado, Kansas, Montana, Nebraska, North Dakota, South Dakota, Utah, and Wyoming
8 Pacific Southwest	California, Nevada

3 Results and Discussion

3.1 Direct and indirect exposure

The five installations with the highest values for the factor characterizing direct and indirect climate change exposure, ordered high-to-low were Fort Sill, OK, Fort Eustis, VA, Fort Carson, CO, Camp Williams, UT, and Camp Dodge, IA (Table 5). Military Training Center-Heavy (MTC-H) Camp Roberts, CA and Fort Hunter Liggett, CA had the lowest estimated scores for this factor. Many coastal portions of the country are projected to experience smaller changes in temperatures and precipitation than the interior United States, due to buffering effects of the oceans. Land tends to cool and heat more rapidly than water, such that areas away from large bodies of water experience greater seasonal extremes of temperature than do coastal communities. Proximity to large bodies of water also tends to positively influence precipitation levels; coastal locations receive generally higher amounts than interior areas primarily due to higher levels of evaporation. Also, the indirect impacts of sea level rise are projected to be greater along the Atlantic and Gulf coasts, than along the Pacific coast (Krasting et al. 2016).

Table 5. Weighted scores for five factors used to estimate climate change vulnerability of 43 CONUS Army ITAM installations. Installations are ordered high-to-low based on vulnerability.

Installation	State	Exposure*	Sensitivity	Adaptive Capacity	Species	Installation Rank	Vulnerability
Parks Reserve Forces Training Area	CA	0.106	0.235	0.229	0.476	0.090	1.137
Joint Base Lewis-McCord	WA	0.108	0.181	0.211	0.454	0.150	1.103
Fort Huachuca	AZ	0.170	0.198	0.094	0.473	0.030	0.965
Fort Bliss	TX	0.162	0.187	0.091	0.311	0.150	0.901
Fort Carson	CO	0.264	0.144	0.137	0.142	0.150	0.838
Camp Atterbury	IN	0.229	0.096	0.293	0.113	0.090	0.820
Yakima Training Center	WA	0.090	0.203	0.219	0.126	0.150	0.788
Fort Hood	TX	0.211	0.147	0.190	0.086	0.150	0.784
Fort Bragg	NC	0.109	0.113	0.232	0.209	0.120	0.783
MTC-H Camp Roberts	CA	0.063	0.202	0.194	0.260	0.060	0.780
Fort Riley	KS	0.222	0.087	0.232	0.115	0.120	0.776
Fort Campbell	KY	0.214	0.085	0.267	0.072	0.120	0.757
Fort Sill	OK	0.272	0.116	0.240	0.033	0.090	0.750
Fort Benning	GA	0.124	0.117	0.188	0.197	0.120	0.746
Fort Stewart	GA	0.104	0.134	0.195	0.186	0.120	0.739

* Blue = minimum, white = median (50th percentile), red = maximum, and transitional colors represent intermediate percentiles.

Installation	State	Exposure*	Sensitivity	Adaptive Capacity	Species	Installation Rank	Vulnerability
Camp Bullis	TX	0.176	0.163	0.267	0.036	0.090	0.733
Fort Eustis	VA	0.265	0.126	0.211	0.057	0.030	0.689
Fort Hunter Liggett	CA	0.053	0.174	0.111	0.255	0.090	0.684
Fort Gordon	GA	0.108	0.120	0.199	0.194	0.060	0.680
Fort Knox	KY	0.217	0.093	0.215	0.063	0.090	0.678
Fort Rucker	AL	0.134	0.117	0.213	0.109	0.090	0.663
Fort McCoy	WI	0.197	0.070	0.241	0.060	0.090	0.657
White Sands Military Range	NM	0.171	0.173	0.051	0.257	0.000	0.653
CTC Fort Custer TS	MI	0.207	0.108	0.280	0.050	0.000	0.646
Aberdeen Proving Ground	MD	0.204	0.106	0.237	0.090	0.000	0.637
Camp Dodge Joint Maneuver Training Center	IA	0.237	0.087	0.269	0.010	0.030	0.633
Military Training Area-Light (MTA-L) Camp Williams	UT	0.254	0.162	0.119	0.015	0.060	0.610
Fort Drum	NY	0.166	0.076	0.175	0.056	0.120	0.592
Fort Leonard Wood	MO	0.227	0.085	0.088	0.102	0.090	0.592
Camp Joseph T Robinson	AR	0.195	0.094	0.239	0.000	0.060	0.588
Fort Polk	LA	0.185	0.101	0.141	0.038	0.120	0.585
Fort Chaffee Military Training Center (MTC)	AR	0.212	0.087	0.152	0.035	0.090	0.576
Fort Dix	NJ	0.129	0.093	0.252	0.009	0.090	0.572
MTC-H Camp Grayling	MI	0.184	0.101	0.153	0.072	0.060	0.571
Fort Jackson	SC	0.117	0.122	0.220	0.050	0.060	0.570
Fort Pickett, ARNG MTC	VA	0.131	0.102	0.165	0.079	0.090	0.567
Fort A P Hill	VA	0.138	0.108	0.193	0.034	0.090	0.563
Military Training Area (MTA) Fort Wm Henry Harrison	MT	0.216	0.164	0.122	0.000	0.060	0.561
Fort Irwin	CA	0.102	0.195	0.015	0.065	0.120	0.498
Fort Lee	VA	0.120	0.108	0.189	0.079	0.000	0.496
Yuma Proving Ground	AZ	0.089	0.207	0.089	0.077	0.000	0.463
Camp Navajo	AZ	0.228	0.139	0.001	0.043	0.030	0.440
Dugway Proving Ground	UT	0.199	0.143	0.016	0.077	0.000	0.435

* Blue = minimum, white = median (50th percentile), red = maximum, and transitional colors represent intermediate percentiles.


Differences in direct and indirect climate change exposure were identified across USFWS regions. Region 6 (Mountain-Prairie) had the largest mean exposure score, which was significantly higher than mean scores in Regions 1 (Pacific), 8 (Pacific Southwest) and 4 (Southeast). Mean exposure scores for Regions 2 (Southwest) and 3 (Midwest) were also notably high, being significantly higher than one, or more of the other seven regions (Table 6).

Table 6. Mean scores and ranges (in parentheses) for five climate change factors and the overall vulnerability of 43 CONUS Army ITAM installations within seven USFWS regions. Means designated by the same letter within columns do not differ at $P = 0.05$ by Tukey's Honestly Significant Difference in an across-region comparison.

USFWS Region	Count	Exposure	Sensitivity	Adaptive Capacity	Species	Installation	Vulnerability
1	2	0.099 (0.090 - 0.108) ab	0.192 (0.181 - 0.203) a	0.215 (0.211 - 0.219) a	0.289 (0.125 - 0.454) a	0.150 (0.150 - 0.150) a	0.946 (0.788 - 1.103) a
2	8	0.185 (0.089 - 0.272) c	0.166 (0.116 - 0.207) b	0.128 (0.001 - 0.267) a	0.164 (0.036 - 0.473) a	0.068 (0.000 - 0.150) a	0.711 (0.440 - 0.965) a
3	6	0.214 (0.184 - 0.237) ad	0.091 (0.069 - 0.108) abc	0.221 (0.088 - 0.293) a	0.068 (0.010 - 0.113) a	0.060 (0.000 - 0.090) a	0.653 (0.571 - 0.820) a
4	11	0.156 (0.104 - 0.217) e	0.107 (0.085 - 0.122) abd	0.205 (0.141 - 0.267) a	0.105 (0.000 - 0.201) a	0.095 (0.060 - 0.120) a	0.669 (0.569 - 0.783) a
5	7	0.165 (0.119 - 0.265) f	0.103 (0.076 - 0.126) Abe	0.203 (0.165 - 0.237) a	0.056 (0.009 - 0.090) a	0.060 (0.000 - 0.120) a	0.588 (0.496 - 0.689) a
6	5	0.231 (0.198 - 0.264) beg	0.140 (0.087 - 0.162) cf	0.125 (0.016 - 0.231) a	0.069 (0.000 - 0.142) a	0.078 (0.000 - 0.150) a	0.644 (0.435 - 0.838) a
8	4	0.081 (0.053 - 0.106) cdg	0.202 (0.174 - 0.235) cdef	0.137 (0.015 - 0.229) a	0.264 (0.065 - 0.476) a	0.090 (0.060 - 0.090) a	0.775 (0.498 - 1.137) a

The variability in climate change exposure identified across installations and regions suggests different mitigation strategies will likely be needed. First, installations with relatively high exposure but lower scores for sensitivity and adaptive capacity (e.g., Fort Leonard Wood, MO and Fort Chaffee MTC, AR) can attempt to reduce the local consequences of climate change on the specific habitat needs of affected species when feasible (Table 5, Figure 1, Class 1). For example, water could be added to breeding ponds of at-risk amphibian species to extend the hydroperiod, facilitating successful metamorphosis and juvenile dispersal. Installations and species that are likely to be exposed to climate change, but that do not suffer from high sensitivity or limited adaptive capacity, have somewhat optimistic prospects for near-term local population persistence and/or climate change adaptation. If feasible, efforts should focus on reducing climate change exposure and maintaining low sensitivity and adaptive capacity scores.

Figure 1. Conceptual diagram of combinations of climate change exposure, sensitivity, and adaptive capacity that identify seven classes of vulnerability having different implications for prioritization and strategic planning.



Second, installations and species subjected to relatively high climate change exposure and sensitivity, but low adaptive capacity scores, can be classified as potential adapters (see Foden et al. 2013, Figure 1, Class 2). These results suggest that Fort Carson, CO, MTA-L Camp Williams, MTA, UT, Fort William Henry Harrison, MT, and Camp Navajo, AZ, and possibly Dugway Proving Ground, UT, will likely face this combination of factors (Table 5). In this situation, efforts should focus on reducing exposure and sensitivity, while maintaining adaptive capacity. Although this approach used historical climate variables as a means of characterizing sensitivity for multiple species on installations, the mitigation efforts used by installations to either reduce or maintain this factor will need to target the sensitivity of specific species. For example, climate change-related increases in species' sensitivity to the impacts of certain invasive species could be mitigated by implementing control efforts that specifically target this source of sensitivity (Abatzoglou and Kolden 2011).

Third, where exposure and adaptive capacity scores are high, but sensitivity is low, one would anticipate some potential for species persistence (Figure 1, Class 3; Foden et al. 2013). These results suggest approximately eight of the study installations will likely face this combination of factors, including: Camp Atterbury, IN; Fort Riley, KS; Fort Campbell, KY; Fort Knox, KY; Fort McCoy, WI; CTC Fort Custer, MI; Aberdeen Proving Ground, MD; and Camp Dodge Joint Maneuver Training Center, IA (Table 4). In this situation, efforts should focus on reducing the impacts of climate change exposure and constraints on adaptive capacity, while maintaining low sensitivity.

Fourth, installations and species subjected to relatively high climate change exposure, sensitivity, and adaptive capacity scores will likely be the most vulnerable. Only Fort Eustis, VA was found to have relatively high scores for these three factors (Table 5; Figure 1, Class 7). To minimize the loss of unrestricted access to training land use on installations due to the additive effects of climate change and listed and at-risk species, efforts should target: (1) the conservation of any existing at-risk species to help eliminate the need for listing, and (2) negotiations with the USFWS to evaluate the merits and feasibility of conserving populations of listed species on the installation.

Fifth, installations and species subjected to lower climate change exposure, sensitivity, and adaptive capacity scores are less likely to be vulnerable in contrast with other installations. Fort Pickett, VA, and to a lesser degree Fort A.P. Hill, VA, were found to have lower scores for all three factors (Table 5). One might anticipate some potential for species persistence under this scenario.

It is worth noting that this work used climate projections made available by the ClimateWizard team and NatureServe, however the approach can readily use other sources of climate change data as they may become available. It is also important to remember that the adopted approach characterized climate change as relative change, but no explicit interpretation of the magnitude of change was made. Consequently, it is possible that high relative exposure may in fact represent a projected change in climate that does not warrant great concern. That being said, Army installations from across the Continental United States were included to ensure that some of the greatest and smallest changes in climate anticipated for the country would be represented in the analysis, and would therefore be likely to reflect actionable val-

ues. Still, there is also potentially great benefit in conducting species-specific climate change vulnerability analyses at multiple scales before finalizing and targeting mitigation actions (e.g., Hohmann and Wall 2016).

3.2 Sensitivity

The sensitivity factor ranged from 0.070 to 0.235 for Fort McCoy, WI and Parks Reserve Forces Training Area, CA, respectively (Table 5). Regions that receive higher amounts of precipitation are anticipated to be less affected by projected changes in precipitation than are regions that receive little precipitation. Likewise, regions with high seasonal temperature variation are expected to be less impacted by temperature changes than regions with less seasonal temperature variation. Consequently, many of the installations with the highest sensitivity scores are from arid regions, such as the Southwest, while the lowest sensitivity scores are from mesic, continental interior regions.

Differences in mean sensitivity scores were identified across USFWS regions, with Army installations in Regions 1 (Pacific), 2 (Southwest), and 8 (Pacific Southwest) having higher sensitivity scores than installations in Regions 3 (Great Lakes-Big Rivers), 4 (Southeast), and 5 (Northeast). Additionally, sensitivity scores were higher in Region 8 than Region 6 (Mountain-Prairie) and higher in Region 6 than Region 3 (Table 6). The variability in climate change sensitivity identified across installations and regions suggests different mitigation strategies will likely be needed.

First, installations and species that are likely to be sensitive to climate change, but that are not likely to suffer from high exposure or limited adaptive capacity have optimistic prospects for near-term local population persistence and/or climate change adaptation. Installations fitting this classification, include Fort Huachuca, CA, Fort Bliss, TX, Fort Hunter Liggett, CA, Fort Irwin, CA, and Yuma Proving Ground, AZ. These installations would benefit from efforts to maintain lower levels of exposure and lessen constraints on adaptive capacity (Table 5, Figure 1, Class 4). They can also attempt to reduce the local consequences of high sensitivity for affected species when feasible. As suggested above, efforts to reduce sensitivity should focus on actionable reduction of climate change-related sensitivity such as invasive species (Hellman et al. 2008), predation risk (e.g., Cox et al. 2013), or temperature dependent sex determination (Janzen

1994). Efforts could also focus on maintaining low climate change exposure and adaptive capacity scores.

Second, installations and species having relatively high sensitivity and adaptive capacity scores, but low climate change exposure can be classified as having high latent risk (Foden et al. 2013; Figure 1, Class 5). These installations pose a potential future risk should climate change projections be underestimated, or for time periods beyond 2050. These results suggest that Parks Reserve Forces Training Area, CA; Joint Base Lewis-McCord, WA; and Yakima Training Center, WA have high latent risk (Table 5). In this situation, efforts should focus on reducing exposure and sensitivity, while maintaining adaptive capacity.

The consequences of sensitivity are undoubtedly tied to the magnitude and timing of climate change exposure and will vary among species. Species-centric climate change vulnerability assessments typically include many additional considerations that focus on habitat microsite, physiology, diet, inter-specific interactions, phenology, etc. (e.g., Young et al. 2015), as well as other measures of exposure, such as habitat change and climate velocity (Loarie et al. 2009, Dichinson et al. 2014). This effort only examined historical precipitation and seasonal temperature variation. Consequently, there is potentially great benefit in conducting species-specific climate change vulnerability analyses at multiple scales before finalizing and targeting mitigation actions (e.g., Hohmann and Wall 2016).

3.3 Adaptive capacity

The adaptive capacity factor ranged from 0.001 to 0.293 for Camp Navajo, AZ and Camp Atterbury, IN, respectively (Table 5). Camp Navajo, AZ, like many other installations with low adaptive capacity scores, is located in the western United States, where there are large tracts of Federal lands and much of the landscape has not been transformed by development. In contrast, Camp Atterbury, IN, which had a higher score, is located in the Midwest, where much of the landscape is under intensive agricultural use and comparatively little area is under Federal stewardship.

No differences in adaptive capacity scores were identified across USFWS regions (Table 6). This was somewhat surprising given that the percent of developed landcover was expected to be higher and the percent of Federal

lands was expected to be lower in the eastern United States compared to the western United States. It could be that local encroachment on installations in the western United States has increased to a greater degree than on eastern installations, and that this has offset any potential regional differences in overall adaptive capacity.

The variability in adaptive capacity identified across installations suggests that different mitigation strategies will likely be needed. Installations and species that are likely to experience constraints on adaptive capacity (Figure 1, Class 6), but that will not suffer from high exposure or sensitivity, have multiple options for mitigating climate change impacts. High adaptive capacity scores suggest constraints on species dispersal and conservation partnering opportunities with other Federal landowners. Although other Federal land managers are preferred partners, the conservation achievements that can be accomplished by partnering with state and private landowners should not be discounted (e.g., Candidate Conservation Agreement, Safe Harbor Agreement). Installations having high adaptive capacity scores, can promote regional conservation with private partners via the DoD Readiness and Environmental Protection Integration (REPI) and Army Compatible Use Buffer (ACUB) programs, which have contributed greatly to military readiness and the environmental protection (Messer et al. 2016). Of the targeted installations, only Fort Dix, NJ fit this classification (Table 5).

This approach to estimating constraints on species and installation adaptive capacity used the percentage of land managed by Federal agencies and the percentage eliminated as potential habitat by urban development and other intensive human land uses within a 50 km buffer of installation boundaries. Although this approach is appropriate for screening a large number of species on multiple installations, its suitability for evaluating the adaptive capacity of any individual species is uncertain. For example, species endemic to a small geographic range, dependent on highly localized and rare edaphic conditions, and having limited dispersal ability (e.g., many cave and plant species) would not be expected to shift their range under climate change even if much of the landscape was undeveloped. Species have unique habitat needs and dispersal abilities; consequently, strategies used by installations to improve adaptive capacity should strive to meet the specific needs of target species. An example of a more species-centric approach would be to specifically evaluate the availability and connectedness of habitats used by focal species (e.g., McRae and Shah 2009).

Additionally, there are likely benefits to adopting a dynamic as opposed to a static assessment of regional land use. It is anticipated that the land use within the landscapes surrounding installations will change over time due to urbanization as well as climate (Ordonez et al. 2014). The location and magnitude of these changes will vary based on shifts in the population, zoning regulations, resource availability, land form constraints, etc. Detailed installation-specific assessments will likely want to incorporate projections of future population densities and land use change. Several methods that are available include the ERDC-CERL-maintained Regional Urban Growth model (RUG) (Westervelt et al. 2011), the Land-use Evolution and Impact Assessment Model (LEAM) (Deal and Pallathucheril 2009), and the U.S. Environmental Protection Agency's (USEPA's) Integrated Climate and Land Use Scenarios (ICLUS) model (Bierwagen and Morefield 2014).

3.4 Listed and at-risk species

The scores for the listed and at-risk species factor ranged from 0.000 to 0.476 (Table 5). Neither MTA Fort William Henry Harrison, MT, nor Camp Joseph T Robinson, AR, are known to have any listed or at-risk species and consequently had a score of zero. The installation having the highest score for this factor was Parks Reserve Forces Training Area, CA, which has a high density of listed and at-risk species. Other installations with notably high scores for this factor included Fort Huachuca, CA, and Joint Base Lewis-McCord, WA.

No differences in mean listed and at-risk species scores were identified across USFWS regions (Table 6). Region 5 (Northeast) had the lowest (0.058) and Region 1 (Pacific) had the highest (0.289) mean scores for the listed and at-risk species factor. Fortuitously, the latter region also had one of the lowest climate change exposure scores (Table 6).

This assessment used information about the number and density of listed and at-risk species found on installations based on a 2010 summary of listed species and a more recently completed summary of at-risk species. However this approach is potentially limiting in a number of ways.

First, the listed and at-risk species represented on installations is ever changing due to listing decisions, down listing, species surveys, and changes in the conservation status of species. This is particularly true at

the present time due to the large number of listing petitions that the USFWS is actively reviewing. Sperry et al. (2016) estimated that as many as 230 species under active review potentially occur on Army installations.

Second, the number and density of species may or may not directly translate into training restrictions, which are instead influenced by the number and size of populations, as well as their overlap and compatibility with training land use. One could explicitly evaluate this overlap where data are available, or one could employ species distribution modeling to better characterize potential overlap. Consequently, it would be appropriate to consider the scores presented for this factor as dynamic estimates. Also, local scale assessments are likely necessary to inform installation management decisions.

3.5 Installation ranks

Scores for the installation rank factor represent a simple transformation of ITAM rankings and thus provide no novel information when examined separately. Since mean installation rank scores did not differ among USFWS regions (Table 6), no additional discussion of this factor is presented here. However, it is worth mentioning that, from a BRAC perspective, a comparison of the vulnerability score to the ITAM rankings could provide a way to define high value installations that are low risk versus high/high, low/low, or low/high (Value/ITAM). In BRAC, the low vulnerability/high ITAM installation would be considered of greatest value. Appendix C provides vulnerability scores calculated without installation ranks.

3.6 Threatened and endangered species related climate change vulnerabilities of installations

Previous sections summarized the scores of each of the five factors used to estimate installation climate change vulnerabilities. This section summarizes results for the aggregate climate change vulnerability scores estimated after applying weights to the factors and summing for each installation (see Section 2.7). The ranking of installations and their relative values are potentially useful for: (1) identifying the most vulnerable installations, (2) planning additional funding needs beyond those that might already be identified for installations based on currently listed and at-risk species, (3) prioritizing limited funds available for abatement, or (4) informing the BRAC process.

Installation climate change vulnerability scores ranged from 0.435 to 1.137 for Dugway Proving Ground, UT and Parks Reserve Forces Training Area, CA, respectively (Table 5). The latter installation had a relatively low climate change exposure score, but intermediate to high scores for the remaining four factors. In contrast, Dugway Proving Ground, UT had low scores for adaptive capacity and installation rank factors, but intermediate exposure and sensitivity scores. Fortuitously, nearly all installations with high vulnerability scores (i.e., >0.765, upper quartile) have relatively low scores for one or more of the five component factors, which suggests that mitigation options are likely available. Fort Hood, TX may present the greatest management challenge in the face of climate change, given that the listed species factor and all the explicitly climate change-related factors had intermediate to high scores (Table 5).

Comparisons across USFWS regions did not identify any differences in mean installation climate change vulnerability scores. Still, the differences identified for individual factors (i.e., exposure and sensitivity) may provide insight about the potential need for and advantages of using regional mitigation approaches (see Sections 3.1-3.4). Wilhoit et al. (2016) evaluated seven regionally representative Army installations for vulnerability to climate change based on a simple approach that was comparable to the listed and at-risk species factor calculated in this effort. However, their approach did not include any specific information about climate change vulnerability. Interestingly, the five CONUS installations evaluated by both efforts were ranked similarly, except for Fort Bliss, TX, which was given a much higher ranking when additional climate change variables were considered.

Sperry et al. (2016) also conducted an Army-wide assessment of the potential emerging risk to installations associated with listed and at-risk species. Since their approach did not consider the consequences of climate change on potential future impacts, it is interesting to examine the potential additional effect of climate change vulnerability identified by this approach. Five of the installations included in this effort were among the top 20 installations they determined most likely to be impacted. Considering their ranking of installations relative to that shown in Table 5, it is apparent that climate change vulnerability has the potential to increase impacts on Army training beyond that suggested by Sperry et al. (2016). It may also offer some reassurance to Army managers to know that climate change vulnerability is not likely to exacerbate impacts to all installations.

4 Conclusions and Recommendations

4.1 Conclusions

Mandates for DoD and the Army to address climate change are in place, but specific approaches that might be adopted to address assessment, monitoring, and adaptation strategies are still evolving. Although vulnerability assessments are an important first step in ensuring future conservation successes for listed and at-risk species, this information is not widely available to regional or Headquarters decision makers. This effort developed and demonstrated a relatively simple method for generating climate change vulnerability assessments for installations using factors that characterized the number of listed and at-risk species, climate change exposure, sensitivity, adaptive capacity and installation importance. The approach is appealing in that it: (1) transparently retains the information about the five factors used in ranking, (2) satisfies the need for consistent assessment methods that can be conducted with broadly available data and that allow meaningful comparisons, and (3) complements recent Army-wide, installation-focused evaluations of proposed species' potential impacts on training (Sperry et al. 2016).

4.2 Recommendations

The approach to vulnerability assessment demonstrated here is suitable as a first pass assessment for evaluating whether climate changed driven impacts to listed and at-risk species will likely affect installation resilience. Army decision makers should consider using the results to assess the relative vulnerability of installations, identify where additional more detailed vulnerability assessments might need to be conducted, and integrate climate change considerations into the BRAC process.

Appendix A: Installation Subfactor Values

Table A-1. Values of subfactors used to inform factors in the climate change vulnerability assessments.

State	Installation	Exposure			Sensitivity		Adaptive Capacity		Threatened and Endangered Species		Installation Rank
		aetpet	Temperature Change (°C)	Coastal Vulnerability Index	Mean Annual Precipitation (mm)	Mean Temperature Variation (°C)	% Federal Land in Buffer	% Developed Land in Buffer	# of Species	Species Density	
WA	Yakima Training Center	-0.04632	2.3	0	250	34.61	14.2	32.9	4.95	0.006	1
WA	Joint base Lewis-McCord	-0.08903	2.0	0	1192	24.21	8.3	26.6	16.63	0.049	1
WA	Fort Lewis	-0.08903	2.0	0	1192	24.21	6.0	14.0	#N/A	#N/A	1
TX	Fort Hood	-0.09835	2.7	0	819	33.80	1.6	15.4	3.36	0.005	1
CO	Fort Carson	-0.10701	3.1	0	346	40.84	16.9	4.7	5.21	0.016	1
TX	Fort Bliss	-0.05841	2.8	0	267	36.44	53.2	8.1	12.7	0.005	1
GA	Fort Stewart	-0.06257	2.2	0	1206	29.96	0.8	16.9	7.29	0.010	2
KS	Fort Riley	-0.07525	3.1	0	810	41.56	3.7	31.6	4.23	0.012	2
LA	Fort Polk	-0.09903	2.5	0	1470	30.53	22.6	9.4	1.43	0.003	2
CA	Fort Irwin	-0.01682	2.7	0	125	37.37	90.2	0.7	2.64	0.002	2
NY	Fort Drum	-0.04511	2.9	0	1009	40.24	0.0	9.3	2.07	0.005	2
KY	Fort Campbell	-0.09433	2.8	0	1259	35.50	15.9	50.8	2.23	0.017	2
NC	Fort Bragg	-0.04838	2.4	0	1204	32.74	0.8	30.3	7.72	0.021	2
GA	Fort Benning	-0.06713	2.3	0	1247	31.55	0.2	13.8	7.46	0.016	2
CA	Parks Reserve Forces Training Area	-0.055	2.3	0	529	26.60	2.4	30.1	3.16	0.363	3
OK	Fort Sill	-0.11987	3.0	0	749	38.72	5.4	35.6	1.16	0.004	3
AL	Fort Rucker	-0.07843	2.3	0	1387	29.66	0.2	23.1	2.77	0.039	3
VA	Fort Pickett, ARNG MTC	-0.05333	2.6	0	1132	35.11	0.2	5.9	2.36	0.020	3
WI	Fort McCoy	-0.05793	3.1	0	834	43.45	0.0	33.0	2.22	0.006	3
MO	Fort Leonard Wood	-0.08496	3.0	0	1062	38.29	48.5	4.4	3.4	0.018	3
KY	Fort Knox	-0.09069	2.9	0	1169	35.82	0.0	23.4	2.3	0.007	3
CA	Fort Hunter Liggett	-0.03561	2.1	0	531	34.33	41.1	8.5	9.3	0.029	3
NJ	Fort Dix	-0.045	2.6	0	1172	35.77	1.6	37.9	0.15	0.005	3
AR	Fort Chaffee MTC	-0.08038	2.9	0	1140	36.90	7.2	4.8	1.16	0.006	3

State	Installation	Exposure			Sensitivity			Adaptive Capacity		Threatened and Endangered Species		
		aetpet	Temperature Change (°C)	Coastal Vulnerability Index	Mean Annual Precipitation (mm)	Mean Temperature Variation (°C)	% Federal Land In Buffer	% Developed Land In Buffer	# of Species	Species Density	Installation Rank	
VA	Fort A P Hill	-0.04985	2.7	0	1081	35.12	0.9	16.3	1.07	0.007	3	3
TX	Camp Bullis	-0.08788	2.5	0	830	31.54	1.3	43.1	0.57	0.020	3	3
IN	Camp Atterbury	-0.08454	3.0	0	1062	36.91	4.8	54.2	3.31	0.031	3	3
CA	MTC-H Camp Roberts	-0.037	2.2	0	376	33.00	6.4	19.7	7.58	0.071	4	4
MI	MTC-H Camp Grayling	-0.06117	2.9	0	812	39.77	19.0	11.7	2.76	0.005	4	4
UT	MTA-L Camp Williams	-0.089	3.2	0	369	38.21	29.0	4.7	0.2	0.009	4	4
MT	MTA Fort Wm Henry Harrison	-0.1	2.7	0	314	38.74	36.7	10.0	0	0.000	4	4
SC	Fort Jackson	-0.06102	2.4	0	1145	32.33	2.0	26.4	1.47	0.014	4	4
GA	Fort Gordon	-0.055	2.3	0	1148	32.59	2.8	19.2	6.47	0.035	4	4
AR	Camp Joseph T Robinson	-0.076	2.8	0	1241	34.63	0.9	32.7	0	0.000	4	4
AZ	Fort Huachuca	-0.06578	2.7	0	461	32.34	36.8	0.3	15.77	0.085	5	5
VA	Fort Eustis	-0.048	2.5	14.14	1150	31.81	5.7	25.4	1	0.030	5	5
AZ	Camp Navajo	-0.086	3.0	0	609	37.83	99.1	0.5	1.16	0.014	5	5
IA	Camp Dodge Joint Maneuver Training Center	-0.074	3.2	0	813	41.55	1.0	43.4	0.16	0.005	5	5
AZ	Yuma Proving Ground	-0.01	2.7	0	107	36.09	74.6	19.0	3.16	0.001	6	6
NM	White Sands Military Range	-0.06	2.8	0	259	38.28	76.2	6.2	10.62	0.002	6	6
VA	Fort Lee	-0.05	2.5	0	1145	34.22	1.6	15.2	1	0.050	6	6
UT	Dugway Proving Ground	-0.05	3.2	0	202	43.00	88.6	0.2	3.16	0.001	6	6
MI	CTC Fort Custer TS	-0.0777	2.9	0	896	37.64	0.0	47.1	0.3	0.039	6	6
MD	Aberdeen Proving Ground	-0.054	2.7	5.5	1124	34.71	10.9	37.4	3.16	0.012	6	6

THIS PAGE INTENTIONALLY LEFT BLANK

Appendix B: Lists of Threatened, Endangered, and At-Risk Species by Installation

Table B-1. List of threatened, endangered, and at-risk species found on or near 43 CONUS Army ITAM installations.

Installation	State	Common Name	Scientific Name	Taxonomic Group	Location	Status*
Fort Rucker	AL	American alligator	Alligator mississippiensis	Reptile	On-site	T(SA)
Fort Rucker	AL	Fly's Brickell-bush	Brickellia cordifolia	Plant	On-site	SAR
Fort Rucker	AL	Tapered pigtoe	Fusconaia burkei	Mussel	On-site	C
Fort Rucker	AL	Gopher tortoise	Gopherus polyphemus	Reptile	On-site	C
Fort Rucker	AL	Southern sandshell	Hamia australis	Mussel	On-site	C
Fort Rucker	AL	Choctaw bean	Villosa choctawensis	Mussel	On-site	C
Camp Joseph T. Robinson	AR	NA	NA	NA	NA	NA
Fort Chaffee MTC	AR	American burying beetle	Nicrophorus americanus	Insect	On-site	E
Fort Chaffee MTC	AR	Nuttall's Cornsalad	Valerianella nuttallii	Plant	On-site	SAR
Camp Navajo	AZ	Creeping Milk-Vetch	Astragalus troglodytus	Plant	On-site	SAR
Camp Navajo	AZ	Mexican spotted owl	Strix occidentalis lucida	Bird	On-site	T
Fort Huachuca	AZ	Huachuca Giant-Skipper	Agathymus evansi	Invertebrate	On-site	SAR
Fort Huachuca	AZ	Sonora tiger salamander	Ambystoma tigrinum stebbinsi	Amphibian	On-site	E
Fort Huachuca	AZ	Huachuca Milkvech	Astragalus hypoxylus	Plant	On-site	SAR
Fort Huachuca	AZ	Western Yellow-billed cuckoo	Coccyzus americanus occidentalis	Bird	On-site	C
Fort Huachuca	AZ		Dryopteris knoblochii	Plant	On-site	SAR
Fort Huachuca	AZ		Dryopteris rossii	Plant	On-site	SAR
Fort Huachuca	AZ	Southwestern willow flycatcher	Empidonax traillii eximius	Bird	Contiguous	E
Fort Huachuca	AZ	Arid Throne Fleabane	Erigeron arisoli	Plant	On-site	SAR
Fort Huachuca	AZ	Lemmon's Fleabane	Erigeron lemmonii	Plant	On-site	SAR
Fort Huachuca	AZ	Pringle's Fleabane	Erigeron pringlei	Plant	On-site	SAR
Fort Huachuca	AZ	Eared Quetzal	Euphonia neoxenus	Plant	On-site	SAR
Fort Huachuca	AZ	Santa Catalina Burstwort	Hermannia pauciflora	Plant	On-site	SAR
Fort Huachuca	AZ	Rutter's Golden-Aster	Heterotheca rutteri	Plant	On-site	SAR
Fort Huachuca	AZ	Purple-spiked Coralroot	Hexalectris warnockii	Plant	On-site	SAR
Fort Huachuca	AZ	Pringle's Hawkweed	Hieracium pringlei	Plant	On-site	SAR
Fort Huachuca	AZ	Arizona treefrog	Hyla wrightorum pop. 2	Amphibian	On-site	C

* Species are designated by either E = endangered, T = threatened, UR = under review, C = candidate species, SA = similarity of appearance receives the same protection as listed, or SAR = species at risk. Species are considered to be located on site if they reside within a 2 km distance of the installation (ACSIM 2010, NatureServe 2014).

Installation	State	Common Name	Scientific Name	Taxonomic Group	Location	Status*
Fort Huachuca	AZ	Lesser long-nosed bat	<i>Leptonycteris curasoae yerbabuenae</i>	Mammal	On-site	E
Fort Huachuca	AZ	Huachuca water-umbel	<i>Lilaeopsis schaffneriana ssp. recurva</i>	Plant	On-site	E
Fort Huachuca	AZ	Huachuca Mountain Lupine	<i>Lupinus huachucae</i>	Plant	On-site	SAR
Fort Huachuca	AZ	Spikedace	<i>Meda fulgida</i>	Fish	Contiguous	T
Fort Huachuca	AZ	Box Canyon Muhly	<i>Muhlenbergia dubioides</i>	Plant	On-site	SAR
Fort Huachuca	AZ	Palmer's Muhly	<i>Muhlenbergia palmeri</i>	Plant	On-site	SAR
Fort Huachuca	AZ	Huachuca springsnail	<i>Pygulopsis thompsoni</i>	Snail	On-site	C
Fort Huachuca	AZ	Chiricahua leopard frog	<i>Rana chiricahuensis</i>	Amphibian	On-site	T
Fort Huachuca	AZ	Ramsey Canyon Leopard Frog	<i>Rana subaquavocalis</i>	Amphibian	On-site	SAR
Fort Huachuca	AZ	Chiricahua Mountain Brookweed	<i>Samolus vagans</i>	Plant	On-site	SAR
Fort Huachuca	AZ	Huachuca Groundsel	<i>Senecio multidentatus</i> var. <i>huachucae</i>	Plant	On-site	SAR
Fort Huachuca	AZ	Garden Canyon Talussnail	<i>Sonorella dalli</i>	Mollusk	On-site	SAR
Fort Huachuca	AZ	Mexican spotted owl	<i>Strix occidentalis lucida</i>	Bird	On-site	T
Fort Huachuca	AZ	Santa Rita Mountain American-aster	<i>Symphyotrichum potosinum</i>	Plant	On-site	SAR
Fort Huachuca	AZ	Tepic Flameflower	<i>Talinum marginatum</i>	Plant	On-site	SAR
Fort Huachuca	AZ	Brown Gartersnake	<i>Thamnophis eques megalops</i>	Reptile	On-site	C
Fort Huachuca	AZ	Loach minnow	<i>Tiaroga cobitis</i>	Fish	Contiguous	T
Yuma Proving Ground	AZ	Western Yellow-billed cuckoo	<i>Coccyzus americanus occidentalis</i>	Bird	On-site	T
Yuma Proving Ground	AZ	Desert tortoise (AZ south and east of Colorado River, and Mexico, when found outside of Mexico or said range in AZ)	<i>Gopherus agassizii</i>	Reptile	On-site	T(SA)
Yuma Proving Ground	AZ	Morafka's Desert tortoise	<i>Gopherus morafkai</i>	Turtle	On-site	C
Yuma Proving Ground	AZ	Schott's wire-lettuce	<i>Stephanomeria schottii</i>	Plant	On-site	SAR
Camp Roberts MTCH	CA	Vernal pool fairy shrimp	<i>Branchinecta lynchi</i>	Crustacean	On-site	T
Camp Roberts MTCH	CA	Dwarf Rosinweed	<i>Calycadenia villosa</i>	Plant	On-site	SAR
Camp Roberts MTCH	CA	Hardham's Evening primrose	<i>Camissonia hardhamiae</i>	Plant	On-site	SAR
Camp Roberts MTCH	CA	Purple amole	<i>Chlorogalum purpureum</i>	Plant	On-site	T
Camp Roberts MTCH	CA	One-awned Chorizanthe	<i>Chorizanthe rectispina</i>	Plant	On-site	SAR
Camp Roberts MTCH	CA		<i>Entosthodon kochii</i>	Plant	On-site	SAR

* Species are designated by either E = endangered, T = threatened, UR = under review, C = candidate species, SA = similarity of appearance receives the same protection as listed, or SAR = species at risk. Species are considered to be located on site if they reside within a 2 km distance of the installation (ACSIM 2010, NatureServe 2014).

Installation	State	Common Name	Scientific Name	Taxonomic Group	Location	Status*
Camp Roberts MTC-H	CA	California condor	Gymnogyps californianus	Bird	Contiguous	E
Camp Roberts MTC-H	CA	Santa Lucia Dwarf Rush	Juncus lucensis	Plant	On-site	SAR
Camp Roberts MTC-H	CA	Pale-yellow Layia	Layia heterotricha	Plant	On-site	SAR
Camp Roberts MTC-H	CA	Davidson's Bushmallow	Malacothamnus davidsonii	Plant	On-site	SAR
Camp Roberts MTC-H	CA	Prostrate Navarretia	Navarretia prostrata	Plant	On-site	SAR
Camp Roberts MTC-H	CA	Steelhead (Southern California)	Oncorhynchus mykiss	Fish	On-site	T
Camp Roberts MTC-H	CA	Hooked Popcorn-flower	Plagiobothrys uncinatus	Plant	On-site	SAR
Camp Roberts MTC-H	CA	Thoma's Microseris	Stebbinsoseris decipiens	Plant	On-site	SAR
Camp Roberts MTC-H	CA	Least Bell's vireo	Vireo bellii pusillus	Bird	Contiguous	E
Camp Roberts MTC-H	CA	San Joaquin kit fox	Vulpes macrotis mutica	Mammal	On-site	E
Fort Hunter Liggett	CA	Bristlecone Fir	Abies bracteata	Plant	On-site	SAR
Fort Hunter Liggett	CA	Vernal pool fairy shrimp	Branchinecta lynchi	Crustacean	On-site	T
Fort Hunter Liggett	CA	Arroyo toad	Bufo californicus	Amphibian	On-site	E
Fort Hunter Liggett	CA	Small-flower Calycadenia	Calycadenia micrantha	Plant	On-site	SAR
Fort Hunter Liggett	CA	Dwarf Rosinweed	Calycadenia villosa	Plant	On-site	SAR
Fort Hunter Liggett	CA	Hardham's Evening primrose	Camissonia hardhamiae	Plant	On-site	SAR
Fort Hunter Liggett	CA	Purple amole	Chlorogalum purpureum	Plant	On-site	T
Fort Hunter Liggett	CA	One-awned Chorizanthe	Chorizanthe rectispina	Plant	On-site	SAR
Fort Hunter Liggett	CA	Jolon Clarkia	Clarkia jolonensis	Plant	On-site	SAR
Fort Hunter Liggett	CA	San Antonio Collinsia	Collinsia antonina	Plant	On-site	SAR
Fort Hunter Liggett	CA	Umbrella Larkspur	Delphinium umbraculorum	Plant	On-site	SAR
Fort Hunter Liggett	CA		Didymodon norisii	Mosses	On-site	SAR
Fort Hunter Liggett	CA	Yellow-flower Woolstar	Eriastrum luteum	Plant	On-site	SAR
Fort Hunter Liggett	CA	San Benito Fritillary	Fritillaria viridea	Plant	On-site	SAR
Fort Hunter Liggett	CA	Hardham's Bedstraw	Galium hardhamiae	Plant	On-site	SAR
Fort Hunter Liggett	CA	California condor	Gymnogyps californianus	Bird	On-site	E
Fort Hunter Liggett	CA	Santa Lucia Dwarf Rush	Juncus lucensis	Plant	On-site	SAR
Fort Hunter Liggett	CA	Pale-yellow Layia	Layia heterotricha	Plant	On-site	SAR

* Species are designated by either E = endangered, T = threatened, UR = under review, C = candidate species, SA = similarity of appearance receives the same protection as listed, or SAR = species at risk. Species are considered to be located on site if they reside within a 2 km distance of the installation (ACSIM 2010, NatureServe 2014).

Installation	State	Common Name	Scientific Name	Taxonomic Group	Location	Status*
Fort Hunter Liggett	CA	Abbott's Bushmallow	Malacothamnus abbottii	Plant	On-site	SAR
Fort Hunter Liggett	CA	Davidson's Bushmallow	Malacothamnus davidsonii	Plant	On-site	SAR
Fort Hunter Liggett	CA	Palmer's Monardella	Monardella palmeri	Plant	On-site	SAR
Fort Hunter Liggett	CA	Prostrate Navarretia	Navarretia prostrata	Plant	On-site	SAR
Fort Hunter Liggett	CA	Santa Lucia Pogogyne	Pogogyne clareana	Plant	On-site	SAR
Fort Hunter Liggett	CA	Most Beautiful Jewelflower	Streptanthus albidus ssp. peramoenus	Plant	On-site	SAR
Fort Hunter Liggett	CA	Caper-fruited Tropidocarpum	Tropidocarpum capparideum	Plant	On-site	SAR
Fort Hunter Liggett	CA	Least Bell's vireo	Vireo bellii pusillus	Bird	Contiguous	E
Fort Hunter Liggett	CA	San Joaquin kit fox	Vulpes macrotis mutica	Mammal	On-site	E
Fort Irwin	CA	Lane Mountain milk-vetch	Astragalus jaegerianus	Plant	On-site	E
Fort Irwin	CA	Alkali Mariposally	Calochortus striatus	Plant	On-site	SAR
Fort Irwin	CA	Clokey's Cat's-eye	Cryptantha clokeyi	Plant	On-site	SAR
Fort Irwin	CA	Desert Cymopterus	Cymopterus deserticola	Plant	On-site	SAR
Fort Irwin	CA	Desert tortoise (except AZ south and east of Colorado River, and Mexico)	Gopherus agassizii	Reptile	On-site	T
Fort Irwin	CA	Mohave Ground squirrel	Xeromorphophilus mohavensis	Mammal	On-site	SAR
Parks Reserve Forces Training Area	CA	California Tiger Salamander	Ambystoma Californense	Amphibian	On-site	E
Parks Reserve Forces Training Area	CA	San Joaquin Saltbush	Atriplex joaquiniana	Plant	On-site	SAR
Parks Reserve Forces Training Area	CA	California red-legged frog	Rana aurora draytonii	Amphibian	On-site	T
Parks Reserve Forces Training Area	CA	San Joaquin kit fox	Vulpes Macrotis Mutica	Mammal	On-site	E
Fort Carson	CO	Dwarf milkweed	Asclepias uncialis ssp. uncialis	Plant	On-site	SAR
Fort Carson	CO	Colorado Checkered whiptail	Aspidoscelis neotesselata	Reptile	On-site	SAR
Fort Carson	CO	Front Range milkvetch	Astragalus sparsiflorus	Plant	On-site	SAR
Fort Carson	CO	Arkansas darter	Etheostoma cragini	Fish	On-site	C
Fort Carson	CO	Gold Blazing Star	Mentzelia chrysantha	Plant	On-site	SAR
Fort Carson	CO	Roundleaf Four-o'clock	Mirabilis rotundifolia	Plant	On-site	SAR
Fort Carson	CO	Arkansas Valley Evening primrose	Oenothera harringtonii	Plant	On-site	SAR
Fort Carson	CO	Pueblo goldenweed	Oenopsis sp. 1	Plant	On-site	SAR

* Species are designated by either E = endangered, T = threatened, UR = under review, C = candidate species, SA = similarity of appearance receives the same protection as listed, or SAR = species at risk. Species are considered to be located on site if they reside within a 2 km distance of the installation (ACSIM 2010, NatureServe 2014).

Installation	State	Common Name	Scientific Name	Taxonomic Group	Location	Status*
Fort Carson	CO	Arkansas River feverfew	Parthenium tetraeuris	Plant	On-site	SAR
Fort Carson	CO	Ute Ladies'-tresses	Spiranthes diluvialis	Plant	Contiguous	T
Fort Carson	CO	Mexican spotted owl	Strix occidentalis lucida	Bird	Contiguous	T
Fort Carson	CO	Fendler's Townsend-daisy	Townsendia fendleri	Plant	On-site	SAR
Fort Carson	CO	Pebble's meadow jumping mouse	Zapus hudsonius preblei	Mammal	Contiguous	T
Fort Benning	GA	American alligator	Alligator mississippiensis	Reptile	On-site	T(SA)
Fort Benning	GA	Apalachicola floater	Anodonta heardi	Mollusk	On-site	SAR
Fort Benning	GA	Georgia rock-creep	Arabis georgiana	Plant	On-site	SAR
Fort Benning	GA	Fly's Brickell-bush	Brickellia cordifolia	Plant	On-site	SAR
Fort Benning	GA	Bluestripe shiner	Cyprinella callitaenia	Fish	On-site	SAR
Fort Benning	GA	Gopher tortoise	Gopherus polyphemus	Turtle	On-site	C
Fort Benning	GA	Barbour's Map turtle	Graptemys barbouri	Turtle	On-site	SAR
Fort Benning	GA	Smith's sunflower	Helianthus smithii	Plant	On-site	SAR
Fort Benning	GA	Southern hog-nosed snake	Heterodon simus	Reptile	On-site	SAR
Fort Benning	GA	Boykin's Lobelia	Lobelia boykinii	Plant	On-site	SAR
Fort Benning	GA	Spathulate Seedbox	Ludwigia spathulata	Plant	On-site	SAR
Fort Benning	GA	Wood stork	Mycteria americana	Bird	On-site	E
Fort Benning	GA	Halloween darter	Percina crypta	Fish	On-site	SAR
Fort Benning	GA	Red-cockaded woodpecker	Picoides borealis	Bird	On-site	E
Fort Benning	GA	Relict trillium	Trillium reliquum	Plant	On-site	E
Fort Gordon	GA	Smooth cone-flower	Echinacea laevigata	Plant	On-site	E
Fort Gordon	GA	Bluebarred Pygmy sunfish	Elassoma okatie	Fish	On-site	SAR
Fort Gordon	GA	Gopher tortoise	Gopherus polyphemus	Turtle	On-site	C
Fort Gordon	GA	Southern hog-nosed snake	Heterodon simus	Reptile	On-site	SAR
Fort Gordon	GA	Rough-leaved loosestrife	Lysimachia asperulaefolia	Plant	On-site	E
Fort Gordon	GA	Carolina Birds-in-a-nest	Macbridea caroliniana	Plant	On-site	SAR
Fort Gordon	GA	Wood stork	Mycteria americana	Bird	Contiguous	E
Fort Gordon	GA	Indiana bat	Myotis sodalis	Mammal	On-site	E

* Species are designated by either E = endangered, T = threatened, UR = under review, C = candidate species, SA = similarity of appearance receives the same protection as listed, or SAR = species at risk. Species are considered to be located on site if they reside within a 2 km distance of the installation (ACSIM 2010, NatureServe 2014).

Installation	State	Common Name	Scientific Name	Taxonomic Group	Location	Status*
Fort Gordon	GA	Red-cockaded woodpecker	Picoides borealis	Bird	On-site	E
Fort Stewart	GA	Shortnose sturgeon	Acipenser brevirostrum	Fish	On-site	E
Fort Stewart	GA	Flatwoods salamander	Ambystoma cingulatum	Amphibian	On-site	T
Fort Stewart	GA	Purple Baldpate	Baldpate atropurpurea	Plant	On-site	SAR
Fort Stewart	GA	Autumn Tiger beetle	Cicindela nigror	Invertebrate	On-site	SAR
Fort Stewart	GA	Say's Spiketail	Cordulegaster sayi	Invertebrate	On-site	SAR
Fort Stewart	GA	Eastern indigo snake	Drymarchon corais couperi	Reptile	On-site	T
Fort Stewart	GA	Gopher tortoise	Gopherus polyphemus	Turtle	On-site	C
Fort Stewart	GA	Southern hog-nosed snake	Heterodon simus	Reptile	On-site	SAR
Fort Stewart	GA	Yellow Anisetree	Illicium parviflorum	Plant	On-site	SAR
Fort Stewart	GA	Boykin's Lobelia	Lobelia boykinii	Plant	On-site	SAR
Fort Stewart	GA	Robust Redhorse	Moxostoma robustum	Fish	On-site	SAR
Fort Stewart	GA	Wood stork	Mycteria americana	Bird	On-site	E
Fort Stewart	GA	Red-cockaded woodpecker	Picoides borealis	Bird	On-site	E
Fort Stewart	GA	Wireleaf Dropseed	Sporobolus teretifolius	Plant	On-site	SAR
Fort Stewart	GA	Savannah Liliuput	Toxolasma pullus	Mollusk	On-site	SAR
Camp Dodge Joint Maneuver Training Center	IA	Northern long-eared bat	Myotis septentrionalis	Mammal	On-site	T
Camp Atterbury	IN	Kirtland's snake	Clonophis kirtlandii	Reptile	On-site	UR
Camp Atterbury	IN	Snuffbox	Epioblasma triquetra	Mollusk	On-site	E
Camp Atterbury	IN	Northern long-eared bat	Myotis septentrionalis	Mammal	On-site	T
Camp Atterbury	IN	Indiana bat	Myotis sodalis	Mammal	On-site	E
Camp Atterbury	IN	Rayed bean	Villosa fabalis	Clam	On-site	E
Fort Riley	KS	Piping plover	Charadrius melodus	Bird	On-site	T
Fort Riley	KS	Northern long-eared bat	Myotis septentrionalis	Mammal	On-site	T
Fort Riley	KS	Topeka shiner	Notropis topeka	Fish	On-site	E
Fort Riley	KS	Salina Dewberry	Rubus hancianus	Plant	On-site	SAR
Fort Riley	KS	Regal fritillary butterfly	Speyeria idalia	Insect	On-site	SAR

* Species are designated by either E = endangered, T = threatened, UR = under review, C = candidate species, SA = similarity of appearance receives the same protection as listed, or SAR = species at risk. Species are considered to be located on site if they reside within a 2 km distance of the installation (ACSIM 2010, NatureServe 2014).

Installation	State	Common Name	Scientific Name	Taxonomic Group	Location	Status*
Fort Riley	KS	Least tern	<i>Sterna antillarum</i>	Bird	On-site	E
Fort Campbell	KY	Gray bat	<i>Myotis grisescens</i>	Mammal	On-site	E
Fort Campbell	KY	Indiana bat	<i>Myotis sodalis</i>	Mammal	On-site	E
Fort Campbell	KY	Coleman Cave beetle	<i>Pseudanophthalmus colmanensis</i>	Invertebrate	On-site	C
Fort Campbell	KY	Hall's Bulrush	<i>Schoenoplectus hallii</i>	Plant	On-site	SAR
Fort Knox	KY	Kirtland's snake	<i>Clonophis kirtlandii</i>	Reptile	On-site	SAR
Fort Knox	KY	Gray bat	<i>Myotis grisescens</i>	Mammal	On-site	E
Fort Knox	KY	Northern long-eared bat	<i>Myotis septentrionalis</i>	Mammal	On-site	T
Fort Knox	KY	Louisville crayfish	<i>Orconectes jeffersoni</i>	Invertebrate	On-site	SAR
Fort Knox	KY	Blazing Star Stem Borer	<i>Papaipema beeriana</i>	Invertebrate	On-site	SAR
Fort Polk	LA	Southern Hickorynut	<i>Obovaria jacksoniana</i>	Mollusk	On-site	SAR
Fort Polk	LA	Kisatchie Painted crayfish	<i>Orconectes maletae</i>	Invertebrate	On-site	SAR
Fort Polk	LA	Red-cockaded woodpecker	<i>Picoides borealis</i>	Bird	On-site	E
Fort Polk	LA	Louisiana pine snake	<i>Pituophis ruthveni</i>	Reptile	On-site	C
Aberdeen Proving Ground	MD	Shorthose sturgeon	<i>Acipenser brevirostrum</i>	Fish	On-site	E
Aberdeen Proving Ground	MD	Puritan Tiger beetle	<i>Cicindela puritana</i>	Insect	Contiguous	E
Aberdeen Proving Ground	MD	Maryland darter	<i>Etheostoma sellare</i>	Fish	Contiguous	E
Aberdeen Proving Ground	MD	Chesapeake Logperch	<i>Percina bimaculata</i>	Fish	On-site	SAR
Camp Grayling MTC-H	MI	Kirtland's warbler	<i>Dendroica kirtlandii</i>	Bird	On-site	E
Camp Grayling MTC-H	MI	Appalachian Grizzled Skipper	<i>Pyrgus wyandot</i>	Plant	On-site	SAR
Camp Grayling MTC-H	MI	Eastern Massasauga snake	<i>Sistrurus catenatus catenatus</i>	Reptile	On-site	C
Camp Grayling MTC-H	MI	Houghton's goldenrod	<i>Solidago houghtonii</i>	Plant	On-site	T
Fort Ouster Training Center CTC	MI	Northern Long-eared bat	<i>Myotis septentrionalis</i>	Mammal	On-site	T
Fort Ouster Training Center CTC	MI	Eastern Massasauga snake	<i>Sistrurus catenatus catenatus</i>	Reptile	On-site	C
Fort Leonard Wood	MO	Hubricht's Long-tailed Amphipod	<i>Alloicranonyx hubrichti</i>	Invertebrate	On-site	SAR
Fort Leonard Wood	MO	Spectaclecase	<i>Cumberlandia Monodonta</i>	Mollusk	On-site	E
Fort Leonard Wood	MO	Gray bat	<i>Myotis grisescens</i>	Mammal	On-site	E
Fort Leonard Wood	MO	Northern long-eared bat	<i>Myotis septentrionalis</i>	Mammal	On-site	T

* Species are designated by either E = endangered, T = threatened, UR = under review, C = candidate species, SA = similarity of appearance receives the same protection as listed, or SAR = species at risk. Species are considered to be located on site if they reside within a 2 km distance of the installation (ACSIM 2010, NatureServe 2014).

Installation	State	Common Name	Scientific Name	Taxonomic Group	Location	Status*
Fort Leonard Wood	MO	Indiana bat	Myotis sodalis	Mammal	On-site	E
Fort Leonard Wood	MO	Bluestripe darter	Percina cyathotaenia	Fish	On-site	SAR
Fort William Henry Harrison MTA	MT	NA	NA	NA	NA	NA
Fort Bragg	NC	Dusky Roadside-Skipper	Amblyscirtes alternata	Invertebrate	On-site	SAR
Fort Bragg	NC	Savannah Campylopus	Campylopus caroliniae	Moss	On-site	SAR
Fort Bragg	NC	Thinlip chub	Cyprinella sp. 1	Fish	On-site	SAR
Fort Bragg	NC	Hidden-flower Witchgrass	Dichanthelium cryptanthum Sp. 9	Plant	On-site	SAR
Fort Bragg	NC	Southern hog-nosed snake	Heterodon simus	Reptile	On-site	SAR
Fort Bragg	NC	Winter quillwort	Isoetes hyemalis	Plant	On-site	SAR
Fort Bragg	NC	Panhandle Lily	Lilium iridollae	Plant	On-site	SAR
Fort Bragg	NC	Sandhills Lily	Lilium pyrophilum	Plant	On-site	SAR
Fort Bragg	NC	Bog Spicebush	Lindera subcoriacea	Plant	On-site	UR
Fort Bragg	NC	Boykin's Lobelia	Lobelia boykinii	Plant	On-site	SAR
Fort Bragg	NC	Rough-leaved loosestrife	Lysimachia asperulaefolia	Plant	On-site	E
Fort Bragg	NC		Melanoplus nubilus	Invertebrate	On-site	SAR
Fort Bragg	NC	Northern long-eared bat	Myotis septentrionalis	Mammal	On-site	T
Fort Bragg	NC	Saint Francis' satyr butterfly	Neonympha mitchellii francisci	Insect	On-site	E
Fort Bragg	NC	Red-cockaded woodpecker	Picoides borealis	Bird	On-site	E
Fort Bragg	NC	Michaux's sumac	Rhus michauxii	Plant	On-site	E
Fort Bragg	NC	Hairy-peduncled beakrush	Rhynchospora crinipes	Plant	On-site	UR
Fort Bragg	NC	American chaffseed	Schwalbea americana	Plant	On-site	E
Fort Dix	NJ	Buchholz's Dart Moth	Agrotis buchholzi	Insect	On-site	SAR
Fort Dix	NJ		Richia sp. 2	Insect	On-site	SAR
White Sands Military Range	NM	Cliff bristlebush	Apacheria chricahuensis	Plant	On-site	SAR
White Sands Military Range	NM	Goat Mountain Woodlandsnail	Ashmunella harrisi	Mollusk	On-site	SAR
White Sands Military Range	NM	Gray wolf	Canis lupus	Mammal	Contiguous	E
White Sands Military Range	NM	Samalayuca Dune grasshopper	Chloracris samalayuca	Invertebrate	On-site	SAR
White Sands Military Range	NM	White Sands pupfish	Cyprinodon tularosa	Fish	On-site	UR

* Species are designated by either E = endangered, T = threatened, UR = under review, C = candidate species, SA = similarity of appearance receives the same protection as listed, or SAR = species at risk. Species are considered to be located on site if they reside within a 2 km distance of the installation (ACSIM 2010, NatureServe 2014).

Installation	State	Common Name	Scientific Name	Taxonomic Group	Location	Status*
White Sands Military Range	NM	Southwestern willow flycatcher	<i>Empidonax traillii extimus</i>	Bird	On-site	E
White Sands Military Range	NM	Organ Mountain Foxtail Cactus	<i>Escobaria organensis</i>	Plant	On-site	SAR
White Sands Military Range	NM	San Andres Mountain Foxtail Cactus	<i>Escobaria sandbergii</i>	Plant	On-site	SAR
White Sands Military Range	NM	Northern Aplomado falcon	<i>Falco femoralis septentrionalis</i>	Bird	On-site	E
White Sands Military Range	NM	Poling's Hairstreak	<i>Ficksenia polingii</i>	Insect	On-site	SAR
White Sands Military Range	NM	Bolson tortoise	<i>Gopherus flavomarginatus</i>	Reptile	Contiguous	E
White Sands Military Range	NM	White Mountain False Pennyroyal	<i>Hedeoma pulcherrima</i>	Plant	On-site	SAR
White Sands Military Range	NM	Todsen's Pennyroyal	<i>Hedeoma todsenii</i>	Plant	On-site	E
White Sands Military Range	NM	Vasey's Bitterweed	<i>Hymenoxys vaseyi</i>	Plant	On-site	SAR
White Sands Military Range	NM	Organ Mountain Evening primrose	<i>Oenothera organensis</i>	Plant	On-site	SAR
White Sands Military Range	NM	Jaguar	<i>Panthera onca</i>	Mammal	On-site	E
White Sands Military Range	NM	Brown pelican	<i>Pelecanus occidentalis</i>	Bird	On-site	E
White Sands Military Range	NM	Least tern	<i>Sterna antillarum</i>	Bird	On-site	E
White Sands Military Range	NM	Mexican spotted owl	<i>Strix occidentalis lucida</i>	Bird	Contiguous	T
Fort Drum	NY	Northern long-eared bat	<i>Myotis septentrionalis</i>	Mammal	On-site	T
Fort Drum	NY	Indiana bat	<i>Myotis sodalis</i>	Mammal	On-site	E
Fort Drum	NY	Tomah mayfly	<i>Siphonisca aerodromia</i>	Invertebrate	On-site	SAR
Fort Sill	OK	Taper-tip Dodder	<i>Cuscuta attenuata</i>	Plant	On-site	SAR
Fort Sill	OK	Black-capped vireo	<i>Vireo atricapilla</i>	Bird	On-site	E
Fort Jackson	SC	Purple Baldpate	<i>Baldpate atropurpurea</i>	Plant	On-site	SAR
Fort Jackson	SC	Southern hog-nosed snake	<i>Heterodon simus</i>	Reptile	On-site	SAR
Fort Jackson	SC	Sandhills lily	<i>Lilium pyrophilum</i>	Plant	On-site	SAR
Fort Jackson	SC	Bog Spicebush	<i>Lindera subcoriacea</i>	Plant	On-site	SAR
Fort Jackson	SC	Rough-leaved loosestrife	<i>Lysimachia asperulaefolia</i>	Plant	On-site	E
Fort Jackson	SC	Red-cockaded woodpecker	<i>Picoides borealis</i>	Bird	On-site	E
Camp Bullis	TX	Texas salamander	<i>Eurycea neotenes</i>	Amphibian	On-site	SAR
Camp Bullis	TX	Comal Blind salamander	<i>Eurycea tridentifera</i>	Amphibian	On-site	SAR
Camp Bullis	TX	Bracted Twistflower	<i>Streptanthus bracteatus</i>	Plant	On-site	C

* Species are designated by either E = endangered, T = threatened, UR = under review, C = candidate species, SA = similarity of appearance receives the same protection as listed, or SAR = species at risk. Species are considered to be located on site if they reside within a 2 km distance of the installation (ACSIM 2010, NatureServe 2014).

Installation	State	Common Name	Scientific Name	Taxonomic Group	Location	Status*
Fort Bliss	TX	Cliff brittlebush	Apacheria chricahuensis	Plant	On-site	SAR
Fort Bliss	TX	Prickly-poppy, Sacramento	Argemone pleiacantha ssp. pinnatisecta	Plant	Contiguous	E
Fort Bliss	TX	Boulder Canyon Woodland snail	Ashmunella auriculata	Mollusk	On-site	SAR
Fort Bliss	TX	Goat Mountain Woodlandsnail	Ashmunella harrisi	Plant	On-site	SAR
Fort Bliss	TX	Organ Mountain Woodlands snail	Ashmunella organensis	Plant	On-site	SAR
Fort Bliss	TX	Maple Canyon Woodlands snail	Ashmunella todsoni	Mollusk	On-site	SAR
Fort Bliss	TX	Piping plover	Charadrius melodus	Bird	On-site	T
Fort Bliss	TX	Mountain plover	Charadrius montanus	Bird	On-site	T
Fort Bliss	TX	Samalayuca Dune grasshopper	Cibolacris samalayuca	Insect	On-site	SAR
Fort Bliss	TX	Yellow-billed Cuckoo [western distinct population]	Coccyzus americanus	Bird	On-site	T
Fort Bliss	TX	Cactus, Sneed pincushion	Coryphantha sneedii var. sneedii	Plant	On-site	E
Fort Bliss	TX	Kuenzler hedgehog cactus	Echinocereus fendleri var. kuenzleri	Plant	Contiguous	E
Fort Bliss	TX	Southwestern Willow flycatcher	Empidonax traillii eximius	Bird	Contiguous	E
Fort Bliss	TX	Organ Mountain Pincushion cactus	Escobaria organensis	Plant	On-site	SAR
Fort Bliss	TX	Northern Aplomado falcon	Falco femoralis septentrionalis	Bird	On-site	E
Fort Bliss	TX	Anthony Blister Beetle	Lytta mirifica	Insect	On-site	SAR
Fort Bliss	TX	Organ mountain Evening primrose	Oenothera organensis	Plant	On-site	SAR
Fort Bliss	TX	Sand prickly pear	Opuntia arenaria	Plant	On-site	SAR
Fort Bliss	TX	Nodding rock daisy	Perityle cernua	Plant	On-site	SAR
Fort Bliss	TX	Hueco Mountains Rockdaisy	Perityle huecoensis	Plant	On-site	SAR
Fort Bliss	TX	Smooth figwort	Scrophularia laevis	Plant	On-site	SAR
Fort Bliss	TX	Plank's Catchfly	Silene plankii	Plant	On-site	SAR
Fort Bliss	TX	Franklin Mountain Talussnail	Sonorella metcalfi	Mollusk	On-site	SAR
Fort Bliss	TX	Interior least tern	Sterna antillarum athalassos	Bird	Contiguous	E
Fort Hood	TX	Texabama Croton	Croton alabamensis texensis	Plant	On-site	SAR
Fort Hood	TX	Golden-cheeked warbler	Dendroica chrysoparia	Bird	On-site	E
Fort Hood	TX	Whooping crane	Grus americana	Bird	On-site	E
Fort Hood	TX	Smooth Pimpleback	Quadrula houstonensis	Mollusk	On-site	SAR

* Species are designated by either E = endangered, T = threatened, UR = under review, C = candidate species, SA = similarity of appearance receives the same protection as listed, or SAR = species at risk. Species are considered to be located on site if they reside within a 2 km distance of the installation (ACSIM 2010, NatureServe 2014).

Installation	State	Common Name	Scientific Name	Taxonomic Group	Location	Status*
Fort Hood	TX	Black-capped vireo	Vireo atricapilla	Bird	On-site	E
Camp Williams MTA-L	UT	Green River Pebblesnail	Fluminicola coloradoensis	Mollusk	On-site	SAR
Dugaway Proving Ground	UT	Western snowy plover	Charadrius alexandrinus nivosus	Bird	On-site	T
Dugaway Proving Ground	UT	Least Chub	lotichthys phlegethontis	Fish	On-site	SAR
Dugaway Proving Ground	UT	Oregon spotted frog	Rana pretiosa	Amphibian	Contiguous	T
Dugaway Proving Ground	UT	Ladies-tresses Ute	Spiranthes diluvialis	Plant	Contiguous	T
Fort A.P. Hill	VA	Swamp pink	Helonias bullata	Plant	On-site	T
Fort A.P. Hill	VA	Small whorled pogonia	Isotria medeoloides	Plant	On-site	T
Fort A.P. Hill	VA	Rappahannock Spring amphipod	Stygobromus sp. 21	Invertebrate	On-site	SAR
Fort Lee	VA	Sensitive joint-veitch	Aeschynomene virginica	Plant	On-site	T
Fort Lee	VA	Northern Long-eared bat	Myotis septentrionalis	Mammal	On-site	T
Fort Pickett MTC	VA	Sensitive Freshwater mussels	Fusconia masoni	Mollusk	On-site	SAR
Fort Pickett MTC	VA	Roanoke logperch	Percina rex	Fish	On-site	E
Fort Pickett MTC	VA	Torrey's mountain mint	Pycnanthemum torrei	Plant	On-site	SAR
Fort Pickett MTC	VA	Michaux's sumac	Rhus michauxii	Plant	On-site	E
Joint Base Langley - Eustis	VA	Atlantic sturgeon	Acipenser oxyrinchus	Fish	On-site	E
Joint Base Lewis-McCord	WA	Streaked horned lark	Eremophila alpestris strigata	Bird	On-site	T
Joint Base Lewis-McCord	WA	Taylor's checkerspot butterfly	Euphydryas editha taylori	Insect	On-site	C
Joint Base Lewis-McCord	WA	Water Howellia	Howellia aquatilis	Plant	On-site	T
Joint Base Lewis-McCord	WA	Steelhead (Puget Sound Distinct Population Segment, DPS)	Oncorhynchus mykiss	Fish	Contiguous	T
Joint Base Lewis-McCord	WA	Chinook salmon (naturally spawned populations from rivers and streams flowing into Puget Sound)	Oncorhynchus tshawytscha	Fish	On-site	T
Joint Base Lewis-McCord	WA	Mardon skipper	Polites mardon	Insect	On-site	C
Joint Base Lewis-McCord	WA	Oregon spotted frog	Rana pretiosa	Amphibian	On-site	T
Joint Base Lewis-McCord	WA	Bull trout	Salvelinus confluentus	Fish	On-site	T
Joint Base Lewis-McCord	WA	Bocaccio rockfish	Sebastes paucispinis	Fish	Contiguous	E
Joint Base Lewis-McCord	WA	Canary rockfish	Sebastes pinniger	Fish	On-site	T
Joint Base Lewis-McCord	WA	Yelloweye rockfish	Sebastes ruberrimus	Fish	On-site	T

* Species are designated by either E = endangered, T = threatened, UR = under review, C = candidate species, SA = similarity of appearance receives the same protection as listed, or SAR = species at risk. Species are considered to be located on site if they reside within a 2 km distance of the installation (ACSIM 2010, NatureServe 2014).

Installation	State	Common Name	Scientific Name	Taxonomic Group	Location	Status*
Joint Base Lewis-McCord	WA	Northern spotted owl	<i>Strix occidentalis caurina</i>	Bird	Contiguous	T
Joint Base Lewis-McCord	WA	Roy prairie pocket gopher	<i>Thomomys mazama glacialis</i>	Mammal	On-site	T
Joint Base Lewis-McCord	WA	Olympia pocket gopher	<i>Thomomys mazama pugetensis</i>	Mammal	On-site	T
Joint Base Lewis-McCord	WA	Tenino pocket gopher	<i>Thomomys mazama tumuli</i>	Mammal	On-site	T
Joint Base Lewis-McCord	WA	Yelm pocket gopher	<i>Thomomys mazama yelmensis</i>	Mammal	On-site	T
Joint Base Lewis-McCord	WA	Small-flowered Trillium	<i>Trillium parviflorum</i>	Plant	On-site	SAR
Yakima Training Center	WA	Northern Womwood	<i>Artemisia campestris</i> var. <i>wormskioildii</i>	Plant	On-site	SAR
Yakima Training Center	WA	Gray Cryptantha	<i>Cryptantha leucophaea</i>	Plant	On-site	SAR
Yakima Training Center	WA	Basalt Daisy	<i>Erigeron basalticus</i>	Plant	On-site	SAR
Yakima Training Center	WA	Hoover's Desert-parsley	<i>Lomatium tuberosum</i>	Plant	On-site	SAR
Yakima Training Center	WA	Steelhead (Upper Columbia River)	<i>Oncorhynchus mykiss</i> pop. 12	Fish	On-site	T
Yakima Training Center	WA	Steelhead (Middle Columbia River)	<i>Oncorhynchus mykiss</i> pop. 17	Fish	Contiguous	T
Yakima Training Center	WA	Chinook salmon (naturally spawned populations in the Columbia R. tributaries upstream of Rock Island Dam and downstream of Chief Joseph Dam)	<i>Oncorhynchus tshawytscha</i>	Fish	On-site	T
Yakima Training Center	WA	Bull trout	<i>Salvelinus confluentus</i>	Fish	Contiguous	T
Yakima Training Center	WA	Hoover's Tauschia	<i>Tauschia hooveri</i>	Plant	On-site	SAR
Fort McCoy	WI	Red-Tailed Leafhopper	<i>Afelia rubranura</i>	Invertebrate	On-site	SAR
Fort McCoy	WI	Gray wolf	<i>Canis lupus</i>	Mammal	On-site	T
Fort McCoy	WI	Karner blue butterfly	<i>Lyciaides melissa samuelis</i>	Insect	On-site	E
Fort McCoy	WI	Eastern Massasauga snake	<i>Sistrurus catenatus catenatus</i>	Reptile	On-site	C

* Species are designated by either E = endangered, T = threatened, UR = under review, C = candidate species, SA = similarity of appearance receives the same protection as listed, or SAR = species at risk. Species are considered to be located on site if they reside within a 2 km distance of the installation (ACSIM 2010, NatureServe 2014).

Appendix C: Installation Vulnerabilities Calculated Without the Installation Rank Factor

Table C-1. Climate change vulnerability scores for 43 CONUS ITAM installations based on four factors, excluding the installation rank factor. Installations are ordered high-to-low based on vulnerability. Blue = minimum, white = median (50th percentile), red = maximum, and transitional colors represent intermediate percentiles.

Installation	State	Vulnerability
Parks Reserve Forces Training Area	CA	1.047
Joint base Lewis-McCord	WA	0.953
Fort Huachuca	AZ	0.935
Fort Bliss	TX	0.751
Camp Atterbury	IN	0.730
MTC-H Camp Roberts	CA	0.720
Fort Carson	CO	0.688
Fort Bragg	NC	0.663
Fort Sill	OK	0.660
Fort Eustis	VA	0.659
Fort Riley	KS	0.656
White Sands Military Range	NM	0.653
CTC Fort Custer TS	MI	0.646
Camp Bullis	TX	0.643
Yakima Training Center	WA	0.638
Fort Campbell	KY	0.637
Aberdeen Proving Ground	MD	0.637
Fort Hood	TX	0.634
Fort Benning	GA	0.626
Fort Gordon	GA	0.620
Fort Stewart	GA	0.619
Camp Dodge Joint Maneuver Training Center	IA	0.603
Fort Hunter Liggett	CA	0.594
Fort Knox	KY	0.588
Fort Rucker	AL	0.573
Fort McCoy	WI	0.567
MTA-L Camp Williams	UT	0.550
Camp Joseph T Robinson	AR	0.528
MTC-H Camp Grayling	MI	0.511
Fort Jackson	SC	0.510

Installation	State	Vulnerability
Fort Leonard Wood	MO	0.502
MTA Fort Wm Henry Harrison	MT	0.501
Fort Lee	VA	0.496
Fort Chaffee MTC	AR	0.486
Fort Dix	NJ	0.482
Fort Pickett, ARNG MTC	VA	0.477
Fort A P Hill	VA	0.473
Fort Drum	NY	0.472
Fort Polk	LA	0.465
Yuma Proving Ground	AZ	0.463
Dugway Proving Ground	UT	0.435
Camp Navajo	AZ	0.410
Fort Irwin	CA	0.378

References

- Abatzoglou, J. T., and C. A. Kolden. 2011. Climate change in western US deserts: Potential for increased wildfire and invasive annual grasses. *Rangeland Ecology and Management* 64:471-478.
- Assistant Chief of Staff for Installation Management (ACSIM). 2010. *Threatened, Endangered, and at-Risk Species Interagency Research Coordination Manual*, http://www.dodnaturalresources.net/files/TER-S_Interagency_Research_Coordination_Manual.pdf
- Bagne, K. E., M. M. Friggens, and D. M. Finch. 2011. *A System for Assessing Vulnerability of Species (SAVS) To Climate Change*. General Technical Report RMRS-GTR-257. Fort. Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station.
- Bierwagen, Britta, and Phil Morefield. 2014. *Integrated Climate and Land Use Scenarios (ICLUS)*. Accessed September 25, <http://134.67.99.51/ICLUSonline/>
- Cox, W. A., F. R. Thompson III, and J. L. Reidy. 2013. The Effects of temperature on nest predation by mammals, birds, and snakes. *The Auk* 130:784-790. doi: <http://dx.doi.org/10.1525/auk.2013.13033>
- Dawson, T. P., S. T. Jackson, J. I. House, I. C. Prentice, and G. M. Mace. 2011. Beyond predictions: Biodiversity conservation in a changing climate. *Science* 332:53-58.
- Deal, B., and V. Pallathucheril. 2009. A use-driven approach to large-scale urban modelling and planning support. *Planning Support Systems Best Practice and New Methods*, pp 29–51. Springer Netherlands, http://link.springer.com.proxy2.library.illinois.edu/chapter/10.1007/978-1-4020-8952-7_2
- Eastman, J. R., P. A. K. Kyem, J. Toledano, and W. Jin. 1993. *GIS and Decision Making*. Geneva: UNITAR.
- Ellenwood, M. S., L. Dilling, and J.B. Milford. 2012. Managing United States public lands in response to climate change: A view from the ground up. *Environmental Management* 49:954-967.
- Environmental Systems Research Institute, Inc. (ESRI). 2016. ArcGIS 10.2.2 (ESRI, Inc., Redlands, CA).
- Hellmann, J. J., J. E. Byers, B. G. Bierwagen, and J. S. Dukes. 2008. Five potential consequences of climate change for invasive species. *Conservation Biology* 22:534-543.
- Hohmann, M. G., and W. A. Wall. 2017. Multiscale Assessment of Listed and At-Risk Species: Climate Change Vulnerabilities. ERDC/CERL TR-17-21. Champaign, IL: Engineer Research and Development Center, Construction Engineering Research Laboratory (ERDC-CERL).

- Homer, C. G., J. A. Dewitz, L. Yang, S. Jin, P. Danielson, G. Xian, J. Coulston, N.D. Herold, J.D. Wickham, and K. Megown. 2015. Completion of the 2011 National Land Cover Database for the conterminous United States-Representing a decade of land cover change information. *Photogrammetric Engineering and Remote Sensing* 81:345-354.
- Janzen, F. J. 1994. Climate change and temperature-dependent sex determination in reptiles. *Proceedings of the National Academy of Sciences* 91:7487-7490.
- Jenkinsa, C. N., K. S. Van Houtanb, S. L. Pimm, and J. O. Sexton. 2015. US protected lands mismatch biodiversity priorities. *Proceedings of the National Academy of Science* 112:5081-5086.
- Kirkwood, C. W. 1997. *Strategic Decision Making: Multiobjective Decision Analysis with Spreadsheets*. Belmont, CA: Brooks/Cole, Cengage Learning.
- Krasting, J. P., J. P. Dunne, R. J. Stouffer, and R. W. Hallberg. 2016. Enhanced Atlantic sea-level rise relative to the Pacific under high carbon emission rates. *Nature Geoscience* 9:210-214.
- Loarie, S. R., P. B. Duffy, H. Hamilton, G. P. Asner, C. B. Field, and D. D. Ackerly. 2009. The velocity of climate change. *Nature* 462:1052-1057. doi:10.1038/nature08649
- Malczewski, J. 2000. On the use of weighted linear combination method in GIS: Common and best practice approaches. *Transactions in GIS* 4:5-22.
- Martinuzzi, S., V. C. Radeloff, L. N. Joppa, C. M. Hamilton, D. P. Helmers, A. J. Plantinga, and D. J. Lewis. 2015. Scenarios of future land use change around US protected areas. *Biological Conservation* 184:446-455.
- Messer, K. D., M. Kecinski, Z. Liu, M. A. Korch, and T. Bounds. 2016. Military readiness and environmental protection through cost-effective land conservation. *Land Economics* 92:433-449.
- NatureServe. 2014. *Species at Risk on Department of Defense Lands: Updated Analysis, Report and Maps*. DoD Legacy Project 10-247.
- Ordóñez, A., S. Martinuzzi, V. Radeloff, and J. W. Williams. 2014. Combined speeds of climate and land-use change of the conterminous US until 2050. *Nature Climate Change* 4:811-816.
- Shaw, M. R., K. Klausmeyer, D. R. Cameron, J. Mackenzie, and P. Roehrdanz. 2012. Economic costs of achieving current conservation goals in the future as climate changes. *Conservation Biology* 26:385-396.
- Sperry, J. H., W. A. Wall, and M. G. Hohmann. 2016. *Evaluation of 757 Species Under U.S. Endangered Species Act Review on Department of Defense Lands and Their Potential Impact on Army Training*. ERDC/CERL TR-16-3. Champaign, IL: ERDC-CERL.
- Stein, B. A., C. Scott, and N. Benton. 2008. Federal lands and endangered species: The role of military and other federal lands in sustaining biodiversity. *BioScience* 58:339-347.

- Thieler, E. R., and E. S. Hammar-Klose. 2000. *National Assessment of Coastal Vulnerability to Sea-Level Rise: Preliminary Results for the U.S. Atlantic Coast*. U.S. Geological Survey Open-File Report 99-593.
- Tomanek, L. 2008. The importance of physiological limits in determining biogeographical range shifts due to global climate change: The heat-shock response. *Physiological and Biochemical Zoology* 81:709-717.
- Urban, M. C. 2015. Accelerating extinction risk from climate change. *Science* 348:571-573.
- U.S. Department of Defense (DoD). 2010. *Quadrennial Defense Review*. Washington, DC: DoD, http://www.defense.gov/Portals/1/features/defenseReviews/QDR/QDR_as_of_29JAN10_1600.pdf
- . 2011. DoD Instruction (DODI) 4715.03. Subject: DoD Instruction. Washington, DC: Office of the Under Secretary of Defense, Acquisition, Technology and Logistics (OUSD[AT&L]), <http://www.dtic.mil/whs/directives/corres/pdf/471503p.pdf>
- . 2012. *2012 Climate Change Adaptation Roadmap*. Washington, DC: DoD.
- . 2013. U.S. Department of Defense Manual. Subject: Integrated Natural Resources Management Plan (INRMP) Implementation Manual. DODM 4715.03. Washington, DC: OUSD(AT&L), <http://www.dtic.mil/whs/directives/corres/pdf/471503m.pdf>
- . 2014a. *2014 Climate Change Adaptation Roadmap*. Washington, DC: DoD, <http://reliefweb.int/sites/reliefweb.int/files/resources/CCARprint.pdf>
- U.S. Environmental Protection Agency (USEPA). 2009. *A Framework for Categorizing the Relative Vulnerability of Threatened and Endangered Species to Climate Change*. EPA/600/R-09/011. Washington, DC: National Center for Environmental Assessment, available from the National Technical Information Service, Springfield, VA and online at <http://www.epa.gov/ncea>.
- U.S. Government Accountability Office (GAO). 2009. *Climate Change Adaptation: Strategic Federal Planning Could Help Government Officials Make More Informed Decisions*. GAO-10-113. Washington, DC: GAO.
- . 2014. *Climate Change Adaptation: DoD Can Improve Infrastructure Planning and Process to Better Account for Potential Impacts*. GAO-14-446. Washington, DC: GAO, <http://www.gao.gov/assets/670/663734.pdf>
- . 2014. *Climate Change Adaptation: DOD Can Improve Infrastructure Planning and Process to Better Account for Potential Impacts*. GAO-14-446, 58 pp.
- Vale, C. G., and J. C. Brito. 2015. Desert-adapted species are vulnerable to climate change: Insights from the warmest region on Earth. *Global Ecology and Conservation* 4:369-379.
- Westervelt, J., T. BenDor, and J. Sexton. 2011. A technique for rapidly forecasting regional urban growth. *Environment and Planning B: Planning and Design* 38:61–81. doi:10.1068/b36029.

- White House, The. 2009. *Federal Leadership in Environmental, Energy and Economic Performance*. Executive Order (EO) 13514. Washington, DC: The White House, Office of the Press Secretary, <https://www.whitehouse.gov/the-press-office/president-obama-signs-executive-order-focused-federal-leadership-environmental-ener>
- White House, The. 2013. *Preparing the United States for the Impacts of Climate Change*. EO 13653. Washington, DC: The White House, Office of the Press Secretary, <https://www.whitehouse.gov/the-press-office/2013/11/01/executive-order-preparing-united-states-impacts-climate-change>
- Wilhoit, J., S. Tweddale, M. G. Hohmann, D. Delaney, M. E. Swearingen, and J. Westervelt. 2015. *Effects of Climate Change, Urban Development and Threatened and Endangered Species Management on Army Training Capabilities*. ERDC/CERL TR-16-29. Champaign, IL: ERDC-CERL.
- Williams, S. E., L. P. Shoo, J. L. Isaac, A. A. Hoffmann, and G. Langham. 2008. Towards an integrated framework for assessing the vulnerability of species to climate change. *PLoS Biol* 6(12):e325. doi:10.1371/journal.pbio.0060325.
- Wilson, T. S., B. M. Sleeter, R. R. Sleeter, and C. E. Soulard. 2014. Land-use threats and protected areas: A scenario-based, landscape level approach. *Land* 3:362-389. doi:10.3390/land3020362.
- Young, B. E., N. S. Dubois, and E. L. Rowland. 2014. Using the Climate Change Vulnerability Index to inform adaptation planning: Lessons, innovations, and next steps. *Wildlife Society Bulletin* DOI:10.1002/wsb.478.
- Young, B. E., E. Byers, G. Hammerson, A. Frances, L. Oliver, and A. Treher. 2015. *Guidelines for Using the NatureServe Climate Change Vulnerability Index*. Release 3.0, Arlington VA, 63 pp.

Acronyms and Abbreviations

Term	Definition
ACSIM	Assistant Chief of Staff for Installation Management
ACUB	Army Compatible Use Buffer
AET:PET	Actual Evapotranspiration Potential Evapotranspiration
ANSI	American National Standards Institute
ASA(ALT)	Office of the Assistant Secretary of the Army for Acquisition, Logistics, and Technology
BRAC	Base Realignment and Closure
CCAR	Climate Change Adaptation Roadmap
CEERD	US Army Corps of Engineers, Engineer Research and Development Center
CERL	Construction Engineering Research Laboratory
CONUS	Continental United States
CTC	Combat Training Center
DoD	U.S. Department of Defense
DODI	Department of Defense Instruction
DODM	Department of Defense Manual
EO	Executive Order
ERDC	U.S. Army Engineer Research and Development Center
ERDC-CERL	Engineer Research and Development Center, Construction Engineering Research Laboratory
ESA	U.S. Endangered Species Act
ESRI	Environmental Systems Research Institute, Inc.
ICLUS	Integrated Climate and Land Use Scenarios
INRMP	Integrated Natural Resources Management Plans
ITAM	Integrated Training Area Management
LEAM	Land-use Evolution and impact Assessment Model
MTA	Military Training Area
MTA-L	Military Training Area-Light
MTC	Military Training Center
MTC-H	Military Training Center-Heavy
NSN	National Supply Number
OACSIM	Office of the Assistant Chief of Staff for Installation Management
OMB	Office of Management and Budget
PO	Post Office
REPI	Readiness and Environmental Protection Integration
RUG	Regional Urban Growth (model)
SAR	Same As Report
TS	Training Support
USEPA	U.S. Environmental Protection Agency

Term	Definition
USFWS	US Fish and Wildlife Service
USGS	U.S. Geological Survey

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE (DD-MM-YYYY) 07/12/2017		2. REPORT TYPE Final		3. DATES COVERED (From - To)	
4. TITLE AND SUBTITLE Climate Change Vulnerability of Army Installations Attributable to Listed and At-Risk Species				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT 622720A896	
6. AUTHOR(S) Matthew G. Hohmann, David K. Delaney, and Wade A. Wall				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER F94KF0	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. Army Engineer Research and Development Center (ERDC) Construction Engineering Research Laboratory (CERL) PO Box 9005, Champaign, IL 61826-9005				8. PERFORMING ORGANIZATION REPORT NUMBER ERDC/CERL TR-17-22	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of the Assistant Secretary of the Army for Acquisition, Logistics, and Technology 103 Army Pentagon Washington, DC 20310-0103				10. SPONSOR/MONITOR'S ACRONYM(S) OACSIM-ISE	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Climate change is an important emerging concern for the U.S. Department of Defense (DoD) and the Army. Key among the issues is climate change-driven increases in the number of species listed under the U.S. Endangered Species Act (ESA) and stress to Federally listed species already listed, because both pose additional management requirements and challenges, as well as potential restrictions on training land use. This work developed an approach for characterizing this component of installation climate change vulnerability that integrates multiple factors related to exposure, sensitivity, adaptive capacity, and number of listed and at-risk species. The approach was applied to Army installations in the Continental United States that have Integrated Training Area Management (ITAM) programs. Additionally, the assessment was used to rank installations across the Army based on the aggregate vulnerabilities of species. The approach to vulnerability assessment demonstrated here is suitable for evaluating whether climate change-driven impacts to listed and at-risk species is likely to affect installation resilience.					
15. SUBJECT TERMS Endangered species--United States, Military bases, Climatic changes, Climatic changes--Risk assessment, Environmental management					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT SAR	18. NUMBER OF PAGES 60	19a. NAME OF RESPONSIBLE PERSON
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified			19b. TELEPHONE NUMBER (include area code)